

Intended for  
**Maersk Oil**

Document type  
**Report**

Date  
**August, 2015**

# **MAERSK OIL ESIA-16 ENVIRONMENTAL AND SOCIAL IMPACT STATEMENT - DAN**

**MAERSK OIL ESIA-16  
ENVIRONMENTAL AND SOCIAL IMPACT STATEMENT -  
DAN**

Revision **3**  
Date **25/08/2015**  
Made by **DMM, MIBR, HEH**  
Checked by **CFJ, HEH**  
Approved by **CFJ**  
Description **Environmental and Social Impact Statement - DAN**

Ref ROGC-S-RA-000229

## CONTENTS

<b>1.</b>	<b>Introduction</b>	<b>1</b>
1.1	Background	1
<b>2.</b>	<b>Legal Background</b>	<b>3</b>
2.1	EU and national legislation	3
2.2	International conventions	4
2.3	Industry and national authority initiatives	5
<b>3.</b>	<b>Description of the project</b>	<b>6</b>
3.1	Existing facilities	6
3.2	Planned activities	12
3.3	Accidental events	18
3.4	Project alternatives	18
<b>4.</b>	<b>Methodology</b>	<b>19</b>
4.1	Rochdale envelope approach	19
4.2	Methodology for assessment of impacts	19
<b>5.</b>	<b>Environmental and social baseline</b>	<b>23</b>
5.1	Climate and air quality	23
5.2	Bathymetry	23
5.3	Hydrographic conditions	24
5.4	Water quality	25
5.5	Sediment type and quality	26
5.6	Plankton	27
5.7	Benthic communities	28
5.8	Fish	30
5.9	Marine Mammals	34
5.10	Seabirds	36
5.11	Cultural heritage	38
5.12	Protected areas	38
5.13	Marine spatial use	39
5.14	Fishery	40
5.15	Tourism	42
5.16	Employment	42
5.17	Tax revenue	43
5.18	Oil & gas dependency	43
<b>6.</b>	<b>Impact assessment: Planned activities</b>	<b>44</b>
6.1	Impact mechanisms and relevant receptors	44
6.2	Assessment of potential environmental impacts	47
6.3	Assessment of potential social impacts	72
6.4	Summary	76
<b>7.</b>	<b>Impact assessment: Accidental events</b>	<b>77</b>
7.1	Impact mechanisms and relevant receptors	77
7.2	Assessment of potential environmental impacts	88
7.3	Assessment of potential social impacts	95
7.4	Summary	98

<b>8.</b>	<b>Mitigating measures</b>	<b>99</b>
8.1	Mitigating for planned activities	99
8.2	Mitigating of accidental events	100
<b>9.</b>	<b>Environmental standards and procedures in Maersk Oil</b>	<b>101</b>
9.1	Environmental management system	101
9.2	Environmental and social impact in project maturation	101
9.3	Demonstration of BAT/BEP	101
9.4	Oil spill contingency plan	102
9.5	Ongoing monitoring	103
<b>10.</b>	<b>Natura 2000 screening</b>	<b>105</b>
10.1	Introduction	105
10.2	Designated species and habitats	105
10.3	Screening	106
10.4	Conclusion	107
<b>11.</b>	<b>Transboundary impacts</b>	<b>108</b>
11.1	Introduction	108
11.2	ESPOO convention	108
11.3	The DAN project	108
11.4	Identified impacts – planned activities	110
11.5	Identified impacts – accidental events	111
<b>12.</b>	<b>Lack of information and uncertainties</b>	<b>112</b>
12.1	Project description	112
12.2	Environmental and social baseline	112
12.3	Impact assessment	112
<b>13.</b>	<b>References</b>	<b>114</b>

## APPENDICES

### Appendix 1

Technical sections

## LIST OF FIGURES

Figure 1-1 Matrix for Maersk Oil ESIA-16, showing the 7 generic technical sections and the five ESIS.....	1
Figure 1-2 Project-specific environmental and social impact statement (ESIS) are prepared for the North Sea projects TYRA, HARALD, DAN, GORM and HALFDAN, respectively. ....	2
Figure 3-1 Overview of existing facilities at the DAN project (not to scale). ....	6
Figure 3-2 Dan Bravo .....	8
Figure 3-3 Dan F and Dan E .....	8
Figure 3-4 Kraka.....	9
Figure 3-5 Regnar subsea frame, as seen from rig during deployment.....	9
Figure 3-6 Simplified diagram of the process at Dan F .....	10
Figure 3-7 Sketch of fixed platform (option 1) .....	14
Figure 3-8 Maximum total expected production of oil, gas and water from the DAN project. Oil and water rate are provided as standard barrels per day, while the gas rate is provided as 1000 standard cubic feet of gas per day. The production forecast also accounts for the development projects. ....	15
Figure 3-9 Amount of oil discharged with produced water for the DAN project (based on minimum forecast of 8.5 mg/l and maximum forecast of 13 mg/l). Note that we have included .....	16
Figure 5-1 Bathymetry of the North Sea. Figure redrawn from Maersk Oil Atlas /3/.....	24
Figure 5-2 Left: General water circulation in the North Sea. The width of arrows is indicative of the magnitude of volume transport /10/. Right: Potential for hydrographic fronts in the North Sea /10//2/.....	25
Figure 5-3 Seabed sediments in the North Sea. Figure redrawn from North Sea Atlas /3/.....	26
Figure 5-4 Phytoplankton colour index (PCI) for the North Sea. Figure redrawn from North Sea Atlas /3/. ....	27
Figure 5-5 Assemblages of the benthic fauna in the North Sea. Figure redrawn from North Sea Atlas /3/. ....	30
Figure 5-6 Spawning grounds for cod and whiting in the North Sea. Figure redrawn from North Sea Atlas /3/. ....	33
Figure 5-7 Distribution of harbour porpoise in the North Sea. Figure redrawn from North Sea Atlas /3/. ....	35
Figure 5-8 Protected areas. Figure redrawn from North Sea Atlas /3/. ....	38
Figure 5-9 Ship traffic and infrastructure in 2012. Figure redrawn from North Sea Atlas /3/. Ship traffic is based on all ships fitted with AIS system i.e. ships of more than 300 gross tonnage engaged on international voyages, and cargo ships of more than 500 gross tonnage not engaged on international voyages and all passenger ships irrespective of size. Missing data in the middle of the North Sea is due to poor AIS receiving coverage and not lack of ships. Germany does not participate in the North Sea AIS data sharing program. ...	40
Figure 5-10 Employment per sector in Denmark in 2013 /39/.....	42
Figure 6-1 Sedimentation of discharged water based drilling mud modelled for a typical well /1/.....	54
Figure 6-2 Sedimentation of water based drill cuttings modelled for a typical well /1/. ....	54
Figure 7-1 Accidental oil, diesel and chemical spills from Maersk Oil platforms in the North Sea. ....	78
Figure 7-2 Probability that a surface a 1 km cell could be impacted by oil in case of full pipeline rupture /152/. ....	80

Figure 7-3 Location of two Maersk Oil production wells Svend and Halfdan (and a pipeline midpoint), for which oil spill modelling has been undertaken. Halfdan is located close to the DAN project, and modelling for Halfdan are considered representative to the DAN project. The HALFDAN and DAN platforms are located within 7 km from each other ..... 82

Figure 7-4 Probability that a surface cell could be impacted in a surface blowout at Halfdan well /152/..... 84

Figure 7-5 Probability that a water column cell could be impacted in a surface blowout at Halfdan well /152/..... 85

Figure 7-6 Probability that a shoreline cell could be impacted in a surface blowout at Halfdan well /152/..... 86

Figure 7-7 Maximum time-averaged total oil concentration for a surface blowout at Halfdan well /152/..... 87

Figure 9-1 Illustration of best available technique ..... 102

Figure 9-2 Acoustic monitoring of marine mammals (Photo: Aarhus University, DCE). ..... 104

Figure 10-1 Natura 2000 sites in the North Sea. .... 105

Figure 11-1 Maersk Oil North Sea projects TYRA, HARALD, DAN, GORM and HALFDAN. .... 109

## LIST OF ABBREVIATIONS

ALARP	As low as reasonably practicable
API	American Petroleum Institute gravity. An industry standard used to determine and classify of oil according to their density
BAT	Best available technique
BEP	Best environmental practice
BOPD	Barrels of oil per day
BWPD	Barrels of water per day
CO <sub>2</sub>	Carbon dioxide
DEA	Danish Energy Agency [Energistyrelsen]
DEPA	Danish Environmental Protection Agency [Miljøstyrelsen]
DNA	Danish Nature Agency [Naturstyrelsen]
DUC	Danish Underground Consortium, a joint venture with A. P. Møller – Mærsk, Shell, Chevron and the Danish North Sea Fund
EIA	Environmental impact assessment
EIF	Environmental impact factor
ESIA	Environmental and social impact assessment
ESIS	Environmental and social impact statement
FTEE	Full time employee equivalent
GBS	Gravity-based structure
Hz	Hertz
ITOPF	International tanker owners pollution federation
KSCF	1000 standard cubic foot of gas
MBES	Multibeam echo sounder
MMO	Marine mammal observer
MMSCFD	Million standard cubic feet of gas per day
NM VOC	Non methane volatile organic compounds.
NORM	Naturally occurring radioactive material
NO	Nitric oxide
NO <sub>2</sub>	Nitrogen dioxide
NO <sub>x</sub>	NO <sub>x</sub> is a generic term for mono-nitrogen oxides NO and NO <sub>2</sub> (nitric oxide and nitrogen dioxide)
OSPAR	Oslo-Paris convention for the protection of the marine environment of the North-East Atlantic
PAM	Passive acoustic monitoring
PEC	Predicted environmental concentration
PNEC	Predicted no-effect concentration based on ecotoxicity data
PLONOR	Pose little of no risk
PM <sub>2.5</sub>	Particulate Matter less than 2.5 microns in diameter
PPM	Parts per million
RBA	Risk-based approach
ROV	Remote operated vehicle
SO <sub>2</sub>	Sulphur dioxide
SO <sub>x</sub>	Refers to all sulphur oxides, the two major ones being sulphur dioxide and sulphur trioxide
SSS	Side scan sonar
STB	Standard barrels (of oil)


# 1. INTRODUCTION

## 1.1 Background

In connection with ongoing and future oil and gas exploration, production and decommissioning activities by Maersk Oil in the Danish North Sea, an environmental and social impact assessment (ESIA-16) is prepared. The overall aim of the ESIA-16 is to identify and assess the impact of the Maersk Oil activities on environmental and social receptors.


ESIA-16 shall replace the EIA conducted in 2010 /1/ which is valid for the period 1<sup>st</sup> January 2010 to 31<sup>st</sup> December 2015. The ESIA-16 covers the remaining lifetime of the ongoing projects, and the whole life time from exploration to decommissioning for planned projects.

The ESIA-16 consists of five independent project-specific environmental and social impact statements (ESIS) for TYRA, HARALD, DAN, GORM and HALFDAN including seven generic technical sections that describe the typical activities (seismic, pipelines & structures, production, drilling, well stimulation, transport and decommissioning; provided in appendix 1) in ongoing and planned Maersk Oil projects. Drilling of stand alone exploration wells and replacement of existing pipelines are not included in ESIA-16 and are screened separately in accordance with Order 632 dated 11/06/2012.



**Figure 1-1 Matrix for Maersk Oil ESIA-16, showing the 7 generic technical sections and the five ESIS.**

The environmental and social impact statement for the DAN project covers the activities related to existing and planned projects for Dan F and its satellite platforms Dan B, Dan E, Kraka and the subsea wellhead Regnar. The platforms are located in the North Sea about 220 km from the west coast of Jutland, Denmark (Figure 1-2).



**Figure 1-2 Project-specific environmental and social impact statement (ESIS) are prepared for the North Sea projects TYRA, HARALD, DAN, GORM and HALFDAN, respectively.**

## 2. LEGAL BACKGROUND

### 2.1 EU and national legislation

#### 2.1.1 Environmental impact assessment directive (EIA directive)

The directive on the assessment of the effects of certain public and private projects on the environment (directive 85/337/EEC), as amended by directives 7/11/EC, 2003/35/EC and 2009/31/EC, requires an assessment of the environmental impacts prior to consent being granted. For offshore exploration and recovery of hydrocarbons this directive is implemented in Denmark as executive order 632 dated 11/06/2012. The order is under revision to implement amendments following directive 2014/52.

This ESIA-16 has been prepared in accordance with order 632 dated 11/06/2012 on environmental impact assessment (EIA) and appropriate assessment (AA) for the hydrocarbon activities [Bekendtgørelse om VVM, konsekvensvurdering vedrørende internationale naturbeskyttelsesområder og beskyttelse af visse arter ved efterforskning og indvinding af kulbrinter, lagring i undergrunden, rørledninger, m.v. offshore].

- Transboundary significant adverse impacts are addressed (section 8), in accordance with article 8 and the ESPOO convention.
- Protection of certain species mentioned in the directive article 12 (section 6).
- A Natura 2000 screening is presented in this ESIS (section 10), in accordance with article 9 and 10.

The ESIS and its non-technical summary shall be made available for public consultation on the web page of the Danish Energy Agency. Public consultation shall be for a period of at least 8 weeks, in accordance with article 6.

#### 2.1.2 Protection of the marine environment

The consolidation act 963 dated 03/07/2013 on protection of the marine environment aims to protect the environment and ensure sustainable development.

The consolidation act and associated orders regulate e.g. discharges and emissions from platforms. Relevant orders include: Order 394 dated 17/07/1984 on discharge from some marine constructions, order 9840 dated 12/04/2007 on prevention on air pollution from ships, and order 909 dated 10/07/2015 on contingency plans.

#### 2.1.3 Natura 2000 (Habitats and Bird protection directive)

The "Natura 2000" network is the largest ecological network in the world, ensuring biodiversity by conserving natural habitats and wild fauna and flora in the territory of the EU. The network comprises special areas of conservation designated under the directive on the conservation of natural habitats and of wild fauna and flora (Habitats Directive, Directive 1992/43/EEC). Furthermore, Natura 2000 also includes special protection areas classified pursuant to the Birds Directive (Directive 2009/147/EC) and the Ramsar convention. The directives have been transposed to Danish legislation through a number of orders (or regulatory instruments).

The Natura 2000 protection is included in the order 632 dated 11/06/2012 (section 2.1.1).

#### 2.1.4 National emissions ceiling directive

The national emission ceiling directive (directive 2001/81/EC) sets upper limits for each Member State for the total emissions of the four pollutants nitrogen oxide  $\text{NO}_x$ , volatile organic compound (VOC), ammonia ( $\text{NH}_3$ ) and sulphur dioxide ( $\text{SO}_2$ ). The directive is under revision to include Particulate Matter less than 2.5 microns in diameter ( $\text{PM}_{2.5}$ ). The directive has been implemented by order 1325 dated 21/12/2011 on national emissions ceilings.

#### 2.1.5 Marine strategy framework directive

The marine strategy framework directive (Directive 2008/56/EC) aims to achieve "good environmental status" of the EU marine waters by 2020. The directive has been implemented in Denmark by the act on marine strategy (act 522 dated 26/05/2010). A marine strategy has been developed by the Danish Nature Agency with a detailed assessment of the state of the environment, with a definition of "good environmental status" at regional level and the establishment of environmental targets and monitoring programs ([www.nst.dk](http://www.nst.dk)).

#### 2.1.6 Industrial emissions directive

The industrial emissions directive (directive 2010/75/EU) is about minimising pollution from various industrial sources. The directive addresses integrated pollution prevention and control based on best available technique (BAT). The directive has been implemented by the consolidation act 879 dated 26/06/2010 on protection of the environment and with respect to offshore, order 1449 dated 20/12/2012.

#### 2.1.7 Emission allowances

The European Union Emissions Trading Scheme was launched in 2005 to combat climate change and is a major pillar of EU climate policy. Under the 'cap and trade' principle, a cap is set on the total amount of greenhouse gases that can be emitted by all participating installations. The trading scheme is implemented by act 1095 dated 28/11/2012 on CO<sub>2</sub> emission allowances.

#### 2.1.8 Safety directive of offshore oil and gas operations

The directive 2013/30/EU on safety of offshore oil and gas operations aims to ensure that best safety practices are implemented across all active offshore regions in Europe. The directive has recently been implemented by act 1499 dated 23/12/2014 on offshore safety.

## 2.2 International conventions

#### 2.2.1 Espoo convention

The convention on environmental impact assessment in a transboundary context (Espoo Convention) entered into force in 1991. The convention sets out the obligations of Parties to assess the environmental impact of certain activities at an early stage of planning. It also lays down the general obligation of States to notify and consult each other on all major projects under consideration that are likely to have a significant adverse environmental impact across national boundaries.

The Espoo convention is implemented in the EIA Directive. In Denmark, the Ministry of Environment administrate the Espoo Convention rules and is the responsible authority for the process of exchanging relevant information from the project owner to the potentially affected countries and possible comments from those countries in connection with the Espoo Consultation Process.

#### 2.2.2 Convention on the prevention of marine pollution by dumping of wastes and other matter

International maritime organization (IMO) convention on the prevention of marine pollution by dumping of wastes and other matter (London Convention) has been in force since 1975. Its objective is to promote the effective control of all sources of marine pollution and to take all practicable steps to prevent pollution of the sea by dumping of wastes and other matter.

#### 2.2.3 Convention for the control and management of ships' ballast water and sediments

The convention for the control and management of ships' ballast water and sediments (ballast water management convention) was adopted in 2004. The convention aims to prevent the spread of harmful aquatic organisms from one region to another, by establishing standards and procedures for the management and control of ships' ballast water and sediments.

#### 2.2.4 Ramsar convention

The Ramsar convention aims at the conservation and wise use of all wetlands through local and national actions and international cooperation, as a contribution towards achieving sustainable development throughout the world.

#### 2.2.5 The convention for the protection of the marine environment of the North-East Atlantic

The convention for the protection of the marine Environment of the North-East Atlantic (the 'OSPAR Convention') entered into force in 1998. Contained within the OSPAR Convention are a series of Annexes which focus on prevention and control of pollution from different types of activities. OSPAR has a focus on application of the precautionary principle, and on use of best available technique (BAT), best environmental practice (BEP) and clean technologies.

A number of strategies and recommendations from OSPAR are relevant to the DAN project, most notably:

- Annual OSPAR report on discharges, spills and emissions from offshore oil and gas installations.
- Reduction in the total quantity of oil in produced water discharged and the performance standard of dispersed oil of 30 mg/l (OSPAR Recommendation 2001/1).
- Harmonised mandatory control system for the use and reduction of the discharge of Offshore chemicals (OSPAR decision 2005/1).
- List of substances/preparations used and discharged offshore which are considered to pose little or no risk to the environment (PLONOR) (OSPAR decision 2005/1).
- To phase out, by 1 January 2017, the discharge of offshore chemicals that are, or which contain substances, identified as candidates for substitution, except for those chemicals where, despite considerable efforts, it can be demonstrated that this is not feasible due to technical or safety reasons (OSPAR Recommendation 2006/3).
- Risk based approach to assessment of discharged produced water (OSPAR recommendation 20012/5).
- Decision 98/3 on the disposal of disused offshore installations.

#### 2.2.6 Convention on access to information, public participation in decision-making and access to justice in environmental matters

The UNECE convention on access to information, public participation in decision-making and access to justice in environmental matters (Aarhus convention) was adopted in 1998. The convention is about government accountability, transparency and responsiveness. The Aarhus convention grants the public rights and imposes on parties and public authorities obligations regarding access to information and public participation. The Aarhus convention is among others implemented in Denmark by the Subsoil Act 960 dated 13<sup>th</sup> September 2013.

### 2.3 Industry and national authority initiatives

#### 2.3.1 Offshore action plan

An offshore action plan was implemented by the Danish Environmental Protection Agency and the Danish operators in 2005 in order to reduce the discharge of chemicals and oil in produced water. A revised action plan for 2008-2010 was implemented to reduce emissions to air and further reduce discharges.

#### 2.3.2 Action plan on energy efficiency

An action plan on energy efficiency was implemented by the Danish Energy Agency and the Danish oil and gas operators for 2008-2011 and 2012-2014 to improve energy efficiency for the oil and gas industry. More specifically, the action plan included measures on energy management and initiatives to reduce flaring and energy consumption.

### 3. DESCRIPTION OF THE PROJECT

The project description for the DAN project is based on site specific input from Maersk Oil and the technical sections (appendix 1). The DAN project refers to the existing and planned activities for Dan F and its satellite Dan B, Dan E, Kraka and the subsea wellhead Regnar. The DAN project (capital letters) refers to the project, while Dan B, Dan E or Dan F refer to the intallations/platforms.

#### 3.1 Existing facilities

##### 3.1.1 Overview

The DAN project refers to the existing and planned activities for Dan F and its satellite Dan B, Dan E, Kraka and the subsea wellhead Regnar. The production facilities are connected by subsea pipelines, through which oil, gas and water are transported. Pipelines departing from the Dan F, Dan B, Dan E and Kraka and the subsea well head Regnar are considered part of the DAN project.

An overview of the existing pipelines and structures for the DAN project is provided in Figure 3-1.




Figure 3-1 Overview of existing facilities at the DAN project (not to scale).

### 3.1.2 Pipelines and structures

#### 3.1.2.1 Dan B, Dan E and Dan F

Dan B (Figure 3-2), Dan E and Dan F (Figure 3-3) are located in the South-Western part of the Danish sector of the North Sea, approximately 210 km west of Esbjerg. The water depth is 41-42 m. Dan F, Dan B and Dan E are located approximately 1-2 km from each other.

Dan B (Figure 3-2) comprises a processing and accommodation platform (Dan BB), two wellhead platforms (Dan BA and Dan BD) and a flare platform (Dan BC). Today, Dan B functions as a manned satellite to Dan F and holds no processing equipment as the entire production is routed to Dan F. Topsides modifications are planned at Dan B to change the Dan B facility into an unmanned platform in the next few years.

Dan E is a wellhead platform.

The Dan F installation consists of seven platforms, which are connected by bridges:

- Dan FA and FB platforms are wellhead platforms as well as risers for the subsea pipelines to Gorm, Tyra E, Dan E, Kraka and Regnar. Seawater treatment for water injection is installed on the FE bridge module.
- Dan FC is the main platform with accommodation, utility and life support systems as well as support systems for platform operations and production equipment for treatment of oil and produced water and export of oil and gas.
- Dan FD is a STAR platform for support of the flare tower for flaring of produced gas if required.
- Dan FE is a wellhead platform with fire pump and water injection facilities for treatment and injection of seawater.
- Dan FF is a wellhead, process and utility platform. Besides wells, the platform holds separation, compression and dehydration facilities, risers to/from Dan D, Halfdan and Kraka and utilities such as power generation, water injection etc.
- Dan FG is a process and utility platform which holds equipment for separation, gas compression, dehydration and water injection, flare tower, fire water pumps and other utilities as well as provision for a future module on top of the topside.



**Figure 3-2 Dan Bravo**



**Figure 3-3 Dan F and Dan E**

Dan F is the primary processing platform for the entire oil production from the DAN project. The processed gas is sent to Tyra East, while the crude oil is transported to Fredericia via the Gorm E riser platform. The majority of the produced water is discharged to sea at Dan F.

Continuous control and monitoring of the satellite platform Kraka and the subsea wellhead Regnar is carried out from Dan F.

### 3.1.2.2 Kraka

Kraka is situated approximately 9 km south of Dan F. The water depth at Kraka is 44 m.

The Kraka installation comprises an unmanned wellhead STAR platform without a helideck. There are no processing facilities at Kraka, and the production is transported to Dan F for processing.




**Figure 3-4 Kraka**

### 3.1.2.3 Regnar

Regnar is situated approximately 13 km southwest of Dan F. The water depth at Regnar is 43 meters.

Regnar is a subsea-completed well. The hydrocarbons produced are conveyed by pipeline in multiphase flow to Dan F for processing and export ashore. The well is remotely monitored and controlled from the Dan FC platform. Regnar is currently not producing.



**Figure 3-5 Regnar subsea frame, as seen from rig during deployment.**

3.1.2.4 Pipelines

The production facilities are connected by subsea pipelines, through which oil, gas and water are transported. Pipelines are trenched to a depth of 2 m or covered by rocks where above the seafloor. An overview of the existing pipelines and their content is provided in Figure 3-1.

3.1.3 Wells

The DAN project currently has a total of 123 existing wells: 93 at Dan F, 16 at Dan B, 6 at Dan E, 7 at Kraka and 1 at Regnar. Three well slots are available for drilling at Dan F.

3.1.4 Processing capabilities


The processing capability at the DAN project (Dan F) is provided in Table 3-1. The facility is designed for continuous operation 24 hours a day. Maintenance is generally planned, so only part of the facility is shut down, thus only reducing the production. The whole facility is only shut down in case of major emergencies or maintenance operations.

**Table 3-1 Processing capacity at the DAN project (Dan F).**

Process	Unit	Dan F
Primary separation	BOPD	410,000
Gas compression	MMSCFD	390
Produced water treatment	BWPD	340,000
HP water injection	BWPD	777,000
Oil export	BOPD	261,000

There are 3 main processes:

- Separation and stabilisation process,
- Gas compression and dehydration process,
- Water (seawater) injection process.



**Figure 3-6 Simplified diagram of the process at Dan F**

The energy supply to the Dan facility consists of self-produced natural gas from the Dan field, imported natural gas from Tyra and diesel supplied by ship.

Natural gas is used as fuel gas in gas turbines operating as drives for e.g. power generators, gas compressors and high-pressure water injection pumps.

Diesel is used in dual-fuel gas turbines, for cranes and for emergency equipment such as fire pumps.

Flaring of gas at compressor inlet/outlet might be required for short periods of time in relation to planned and controlled process operations (e.g. start up) and for safety reason in relation with unforeseen process upsets which causes overpressure of process equipment and emergency depressurization of platform equipment.

#### 3.1.5 Waste

Maersk Oil transports all waste from its Danish North Sea facilities to shore where it is recycled, incinerated or landfilled in accordance with current legislation. The last five years, an average of about 10,000 tons of waste were collected and brought onshore from all Maersk Oil facilities. In the last five years, about 99 % of the waste was recycled or incinerated. Landfilled waste is partly made up of sandblasting materials. Since 2014, most of the sand is being reused for roads construction and other building materials leading to a significant reduction in the amount of landfilled waste.

Decommissioning of the DAN facilities is expected to generate up to 43,000 tons of waste which will be brought onshore and treated accordingly. The main source of waste is expected to be from the carbon steel from the jacket and the topside facilities.

#### 3.1.6 NORM

Normally occurring radioactive material (NORM) such as sand, scale, cleanup materials from tubing, valves or pipes are collected and brought onshore, where they are treated to remove traces of hydrocarbons or scaleformation. After treatment, the NORM is securely stored. The average quantity of NORM stored in 2013-2014 was approximately 70 tons. The quantity of NORM is expected to increase as fields are maturing and Maersk Oil is currently evaluating the best options for handling of NORM waste.

#### 3.1.7 Discharges

A number of discharges are expected as part of the planned activities, including drilling mud and cuttings, produced water and cooling water. These are described in appendix 1.

In addition, main liquid effluents generated by the vessels and platforms will comprise:

- Greywater (water from culinary activities, shower and laundry facilities, deck drains and other non-oily waste water drains (excluding sewage))
- Treated blackwater (sewage)
- Drainage water
- Service water / vessel engine cooling water.

All discharges will comply with requirements set out in the International Convention for the Prevention of Pollution from Ships, 1973, as modified by the Protocol of 1978 (MARPOL 73/78) and its annexes.

## 3.2 Planned activities

Here, the planned activities for the DAN project are presented with reference to the seven technical sections (appendix 1).

### 3.2.1 Seismic

Seismic surveys are performed to provide information about the subsurface geological structure to identify the location and volume of potential hydrocarbon reserves, and to ensure that seabed and subsurface conditions are suitable for planned activities (e.g. drilling and construction of production facilities).

For the DAN project, several types of seismic data acquisition may be carried out:

- 4D seismic surveys are 3D seismic surveys repeated over a period of time, and can take several months to complete. A 4D seismic covering an area of a few hundred km<sup>2</sup> is planned for 2016 or 2017, and is expected to be repeated every 4 years.
- Drilling hazard site surveys (one per year expected) may include 2D high resolution multi-channel and single-channel seismic, side scan sonar, single and multi-beam echo-sounder, seabed coring and magnetometer. Typical duration of such a survey is 1 week covering an area of 1x1 km.
- Borehole seismic surveys (one per year expected) are conducted with a number of geophones that are lowered into a wellbore to record geological data around the well hole. The duration is usually one to two days.

### 3.2.2 Pipelines and structures

Regular maintenance of the existing pipelines and structures at the DAN project will be undertaken including external visual inspections by remotely operated vehicles (ROVs) and an internal inspection/cleaning of pipelines (pigging). If inspection surveys reveals that the replacement of existing pipelines is necessary, a separate project and environmental screening will be carried out.

For the DAN project, two new development projects are planned: Dan Area redevelopment and the expansion of Dan F accommodation capacity.

#### 3.2.2.1 Dan Area redevelopment

The development project includes the construction of up to 4 unmanned platforms (SLIC platforms): 1 near the existing Kraka platform and 3 in the Dan F area. The new platforms in Dan F area will likely be located approximately 4-7 km from Dan F.

The new platforms are expected to be SLIC platforms with 10 slots platforms in order to allow future development and standardization. SLIC platforms are a minimum facilities concept, where only essential equipment needed for oil production is included. The platforms will be unmanned and designed to reduce the number of maintenance operations. The platforms are designed to be manned using a "walk to work" system or by fast rescue craft in limited circumstances. An offshore support rig will be used for well interventions.

Maersk Oil topside and jacket satellite facilities are mostly constituted of carbon steel. The new satellite platforms may be constructed following the tripod concept with jacket size ranging from 950-1500 t incl. foundation, and topside satellite facilities (SLIC concept) ranging from 70-400 t.

The 4 platforms will be 10 well slot platforms for oil/gas/water production and water injection, i.e. a total of up to 40 wells. Initially, 4 wells are expected to be drilled to develop the DAN B-North area and 5 wells are expected to be drilled to develop the DAN B-South East. Finally, 7 wells are expected to be drilled to develop the DAN B-South and the Kraka area. The remaining slots use and timing has not been planned but may be further expected in case of successful development.

The new platform at Kraka maybe built for disposal of water production/injection - a novelty in this field. A test for introducing water injection at Kraka is currently being planned with possible start up date in 2017.

The trial will consist of injecting water into a selected well at a flowrate of 5,000 barrel of water per day and measuring the change in the production rates over a 1-2 year period.

Three possible ways to do the trial are currently under investigation.

1. Temporary water injection plant on Kraka. This setup includes lift pumps, filtering, chemical injection (oxygen scavenger, biocide, hypochloride) and high pressure injection pumps located on Kraka. Minor discharge of few hundred litres per day at start up may be expected.
2. Injection from boat. In this options a ship with water filtering, chemical injection and pumping capability will be position next to Kraka and will supply Kraka with injection water through a flexible hose.
3. Injection water supply from Dan F through a 9 km temporary pipeline to Kraka. The pipeline can either be a flexible hose of a metal pipe. The pipeline will not be buried and the area will be secured from boat anchoring or trawling. Information to the other seauser will be provided through relevant authorities. In addition, Maerks Oil will commissioned a guard ship that will be positioned to alert other approaching sea-user (e.g. fishing boats).

The production is expected to be transported via two sets of pipelines (North branch and South branch) to a common pipeline to Dan F for processing (oil, gas and produced water) and continued transport onshore. The platforms will be remotely controlled from Dan F. Dan F will handle the expected additional production with its existing processing facilities. Only minor topside adjustments (pig receiver, riser) at Dan F are expected.


It is expected that a 16" multiphase production pipeline, a 12" water injection and a 12" gaslift pipeline and a smaller umbilical for chemicals transport, power and signal would be required.. Maersk Oil uses steel pipelines that weigh between 42 MT (6" linepipe) and 320 MT (24" linepipe) per km depending on the diameter size of the pipeline.

#### 3.2.2.2 Expansion of Dan F accommodation capacity

The current accommodation capacity on Dan FC is not sufficient to support the planned offshore activities on Dan. Therefore Maersk Oil is currently identifying the opportunities to increase and improve the accommodation on Dan F. The objective is to provide a cost efficient and up-to-date solution, which also matches the expected long term Maersk Oil infrastructure development plan for the southern fields.

The following three options are currently being considered for realisation earliest in 2018-2019:

- Option 1: build and install a new 200+ single bed cabin fixed accommodation platform sitting on a 4 legged jackets and linked to the Dan FC by a bridge. The existing Dan FC accommodation module would be sealed off. The top side and jacket would be designed for future reuse in another location .
- Option 2: a 200+ single bed cabin mobile accommodation rig on a long term contract or owned by Maersk. The existing Dan FC accommodation module would be sealed off.
- Option 3: a 100+ single bed cabin accommodation rig on a long term contract or owned by Maersk and carry out "cabin split" and refurbishment of the existing Dan FC accommodation module.




**Figure 3-7 Sketch of fixed platform (option 1)**

### 3.2.3 Production

Production was initiated at Dan B in 1972, then later at Kraka (1991) and Regnar (1993). The total hydrocarbon production for the DAN project peaked around 2000 and is now on a natural decline. This reflects the fact that the fields are in a relatively mature stage in the production cycle. The new development at the DAN project aim to increase the economic life of the fields. The maximum production forecasts of oil and gas from the DAN project are shown in Figure 3-8.


Throughout their productive life, most oil wells produce oil, gas, and water. Initially, the mixture coming from the reservoir may be mostly hydrocarbons but over time, the proportion of water increases and the fluid processing becomes more challenging. Processing is required to separate the fluids produced from the reservoirs.

The maximum production forecasts of produced water from the DAN project is shown in Figure 3-8. There is currently no injection of produced water at the DAN project. These solutions are currently being considered (see section 3.2.2).



**Figure 3-8 Maximum total expected production of oil, gas and water from the DAN project. Oil and water rate are provided as standard barrels per day, while the gas rate is provided as 1000 standard cubic feet of gas per day. The production forecast also accounts for the development projects.**

Maersk Oil uses production chemicals (e.g. H<sub>2</sub>S scavenger, biocides) to optimise the processing of the produced fluids. The inventory of Maersk Oil main chemicals, their general use and partitioning in water/oil phase is presented in appendix 1. A fraction of the oil and chemicals is contained in the treated produced water which is discharged. Discharges of produced water to sea is permitted only after authorisation from the DEPA.



**Figure 3-9 Amount of oil discharged with produced water for the DAN project (based on minimum forecast of 8.5 mg/l and maximum forecast of 13 mg/l). Note that we have included**

The nature, type and quantities of chemicals that are used in production and discharged to sea are expected to be adjusted to follow changes in production and technical development. In 2013-2014, about 7,050 tons of chemicals were used for production at the DAN project and about 7,200 tons of chemicals were discharged to sea at the Dan F platform (note that some chemicals are received from nearby platforms). As a general rule, the amount of chemical used, is somewhat related to the volume of produced water. In the future, Maersk Oil will continue to reduce the risk of impact of the discharges on the marine environment, by reducing of the volume of discharged chemicals, improving of the treatment processes or selecting alternative chemicals (see mitigating measures in section 8).

The expected volume of discharged oil is shown in Figure 3-9. The DAN project contributes to the total amount of oil in produced water discharges to sea. The estimates of oil discharges (average and maximum, Figure 3-9) are based on produced water discharge forecasts and historical oil in water figures at Dan. Oil content in produced water is regulated by OSPAR and the total amount of oil discharged to sea is limited by the DEPA.

Maersk Oil has flowmeters measuring the volume of discharged produced water, and water samples are regularly obtained for analysis of oil and chemical content. The nature, type and quantities chemical used and chemicals and oil discharged to sea are reported to the DEPA.

3.2.4 Drilling

Drilling of wells is necessary for extracting oil and gas resources. Wells are used for transporting the fluid (a mixture of oil, gas, water, sand and non-hydrocarbon gasses) from the geological reservoir to Maersk Oil installations, where fluid processing takes place. Wells are also used for injection of water (seawater or produced water) or gas to increase reservoir pressure and enhance the oil and gas recovery rate.

For the DAN project, drilling can take place in existing well slots. There are a total of 3 free well slots at Dan F. Maersk Oil has not decided whether these free well slots will be drilled. In addition, new development may create up to 40 new wells. Typical well types are presented in appendix 1. It has not been decided which type of well will be applicable for the DAN project. Drilling is performed from a drilling rig, which is placed on the seabed (with an expected area of a few hundred m<sup>2</sup>). A new well will typically take up to 150 days to drill. Different types of drilling mud will be used based on the well and reservoir properties. Water-based mud and cuttings will be discharged to the sea, whereas oil-based mud and cuttings will be brought onshore to be dried

and incinerated. Discharges to sea is permitted only after authorisation from the Environmental Protection Agency. Water-based drilling mud and drill cuttings may contain traces of oil from the reservoir sections. The oil content in the water-based drilling mud and drill cuttings is monitored regularly to ensure it does not exceed 2%, on average. It is estimated that on average 7 tons of oil per 1,000 m reservoir section can be discharged to sea corresponding to a maximum discharge of 28.8 tons of oil per well (type 2 and 4 with a 5,000 m reservoir section).

For the DAN project, 14 wells may be subjected to slot recovery or re-drill. When production from an existing well is no longer profitable, the slots may be re-used to access additional resources. This can be done in two ways: Slot recovery or re-drill. For slot recovery, the redundant well is abandoned and a new well is drilled and completed from a new conductor. For re-drill, sections of the redundant well are re-used. The nature and type of discharges and emissions related to slot recovery or re-drill operations will be less or equivalent to that of a well abandonment and the drilling of a well.

### 3.2.5 Well stimulation

The purpose of well stimulation is to improve the contact between the well and reservoir, thereby facilitating hydrocarbon extraction (for a production well) or water injection (for an injection well). Well testing is performed to evaluate the production potential of a well after stimulation.

At the DAN project, the new wells (up to 57 in total) may be subjected to matrix acid stimulation or acid fracturing. The existing wells at the DAN project may be subjected to matrix acid stimulations (in total up to 2 per year). Use and discharge (e.g. drilling and maintenance) of chemicals are presented in appendix 1. Discharges to sea is permitted only after authorisation from the DEPA.

### 3.2.6 Transport

Personnel and cargo are transported daily to support Maersk Oil's production and drilling operations via helicopters, supply vessels and survey vessels. Standby vessel may be employed in connection with drilling and tasks requiring work over the side of the installation.

Dan F and Dan B is manned at all time, while Kraka, Dan E and Regnar are unmanned.

### 3.2.7 Decommissioning

Decommissioning will be done in accordance with technical capabilities, legislation, industry experience, international conventions and the legal framework at the time of decommissioning. Decommissioning will be planned in accordance with the OSPAR decision 98/3 on the disposal of disused offshore installations.

- Wells will be permanently plugged towards the reservoir and the casing above the seabed will be removed.
- The well head, x-mas tree and protection frame will be removed and brought to shore for dismantling. Hydrocarbons and waste will be sent to shore for disposal.
- Buried pipelines will be cleaned, and left in situ, filled with seawater.

At the DAN project, one decommissioning project is planned for the sub sea well head Regnar, which is expected to be decommissioned within the next three years. However, the 2 pipelines 6" and 2.5" that connect Regnar to Dan F will be preserved for possible future tie-in.

The whole pipeline and well head system will be rinsed with deairedated seawater before the well is abandoned. The water will be treated on Dan F facilities to remove hydrocarbon traces before it is discharged to sea. The well head will be removed and the pipelines interconnected before they are back-filled with inhibited seawater to avoid corrosion damage until re-use.

Pipeline work will be performed by divers, whereas the well abandonment and wellhead removal will be assisted by a drilling rig. The duration of the operation is estimated to require 7 days of diving support vessels for pipelines diver work and 7 days for the rig-assisted work. The safety zone around the pipelines will be maintained according to the Danish Maritime Authority Order No. 939 of 27 November 1992.

### 3.3 Accidental events

The accidental events, considered here, are accidents that could take place during exploration, production and decommissioning activities at the DAN project that can lead to environmental or social impacts.

Accidents might occur as a result of a loss of primary containment event (oil, gas or chemical). Generally, the sequence of events leading to loss of primary containment are complex and a large number of scenarios can be envisioned (e.g. /136//137/).

The scenarios associated with Maersk Oil activities at the DAN project that can lead to major accidents with a risk of major significant impacts are listed in the technical sections and include vessels collisions, pipeline rupture due to corrosion, erosion or impact, well blow out, impact on major platform equipment. Small operational accidental spills of oil or chemical or gas release could also occur.

### 3.4 Project alternatives

Maersk Oil has considered several alternatives for planned activities. The alternatives have been evaluated based on technical, financial, environmental and safety parameters.

#### 3.4.1 0 alternative

The 0 alternative (zero alternative) is a projection of the anticipated future development without project realization, and describes the potential result if nothing is done. For the DAN project, this would mean that the production would cease.

The offshore oil and gas production is important to Danish economy. Thousands of people are employed in the offshore industry, and tax revenue to the state of Denmark is significant. The state's total revenue is estimated to range from DKK 20 to DKK 25 billion per year for the period from 2014 to 2018.

The Danish government has set a target of 30 % of the Danish energy use is provided from renewable energy by 2020. As part of a long-term Danish energy strategy, the oil and gas production is considered instrumental in maintaining high security of supply. Denmark is expected to continue being a net exporter of natural gas up to and including 2025 and Maersk Oil has license to operate until 2042 /35/.

If no production is undertaken by Maersk Oil for the DAN project area in the North Sea, there will be no contribution from the DAN project to the Danish economy or security of supply.

#### 3.4.2 Technical alternatives

Technical alternatives for seismic, pipelines & structures, production, drilling, well stimulation, transport and decommissioning are presented in appendix 1.

## 4. METHODOLOGY

The ESIS is based on the 2014 North Sea Atlas, technical reports, EIAs, peer-reviewed scientific literature, Maersk monitoring reports and industry reports.

### 4.1 Rochdale envelope approach

The adoption of the Rochdale Envelope approach allows meaningful ESIA to take place by defining a 'realistic worst case' scenario that decision makers can consider in determining the acceptability, or otherwise, of the environmental impacts of a project.

The Rochdale Envelope Approach allows a project description to be broadly defined. The project can be described by a series of maximum extents – the 'realistic worst case' scenario. The detailed design of the scheme can then vary within this 'envelope' without invalidating the corresponding ESIA.

Where a range is provided, e.g. amounts of produced water or volume of drilling mud, the most detrimental is assessed in each case. For example, the impact assessment for the DAN project is based on the maximum volume of discharged produced water, the maximum number of wells and realisation of new developments.

### 4.2 Methodology for assessment of impacts

The potential impacts of the DAN project on the environmental and social receptors (e.g. water quality, climate and fishery) are assessed for exploration, production and decommissioning.

The assessment covers the direct and indirect, cumulative and transboundary, permanent or temporary, positive and negative, impacts of the project. Impacts are evaluated based on their nature, type, reversibility, intensity, extent and duration in relation to each receptor (social and environmental).

The proposed methodology used for assessment of impacts includes the following criteria for categorising environmental and social impacts:

- Value of the receptor
- Nature, type and reversibility of impact
- Intensity, geographic extent and duration of impacts
- Overall significance of impacts
- Level of confidence

4.2.1 Value of receptor

Various criteria are used to determine value/sensitivity of each receptor, including resistance to change, rarity and value to other receptors (Table 4-1).

**Table 4-1 Criteria used to assess the value of receptor.**

Value	
<b>Low</b>	A receptor that is not important to the functions/services of the wider ecosystem/socioeconomy or that is important but resistant to change (in the context of project activities) and will naturally or rapidly revert to pre-impact status once activities cease.
<b>Medium</b>	A receptor that is important to the functions/services of the wider ecosystem/socioeconomy. It may not be resistant to change, but it can be actively restored to pre-impact status or will revert naturally over time.
<b>High</b>	A receptor that is critical to ecosystem/socioeconomy functions/services, not resistant to change and cannot be restored to pre-impact status.

4.2.2 Nature, type and reversibility of impacts

Impacts are described and classified according to their nature, type and reversibility (Table 4-2).

**Table 4-2 Classification of impacts: Nature, type and reversibility of impacts**

Nature of impact	
<b>Negative</b>	Impacts that are considered to represent an adverse change from the baseline (current condition).
<b>Positive</b>	Impacts that are considered to represent an improvement to the baseline.
Type of impact	
<b>Direct</b>	Impacts that results from a direct interaction between a planned project activity and the receiving environment.
<b>Indirect or secondary</b>	Impacts which are not a direct result of the project, but as a result of a pathway (e.g. environmental). Sometimes referred to as second level or secondary impacts.
<b>Cumulative</b>	Impacts that result from incremental changes caused by past, present or reasonably foreseeable human activities with the project.
Degree of reversibility	
<b>Reversible</b>	Impacts on receptors that cease to be evident after termination of a project activity.
<b>Irreversible</b>	Impacts on receptors that are evident following termination of a project activity.

#### 4.2.3 Intensity, geographic extent and duration of impacts

Potential impacts are defined and assessed in terms of extent and duration of an impact (Table 4-3).

**Table 4-3 Classification of impacts in terms of intensity, extent and duration**

<b>Intensity of impacts</b>	
<b>None</b>	No impacts on the receptor within the affected area.
<b>Small</b>	Small impacts on individuals/specimen within the affected area, but overall the functionality of the receptor remains unaffected.
<b>Medium</b>	Partial impacts on individuals/specimen within the affected area. Overall, the functionality of the receptor will be partially lost within the affected area.
<b>Large</b>	Partial impacts on individuals/specimen within the affected area. Overall, the functionality of the receptor will be partially or completely lost within and outside the affected area.
<b>Geographical extent of impacts</b>	
<b>Local</b>	Impacts are restricted to the area where the activity is undertaken (within 10 km).
<b>Regional</b>	There will be impacts outside the immediate vicinity of the project area (local impacts), and more than 10 km outside project area.
<b>National</b>	Impacts will be restricted to the Danish sector.
<b>Transboundary</b>	Impacts will be experienced outside of the Danish sector.
<b>Duration of impacts</b>	
<b>Short-term</b>	Impacts throughout the project activity and up to one year after.
<b>Medium-term</b>	Impacts that continue over an extended period, between one and ten years after the project activity.
<b>Long-term</b>	Impacts that continue over an extended period, more than ten years after the project activity.

#### 4.2.4 Overall significance

The definition of the levels of overall significance of impact are separated for environmental and social receptors (Table 4-4).

**Table 4-4 Classification of overall significance of impacts.**

Overall significance	Impacts on environmental receptors	Impacts on social receptors
<b>Positive</b>	Positive impacts on the structure or function of the receptor	
<b>Negligible negative</b>	No measurable impacts on the structure or function of the receptor.	
<b>Minor negative</b>	Impact to the structure or function of the receptor is localised and immediate or short-term. When the activity ceases, the impacted area naturally restores to pre-impact status.	Impact that is inconvenient to a small number of individual(s) with no long-term consequence on culture, quality of life, infrastructure and services. The impacted receptor will be able to adapt to change with relative ease and maintain pre-impact livelihood.
<b>Moderate negative</b>	Impact to the structure or function of the receptor is local or regional and over short- to medium-term. The structure or ecosystem function of the receptor may be partially lost. Populations or habitats may be adversely impacted, but the functions of the ecosystem are maintained. When the activity ceases, the impacted area restores to pre-impact status through natural recovery or some degree of intervention.	Impact that is inconvenient to several individuals on culture, quality of life, infrastructure and services. The impacted receptor will be able to adapt to change with some difficulties and maintain pre-impact livelihood with some degree of support.
<b>Major negative</b>	Impact to the structure or function of the receptor is regional, national or international and medium- to long-term. Populations or habitats and ecosystem function are substantially adversely impacted. The receptor cannot restore to pre-impact status without intervention.	Impact that is widespread and likely impossible to reverse for. The impacted receptors will not be able to adapt or continue to maintain pre-impact livelihood without intervention.

4.2.5 Level of confidence

It is important to establish the uncertainty or reliability of data that are used to predict the magnitude of the effects and the vulnerability of the receptors, as the level of confidence in the overall level of significance depends on it.

There are three levels of confidence for the impact:

- Low: Interactions are poorly understood and not documented. Predictions are not modelled and maps are based on expert interpretation using little or no quantitative data. Information/data have poor spatial coverage/resolution.
- Medium: Interactions are understood with some documented evidence. Predictions may be modelled but not validated and/or calibrated. Mapped outputs are supported by a moderate negative degree of evidence. Information/data have relatively moderate negative spatial coverage/resolution.
- High: Interactions are well understood and documented. Predictions are usually modelled and maps based on interpretations are supported by a large volume of data. Information/data have comprehensive spatial coverage/resolution.

## 5. ENVIRONMENTAL AND SOCIAL BASELINE

The environmental and social baseline contains a general description of each potential receptor, and site-specific information to the DAN project where applicable.

The baseline includes the following potential receptors:

- **Environmental**
  - Climate and air quality
  - Bathymetry
  - Hydrographic conditions
  - Water quality
  - Sediment type and quality
  - Plankton (phytoplankton and zooplankton)
  - Benthic communities (fauna and flora)
  - Fish
  - Marine mammals
  - Seabirds
  - Cultural heritage
  - Protected areas (Natura 2000, UNESCO world heritage, national nature reserves)
- **Social**
  - Marine spatial use
  - Fishery
  - Tourism
  - Employment
  - Tax revenue
  - Oil & Gas dependency

### 5.1 Climate and air quality

The North Sea is situated in temperate latitudes with a climate characterised by large seasonal contrasts. The climate is strongly influenced by the inflow of oceanic water from the Atlantic Ocean and by the large scale westerly air circulation which frequently contains low pressure systems /10/.

Air quality in the North Sea is a combination of global and local emissions. Industrialisation of the coast and inshore area adjacent to certain parts of the central North Sea has led to increased levels of pollutants in these areas which decrease further offshore, though shipping and platforms provide point sources of atmospheric pollution /141/.

### 5.2 Bathymetry

The North Sea is a part of the north-eastern Atlantic Ocean, located between the British Isles and the mainland of north-western Europe. The western part of the Danish North Sea is relatively shallow, with water depths between 20 – 40 m, while the Northern part is deeper (e.g. the Norwegian Trench and the Skagerrak; Figure 5-1).

The DAN project is located in an area with depths ranging from about 41 - 44 m /3/.





Figure 5-1 Bathymetry of the North Sea. Figure redrawn from Maersk Oil Atlas /3/.

### 5.3 Hydrographic conditions

The North Sea is a semi-enclosed sea. The water circulation is determined by inflow from the North Atlantic, water through the English Channel, river outflow from the Rhine and Meuse and the outgoing current from the Baltic Sea through Skagerrak (Figure 5-2). These inputs of water, in close interaction with tidal forces and wind and air pressures, create a complicated flow pattern in the North Sea. The DAN project area is in the central North Sea, where the dominant water circulation is eastward.

Hydrographic fronts are created where different water masses meet, and include areas of upwelling, tidal fronts, and saline fronts. Hydrographic fronts are considered of great importance to the North Sea ecosystems. No potential for hydrographic fronts has been identified in the central North Sea where the DAN project is located.



**Figure 5-2 Left: General water circulation in the North Sea. The width of arrows is indicative of the magnitude of volume transport /10/. Right: Potential for hydrographic fronts in the North Sea /10//2/.**

#### 5.4 Water quality

**Salinity:** Salinity in the North Sea varies from saline water in the west to brackish water along the coastal areas in the East. In the DAN project area, the salinity does not show much seasonal variation with surface and bottom salinity of c. 34-35 /3/.

**Temperature:** Temperature in the North Sea varies seasonally. The lowest temperatures are found in the Northern part of the North Sea, and the highest temperature in the shallower areas in the Southern North Sea. In the DAN project area, the surface temperature is approximately 7 °C in winter (January) and between 15-19 °C in summer (August), while the bottom temperature varies from 6-8 °C in winter (January) and 8-18 °C in summer (August) /3/.

**Nutrients:** Concentrations of nutrients in the North Sea surface layer have been modelled /3/. The concentrations are highest (>0.04 mg/l for phosphate, and >0.30 mg/l for nitrate) along the coastal areas, near output of large rivers. The concentrations in the surface layer in the DAN project area ranges between 0.025-0.035 mg/l for phosphate and between 0.1-0.15 mg/l for nitrate /3/.

**Heavy metals:** Water concentrations of metals in North Sea for cadmium ranges 6-34 ng Cd/l, copper 140-360 ng Cu/l, lead 20-30 ng Pb/l, mercury 0.05-1.3 ng Hg/l and nickel 100-400 ng Ni/l /29/. Metal cycles in the ocean are governed by seasonally variable physical and biological processes. The biologically driven metals (Cd, Cu, Ni) follow nutrient like distributions with higher concentration found in deep water. Certain metals, including Cd and Cu, exhibit higher concentrations near and on the shelf compared to the open sea areas /29/. No site-specific information on metals in seawater is available.

### 5.5 Sediment type and quality

The Danish sector of the North Sea is generally characterized by sediments consisting of sand, muddy sand and mud, with smaller areas of till with coarse sediments. The DAN project area substrate mostly consists of sand and muddy sand (Figure 5-3).




Figure 5-3 Seabed sediments in the North Sea. Figure redrawn from North Sea Atlas /3/.

Monitoring in May 2009 at the Dan F platform shows that the surface consists of fine sand with a median grain size (D50) between 0.15 - 0.19 mm. The silt/clay content of the sediment is approximately 1 % of the dry matter (DM) content. The dry matter content of the sediment is high, about 80%, which is typical for sand. The content of organic matter measured as loss on ignition (LOI) is low and around 0.5% of the dry matter of the sediment. The content of total organic carbon (TOC) is low and varies between 0.53 - 1.5 g/kg DM /6/.

The concentrations of THC in the surface sediment is between 2.1 - 110 mg/kg DM, the concentration of polycyclic aromatic hydrocarbons (PAH) are below 0.08 mg/kg DM while the concentrations of alkylated aromatic hydrocarbons (NPD) are below 0.14 mg/kg DM /6/.

Concentrations of metals (Cd, Cr, Cu, Pb and Zn /6/) are below the Lower Action Levels for dumping of seabed material defined by the DEPA, and thus characterised as having "average background levels or insignificant concentrations with no expected negative impact on marine organisms" /8/.

**5.6 Plankton**

The plankton community may be broadly divided into a plant component (phytoplankton) and an animal component (zooplankton). Plankton constitutes the main primary and secondary biomass in marine ecosystems and plays a fundamental role in marine food-webs.

In the North Sea, the phytoplankton is mainly light-limited in winter and nutrient-limited in the water above the thermocline in summer /10/. Figure 5-4 shows the phytoplankton colour index (PCI) for the North Sea over the course of the year. PCI is a visual estimation directly related to the biomass and abundance of the phytoplankton. The highest biomass and abundance of phytoplankton is found in the Eastern and Southern parts of the North Sea. The DAN project is in an area with an average biomass and abundance in comparison with the rest of the North Sea, and the phytoplankton community in the DAN project area is dominated by dinoflagellates and diatoms /3/.




Figure 5-4 Phytoplankton colour index (PCI) for the North Sea. Figure redrawn from North Sea Atlas /3/.

Zooplankton forms the link in the food web whereby the primary production by phytoplankton is channelled to the highest trophic levels through plankton-feeders such as herring (*Clupea harengus*), mackerel (*Scomber scombrus*), and sandeels (*Ammodytes* spp.). Generally, zooplankton abundance varies between areas owing to differences in production, predation, and transport. Nevertheless, the zooplankton community in the central North Sea is generally homogeneous /12/.

The zooplankton communities in the North Sea are dominated in terms of biomass and productivity by copepods, particularly *Calanus* species such as *C. finmarchicus* and *C. helgolandicus* /3/. Calanoid copepods are large crustaceans (in a planktonic context) which range in size between 0.5 - 6 mm and are an important prey item for many species at higher trophic levels. In the DAN project area, the abundance of copepods is intermediate compared to the North Sea, with 5.5 – 9.5 ind/m<sup>3</sup> of *C. finmarchicus* and 6.5 – 12 ind/m<sup>3</sup> for *C. helgolandicus* /3/.

The larger zooplankton, known as megaplankton, includes euphausiids (krill), thaliacea (salps and doliolids), siphonophores and medusae (jellyfish). Meroplankton comprises the larval stages of benthic organisms and fish that spend a short period of their lifecycle in the pelagic stage before settling on the benthos. Important groups within this category include the larvae of starfish and sea urchins, crabs and lobsters and some fish /11/.

## 5.7 Benthic communities

### 5.7.1 Benthic flora

Macrophytes (macroalgae and higher plants) grow in conditions that feature exceptionally diverse and dynamic light regimes. The water clarity and hydrodynamic conditions have profound effects on the quantity and quality of the light available for marine plants at specific localities, thus directly influencing the biomass and species composition of the benthic communities in the North Sea. The depth of the photic zone for benthic plants is traditionally defined as the depth where 1 % of the surface irradiance is available for photosynthesis /10/.

The water depth at the DAN project and in its vicinity is approximately 40-45 m. At this depth, it is highly unlikely that any macrophytes are to be found.

### 5.7.2 Benthic fauna

The benthic fauna consists of epifauna and infauna (organisms living on or in the seabed, respectively) such as crustaceans, molluscs, annelids, echinoderms.

The 50 m, 100 m, and 200 m depth contours broadly define the boundaries between the main benthic communities in the North Sea, with local community structure further modified by sediment type /13//14/. Descriptions of the spatial distribution of infaunal and epifaunal invertebrates show that the diversity of infauna and epifauna is lower in the southern North Sea than in the central and northern North Sea. Epifaunal communities are dominated by free-living species in the south and sessile species in the north. Large-scale spatial gradients in biomass are less pronounced /15/.

Biological monitoring in May 2009 in the DAN project area recorded a total of 85 species in 154 samples collected around the platform and at the reference stations. With respect to species richness the benthic fauna was dominated by polychaetes followed by bivalves, crustaceans, echinoderms and other taxonomic groups (Table 5.1). Polychaetes accounted for 55% of the benthic abundance while other taxa, including sea anemones, phoronids and nemerteans, contributed with 16%. Echinoderms, bivalves and crustaceans were almost equally abundant and each group contributed with less than 10 % of the abundance. Bivalves were by far the most important component of the benthic biomass (84%) while echinoderms (10%) and gastropods (4.5%) were secondmost important.

**Table 5.1 Composition of the benthic fauna around Dan in May 2009. Rounded average values for abundance and biomass /6/.**

Taxonomic group	Number of species*		Abundance		Biomass	
	2.2 m <sup>-2</sup>	%	ind.m <sup>-2</sup>	%	gDWm <sup>-2</sup>	%
Polychaeta	34	40.0	283	55.4	1.51	0.8
Bivalvia	16	18.8	43	8.5	168.42	84.0
Gastropoda	6	7.1	17	3.4	8.96	4.5
Crustacea	14	16.5	36	7.1	0.67	0.3
Echinodermata	8	9.4	49	9.6	19.95	10.0
Other taxa	7	8.2	82	16.0	0.99	0.5
<b>Total</b>	<b>85</b>	<b>100</b>	<b>511</b>	<b>100</b>	<b>200.5</b>	<b>100</b>

\* Sum of species in the 154 samples collected (143 cm<sup>2</sup> each = 2.2 m<sup>2</sup>)

Figure 5-5 shows benthic fauna in the North Sea as assemblages of benthic fauna in the North Sea.




Figure 5-5 Assemblages of the benthic fauna in the North Sea. Figure redrawn from North Sea Atlas /3/.

### 5.8 Fish

Approximately 230 species of fish are found the North Sea. Fish species diversity is low in the shallow southern North Sea and eastern Channel and increases westwards. Species diversity is also generally higher close to shore as the habitat diversity increases. Most of the variability of the fish stocks is due to variation in egg and larval survival which is thought to be regulated by a number of factors, such as sea temperature and currents affecting larval drift to nursery grounds, as well as density-dependent predation on the eggs and larvae. Annual variability in recruitment of juveniles can differ by a factor of 5 for plaice, 50 for sole and more than 100 for haddock. Most species show annual or inter-annual movements related to feeding and spawning /10/.

A fish survey was carried out in the period from November 2002 to July 2003 at the Halfdan platform located about 8-21 km from the DAN project area. A total of 16 species of fish were registered: eight pelagic or semi-pelagic (Atlantic horse mackerel, Atlantic mackerel, cod, grey gurnard, herring, sandeel, sprat, whiting), and eight benthic species (American plaice, common dab, common dragonet, European plaice, haddock, hooknose/armed bullhead, lemon sole, lumpfish) /19/.

The dominating species were: sprat, herring, whiting, grey gurnard, Atlantic horse mackerel, Atlantic mackerel, common dab, American plaice and European plaice. Herring and sprat were registered during the fall, whereas Atlantic horse mackerel and Atlantic mackerel were registered in the summer period. Common dab, American plaice and grey gurnard were registered all time of the year.

The abundance of fish in the central North Sea is relatively low in comparison to other parts of the North Sea. The fish fauna is characterised by common dab, grey gurnard and whiting /150/.

The biology of the dominating species registered in the area is described in Table 5.2.

**Table 5.2 Distribution and biology of the dominating species registered in the area /23//24/. Further information on spawning areas and catch are presented for selected species in /3/.**

Species	Distribution and biology
<b>Atlantic horse mackerel</b> ( <i>Trachurus trachurus</i> )	Horse mackerel has a restricted distribution during summer, with the greatest densities in the south-eastern North Sea and adults also being found along the shelf edge in the northern North Sea. The species is notably absent from the central North Sea. Juvenile horse mackerel are pelagic feeders that prey on planktonic organisms. Larger individuals feed on small fish (e.g. herring, cod and whiting). Peak spawning in the North Sea falls in May and June. Spawning occurs off the coasts of Belgium, the Netherlands, Germany, and Denmark.
<b>American plaice</b> ( <i>Hippoglossoides platessoides</i> )	American plaice can be found throughout the North Sea. It prefers soft bottoms. Larvae feed on plankton, diatoms and copepods. Preferred food items for larger fish includes sea urchins, brittle stars, polychaetes, crustaceans and small fish. Spawning takes place during spring at 100-200 meter depth.
<b>Atlantic mackerel</b> ( <i>Scomber scombrus</i> )	Mackerel are widespread throughout the North Sea. Mackerel feed on a variety of pelagic crustaceans and small fish. In the North Sea, mackerel overwinter in deep water along the edge of the continental shelf and, in the spring, adult mackerel migrate south to the spawning areas in the central North Sea with extensions along the southern coast of Norway and in the Skagerrak. Spawning takes place between May and July.
<b>Common dab</b> ( <i>Limanda limanda</i> )	Dab is a demersal fish. It lives on sandy bottoms down to depths of about 150 metres. Preferred food items includes sea urchins, brittle stars, polychaetes, crustaceans, mussels and small fish. In the North Sea spawning takes place between April and June.
<b>European plaice</b> ( <i>Pleuronectes platessa</i> )	European plaice has a preference for sandy sediments although older age groups may be found on coarser sand. During summer juvenile plaice are concentrated in the Southern and German Bights and also occur along the east coast of Britain and in the Skagerrak and Kattegat. Juveniles are found at lower densities in the central North Sea and are virtually absent from the north-eastern part. Plaice is an opportunistic species which primarily forage on molluscs and polychaetes. Plaice spawns in winter from January to March. Spawning areas occur in the central part of the North Sea and in the English Channel.
<b>Grey gurnard</b> ( <i>Eutrigla gurnardus</i> )	Grey gurnard occurs throughout the North Sea. Most common on sandy bottoms, but also on mud, shell and rocky bottoms. During winter, grey gurnards are concentrated to the northwest of the Dogger Bank at depths of 50-100 m, while densities are low in areas off the Danish coast, and in the German Bight and eastern part of the Southern Bight. Juveniles feed on a variety of small crustaceans. The diet of older specimens mainly consists of larger crustaceans and small fish. The distribution maps indicate a

Species	Distribution and biology
	marked seasonal northwest-southeast migration pattern that is rather unusual. The population is concentrated in the central western North Sea during winter and spreads into the southeastern part during spring to spawn. In the northern North Sea, such shifts appear to be absent. Spawning takes place in spring and summer.
<b>Herring</b> ( <i>Clupea harengus</i> )	Within the North Sea herring may be found everywhere. The pelagic larvae feed on copepods and other small planktonic organisms while juvenile mainly feeds on Calanoid copepods but euphausiids, hyperiid amphipods, juvenile sandeels and fish eggs are also eaten. Larger herring also consuming predominantly copepods with small fish, arrow worms and ctenophores as an aside. After spending their first few years in coastal nurseries, two-year-old herring move offshore into deeper waters, eventually joining the adult population in the feeding and spawning migrations to the western areas of the North Sea. Herring is a demersal spawner on relatively shallow water depositing sticky eggs on coarse sand, gravel, shells and small stones. The fish congregate on traditional spawning grounds, many of which are on shoals and banks and in relatively shallow water.
<b>Sprat</b> ( <i>Sprattus Sprattus</i> )	Sprat is most abundant south of the Dogger Bank and in the Kattegat. Larvae feed on diatoms, copepods and crustacean larvae. After metamorphosis larger planktonic organisms are also eaten. Spawning occurs in both coastal and offshore waters during spring and late summer, with peak spawning between May and June.
<b>Whiting</b> ( <i>Merlangius merlangus</i> )	High densities of both small and large whiting may be found almost everywhere throughout the North Sea. The species is typically found near the bottom in waters at 10 to 200 m depth. Pelagic larvae feed on nauplii and copepodite stages of copepods. Immature whiting feed on crustaceans such as euphausiids, mysids and crangonid shrimps whereas mature whittings feed almost entirely on fish. Spawning takes place from January in the southern North Sea to July in the northern part.

There are two main forms of spawning: Demersal and pelagic spawning.

Demersal spawners lay their eggs on the seafloor, algae or boulders. The preferred habitat for demersal spawners is species specific.

Pelagic spawners have free floating eggs that are fertilized in the water column. Spawning grounds for pelagic spawners are often large and less well defined as they can move from year to year. Hydrographic conditions that are essential for the pelagic spawning have an important role regulating the boundaries of the spawning grounds. Pelagic spawning takes place mostly at depths of 20-100 m. Pelagic eggs and larvae are more or less passively carried around by ocean currents. Some are carried to nursery areas others stay in the water column. Larval growth and transport of larvae and eggs are regulated by a variety of environmental factors e.g. current, wind and temperature.

A fish survey was carried out in the period from November 2002 to July 2003 at the Halfdan platform located about 8-21 km from the DAN project area. Fish eggs from the following 13 species were registered: Common dab, European plaice, American plaice, cod, lemon sole, Atlantic mackerel, whiting, turbot, greater weever, grey gurnard, Mediterranean scaldfish, Arctic rockling and common dragonet /19/. It is likely that these species also spawn at the DAN project area.

The DAN project is in an area identified as a relatively important spawning ground for cod and whiting. Mackerel and plaice are also know to be spawning in the area (Figure 5-6), but does not seem to be an important spawning and nursery area for other commercial species /3//22/.




Figure 5-6 Spawning grounds for cod and whiting in the North Sea. Figure redrawn from North Sea Atlas /3/.

## 5.9 Marine Mammals

Harbour seal, grey seal, white-beaked dolphin, minke whale and harbour porpoise are the most common marine mammals in the North Sea /28/. The distribution and biology of these species as well as their habitat preference are described in Table 5-3.

**Table 5-3 Distribution and biology of the most common marine mammals; harbour seal, grey seal, harbour porpoise and white-beaked dolphin /30//31//32//33//40/.**

Species	Distribution and biology
<b>Harbour seals</b> <i>(Phoca vitulina)</i>	Harbour seals are one of the most widespread of the pinnipeds. They are found throughout coastal waters of the Northern Hemisphere, from temperate to Polar Regions. Harbour seals are mainly found in the coastal waters of the continental shelf and slope, and are also commonly found in bays, rivers, estuaries and intertidal areas. At sea, they are most often seen alone, but occasionally occur in small groups. Haul-out sites include rocks, sand and shingle beaches, sand bars, mud flats, vegetation and a variety of man-made structures /30/.
<b>Grey seals</b> <i>(Halichoerus grypus)</i>	Grey seals have a cold temperate to sub-Arctic distribution in North Atlantic waters over the continental shelf. They often haul out on land, especially on outlying islands and remote coastlines exposed to the open sea /32/.
<b>White-beaked dolphin</b> <i>(Lagenorhynchus albirostris)</i>	White-beaked dolphins have a wide distribution and inhabit cold temperate to subpolar waters of the North Atlantic. White-beaked dolphins inhabit continental shelf and offshore waters of the cold temperate to subpolar zones, although there is evidence suggesting that their primary habitat is in waters less than 200 m deep. The species is found widely over the continental shelf, but especially along the shelf edge /33/.  Two white-beaked dolphins were observed during aerial surveys in the Southern Maersk area in March 2008. No animals have been registered by acoustic monitoring, and the species is considered uncommon in the Southern Maersk area /40/.
<b>Harbour porpoise</b> <i>(Phocoena phocoena)</i>	Harbour porpoise are found in cold temperate to sub-polar waters of the Northern Hemisphere. They are usually found in continental shelf waters, and frequent relatively shallow bays, estuaries, and tidal channels /31/.  Harbour porpoise is the most common whale species in the North Sea, and the only marine mammal which frequently occurs in the Maersk Oil area /40/. They are mostly found in the eastern, western and southern parts of the North Sea, and generally found in low densities in the central part of the North Sea (Figure 5-7). The DAN project area is not of particular importance to harbour porpoise, and few individuals are observed.  Aerial surveys in the Southern Maersk area in May show densities of 0.25-0.4 harbour porpoises/km <sup>2</sup> near the platforms, and few animals in autumn. However, acoustic monitoring show high activity in autumn /40/. A recent study at the Dan platform /139/ showed that harbour porpoises are present around the platform all year with the highest echolocation activity during fall and winter.
<b>Minke whale</b> <i>(Balaenoptera acutorostrata)</i>	The minke whale is a cosmopolitan species found in all oceans and in virtually all latitudes, including the Northeast Atlantic. Minke whale occurs in both coastal and offshore waters and preys on a variety of species in different areas. Less than 0.025 animals/km <sup>2</sup> is expected in the central North Sea /33/.




Figure 5-7 Distribution of harbour porpoise in the North Sea. Figure redrawn from North Sea Atlas /3/.

The periods where the animals may be vulnerable to disturbance are related to the reproductive cycle (Table 5-3). The reproductive cycle of seals is primarily on land, while harbour porpoise is at sea.

Table 5-4 Time of year where animals are breeding (B), moulting (M) or mating (A). No data available for the other species.

Species	J	F	M	A	M	J	J	A	S	O	N	D
Grey seal		B	BA	A		M	M	M				
Harbour seal						B	BA	M	M			
Harbour porpoise					B	B	A	A				

## 5.10 Seabirds

Seabirds spend most of their life at sea but breed on rocky coasts and cliffs. In the North Sea region, common seabirds include fulmars, gannets and auk species, kittiwakes and skuas.

The spatial distribution of the key species of seabirds is summarised in Table 5-5, based on the distribution presented in the North Sea Atlas /3/ and a three-year aerial seabird monitoring survey in 2006-2008 covering the DAN project area /40/.

**Table 5-5 Spatial distribution of key species /3//40/.**

Species	Spatial distribution and biology in the North Sea
Red and black-throated diver ( <i>Gavia stellata</i> , <i>G. arctica</i> )	The two species, which are sensitive to oil pollution due to their pursuit-diving behaviour and low fecundity rate, are non-breeding visitors to the North Sea. Their sensitivity to oil pollution increases during October-November (Red-throated) and March-April (Black-throated) when the birds are undergoing moult of their flight feathers. In spring, the highest densities of red- and black-throated divers are found along the coast of Denmark, in the Wadden Sea and in the English Channel. In winter, the distribution is more restricted and the highest densities are found along the coast of Denmark and northern part of the shallow area off the Wadden Sea. Almost all birds are found in waters of riverine influence shallower than 35 m, and both species are rare (0 birds/km <sup>2</sup> ) in the DAN project area /3/, with few observations during the aerial survey /40/.
Northern fulmar ( <i>Fulmarus glacialis</i> )	The species is the most abundant seabird in the North Sea. In summer, relatively high densities of Northern fulmar are found at many locations throughout the North Sea with the peak densities located along the southern edge of the Norwegian Trench. In winter, the highest densities are found west of Norway and northwest of Jutland Bank. In the southern part of the North Sea Northern Fulmars are found in lower densities in winter than during summer. In the Southern Maersk Oil activity area, Northern Fulmar occurs at relatively high densities in spring, summer and autumn (up to 24 birds/km <sup>2</sup> /40/ or up to 360 birds/km <sup>2</sup> /3/), and is less abundant in winter (<2 birds/km <sup>2</sup> ) /3//40/.
Northern gannet ( <i>Morus bassanus</i> )	Northern gannets are found in high densities east and north of the UK from spring to autumn. In late summer-autumn high density areas are also found near the German and Dutch coasts. In winter, the northern gannet is patchily distributed and found at low to high densities throughout the North Sea. In the DAN project area, northern gannets occur mainly in low densities (< 1 birds/km <sup>2</sup> ) in winter, spring and summer /3//40/, but relatively high densities (up to 23 birds/km <sup>2</sup> ) were observed during autumn /40/.
Great skua ( <i>Stercorarius skua</i> )	Great skua occurs in low densities from northeast of Greater Fisher Bank to the Norwegian Trench, north of the UK coast, and in few small isolated patches. Unlike in spring-summer, the great skua occurs over much of the North Sea during late summer-autumn. In the DAN project area, the species occurs in low densities (0 birds/km <sup>2</sup> /3/), with few observations during the aerial surveys /40/.
Common gull ( <i>Larus canus</i> )	The common gull is not observed over much of the North Sea, but with intermediate to high densities along the eastern part of the North Sea (e.g. Wadden Sea, German Bight, Jutland Bank, and some isolated patches bordering the eastern UK coast). In the DAN project area, the species is rare (0 birds/km <sup>2</sup> /3/).
Lesser black-backed gull ( <i>Larus fuscus</i> )	Lesser black-backed gulls are largely absent from much of the central and north-western parts of the North Sea, and are concentrated mostly in the eastern parts of the North Sea. In the DAN project area, the species occurs in low densities (0 birds/km <sup>2</sup> /3/).
Herring gull ( <i>Larus argentatus</i> )	The herring gull occurs throughout most of the coastal areas in the eastern North Sea, particularly around Norway and in Skagerrak. Relatively high densities are found in the German Bight, off the coast of the Netherlands, and in winter also in areas further offshore like areas around Dogger Bank. Both the distribution and the

Species	Spatial distribution and biology in the North Sea
	abundance of herring gulls seem mainly to be determined by working trawlers. The species is rare in the Maersk Oil activity areas (0 birds/km <sup>2</sup> /3/).
Great black-backed gull ( <i>Larus marinus</i> )	Like for the herring gull, the distribution and the abundance of great black-backed gull in the activity areas seems mainly to be determined by working trawlers. The species is common throughout the North Sea during winter, and the highest densities are found south and west of the Dogger Bank. In the DAN project area, the species is rare (0 birds/km <sup>2</sup> /3/).
Black-legged Kittiwake ( <i>Rissa tridactyla</i> )	In summer, the species is concentrated primarily in the western North Sea. Outside the breeding season, the species occurs throughout the North Sea with widespread intermediate to high density areas. Most extensive concentrations are found along the southern edges of the Norwegian Trench, northwest of Dogger Bank, off Borkum and in the Channel. In the DAN project area, the species is found in intermediate density (3.5 – 10 birds/km <sup>2</sup> /3//40/), and in spring and autumn large flocks are observed /40/.
Sandwich tern ( <i>Sterna sandvicensis</i> )	The species is mainly distributed in coastal waters on both sides of the North Sea. In spring highest densities are found off the German coast and the Netherlands. In summer-autumn highest densities are shown off the British coast just north of the Wash. In the DAN project area, the species is rare (0 birds/km <sup>2</sup> /3/), and the few observations during the aerial surveys confirm the low densities /40/
Common tern ( <i>Sterna hirundo</i> )	The species is absent throughout most of the offshore parts of the North Sea. In spring highest densities are found off the northern German coast and the Netherlands. In late summer highest densities are found off the Danish coast and the Netherlands. In the DAN project area, the species is rare (0 birds/km <sup>2</sup> /3/), and the few observations during the aerial surveys confirm the low densities /40/
Common guillemot ( <i>Uria aalge</i> )	The common guillemot is the second most abundant seabird in the North Sea. In early summer, high densities are found in the western parts, whereas the species is found in lower densities in other parts of the North Sea. In late summer, the species occurs in high densities in the central and eastern parts as they move across the North Sea to moulting areas south of the Norwegian Trench. The species is very sensitive to oil pollution due to its pursuit diving behaviour, and during August and September both the adults and the accompanying young are flightless, and hence highly sensitive to pollution. As seen for many other species of seabirds, the highest numbers in the activity areas seem to be associated with the areas of lowest water depth. In winter, the species occurs in high densities in the western part of the North Sea. In the Maersk Oil activity areas, the species occurs in low densities in early summer and found in intermediate densities in late summer. In winter, high densities of Common Guillemot are found to the southeast of the area. The highest density of 5 birds/km <sup>2</sup> in the DAN project area is found in March /3/.
Razorbill ( <i>Alca torda</i> )	In early and late summer, the razorbill is largely absent in most of the North Sea and the birds are concentrated in its western part. Higher densities are observed in late- than in early-summer. The razorbill is largely absent in most of the northern and central North Sea in winter when most birds are found in the Skagerrak and Kattegat and off the coasts of the UK and NL. In the DAN project area, the species is found in densities of up to 2.5 birds/km <sup>2</sup>
Little auk ( <i>Alle alle</i> )	The little auk is concentrated along the Norwegian Trench and NW of Dogger Bank during winter, and the species occurs in rather low densities (<5 birds/km <sup>2</sup> /3/) in the DAN project area.

The four species of gulls (common gull, lesser black-backed gull, herring gull, great black-backed gull) are presented as rare (0 birds/km<sup>2</sup>) /3/. However, aerial surveys show frequent observations and density estimates for gulls range up to 11 gulls/km<sup>2</sup>, with the highest densities in autumn /40/.

5.10.1 International Bird Areas (IBAs)

Important Bird Areas (IBAs) are key sites for future conservation. A site is recognised as an IBA only if it meets certain criteria, based on the occurrence of key bird species that are vulnerable to global extinction or whose populations are otherwise irreplaceable. The Wadden Sea (in Dutch, German and Danish waters) and Skagerrak/Southwest Norwegian trench are both recognised as important areas for birds, more than 100 km from the DAN project area. There are no IBAs in the central North Sea /34/.

5.11 Cultural heritage

Cultural heritage in the North Sea includes submerged prehistoric sites that were once land, other coastal features such as early fish-traps, submerged structures from defending coast in the World Wars, and shipwrecks from all ages. Part of the floor of the North Sea is submerged land, and quite a number of villages in the Southern Bight have been submerged by the sea.

5.12 Protected areas

Protected areas are shown in Figure 5-8. Protected areas include Natura 2000 sites, Ramsar sites, UNESCO world heritage sites and nationally designated areas.




Figure 5-8 Protected areas. Figure redrawn from North Sea Atlas /3/.

#### 5.12.1 Natura 2000 sites

The Natura 2000 network comprises:

- Habitats Directive Sites (Sites of Community Importance and Special Areas of Conservation) designated by Member States for the conservation of habitat types and animal and plant species listed in the Habitats Directive
- Bird Directive Sites (Special Protection Areas) for the conservation of bird species listed in the Birds Directive as well as migratory birds

Natura 2000 sites have been designated in the central North Sea for Dogger Banke in UK, the Netherlands and Germany (Figure 5-8). The basis for designation is presented in section 10.

#### 5.12.2 Ramsar sites

Ramsar sites are wetlands of international importance, and are present in coastal areas of the North Sea. The Ramsar Convention requires Contracting Parties to 'formulate and implement their planning so as to promote the conservation of the wetlands included in the List, and as far as possible the wise use of wetlands in their territory' (article 3.1).

All Ramsar sites in the Danish sector of the North Sea are also designated Natura 2000 areas.

#### 5.12.3 UNESCO world heritage sites

The Wadden Sea in Denmark, Germany and the Netherlands have been appointed UNESCO world heritage site (Figure 5-8).

The Wadden Sea is the largest unbroken system of intertidal sand and mud flats in the world. It is a large, temperate, relatively flat coastal wetland environment, formed by the intricate interactions between physical and biological factors that have given rise to a multitude of transitional habitats with tidal channels, sandy shoals, seagrass meadows, mussel beds, sandbars, mudflats, salt marshes, estuaries, beaches and dunes. The area provide a habitat for numerous plant and animal species.


#### 5.12.4 Nationally designated areas

In Denmark, the Wadden Sea is designated as a national park. In addition, several nature reserves ("natur- og vildtreservat") have been appointed in Denmark along the west coast of Jutland, several inshore nature reserves (e.g. Nissum Fjord and Ringkøbing Fjord) (Figure 5-8).

### 5.13 Marine spatial use

The DAN project area is not used as shipping routes for the largest ships equipped with automatic identification systems (Figure 5-9, < approximately 100 per year) /3/.

The infrastructure of oil & gas and wind includes both existing and planned installations. In the North Sea, a number of oil & gas facilities are operational, and additional facilities are planned. Operational wind farms are only present in Danish waters off Esbjerg, while a number of wind farms are planned in UK and German waters. Pipelines and cables connecting platforms are not shown in the figure, but should also be considered when planning new projects.



**Figure 5-9 Ship traffic and infrastructure in 2012. Figure redrawn from North Sea Atlas /3/. Ship traffic is based on all ships fitted with AIS system i.e. ships of more than 300 gross tonnage engaged on international voyages, and cargo ships of more than 500 gross tonnage not engaged on international voyages and all passenger ships irrespective of size. Missing data in the middle of the North Sea is due to poor AIS receiving coverage and not lack of ships. Germany does not participate in the North Sea AIS data sharing program.**

Further spatial restrictions include military areas, dump sites and reclamation areas. Dump sites and reclamation areas are mainly located at a relatively short distance from the coast, and are not present in the central North Sea. Military uses constitute a small part of the sea-borne and coastal activities around the North Sea. There are extensive exercise areas, mainly in the United Kingdom, but also along the west coast of Jutland (Denmark).

**5.14 Fishery**

Fishery is an important industry in the North Sea. The main targets of major commercial fisheries are cod, haddock, whiting, saithe, plaice, sole, mackerel, herring, Norway pout, sprat, sandeel, Norway lobster, and deep-water prawn. Norway pout, sprat and sandeel are predominantly the targets of industrial fisheries for fish meal and oil, while other species are the targets of fisheries for direct human consumption /10/.

A historic overview of production, trade, employment and fleet size for fishery in Denmark is provided in Table 5-6 /36/.

**Table 5-6 Historic overview of production, trade, employment and fleet for fishery in Denmark /36/.**

	1990	2000	2010
<b>Production (thousand tonnes)</b>			
Inland	36	37	23
Marine	1482	1541	840
• Aquaculture	42	44	35
• Capture	1476	1534	828
Total	1518	1578	863
<b>Trade (USD million)</b>			
Import	1116	1806	2958
Export	2166	2756	4140
<b>Employment (thousands)</b>			
Aquaculture	0	0.8	0.4
Capture	6.9	4.6	2.4
Total	6.9	5.4	2.9
<b>Fleet (thousands)</b>			
Total	3.8	4.1	2.8

Landings of sandeel, European plaice, herring, cod, sprat and Norway pout are presented in the North Sea Atlas /3/. The landings are presented for one year (2013), and show that the central North Sea, including the DAN project area, has some importance to the Danish fishery for sandeel. In addition, some fishery takes place in the central North Sea, in particular for cod, sprat and European plaice.

As inter-annual variation can be significant, fishery data for a period of ten years have been extracted from the Danish AgriFish Agency /37/. The data has been extracted for Danish vessels for area IVB, which covers an area of 280,000 km<sup>2</sup> from the west coast of Jutland to the Eastern coast of the UK.

Estimated value for the landing from Danish vessels in the North Sea for the last ten years shows that the area IVB, where the DAN project is located, is important for the fishing industry (Table 5-7) /37/.

**Table 5-7 Total landings and value of fishery, as landed catch for important commercial species in the central North Sea (area IVB) /37/.**

	Overall		Species-species landed catch (tonnes)			
	Total landed catch (tonnes)	Total value (DKK)	Sandeel	Cod	Sprat	European plaice
<b>2005</b>	405,067	824,527,622	129,776	4,365	233,306	9,382
<b>2006</b>	376,174	894,837,171	239,144	3,556	97,208	9,721
<b>2007</b>	239,469	700,252,302	142,309	2,317	64,047	6,918
<b>2008</b>	320,488	696,990,031	231,321	2,596	62,680	6,854
<b>2009</b>	409,143	652,075,835	272,865	2,792	110,650	6,827
<b>2010</b>	344,744	858,381,192	250,676	3,359	68,827	7,837
<b>2011</b>	388,927	990,124,457	263,971	2,736	98,484	9,932
<b>2012</b>	160,556	746,792,906	47,439	2,547	70,907	9,557
<b>2013</b>	263,373	875,992,562	183,330	1,917	46,258	10,707
<b>2014</b>	328,063	855,349,857	147,963	2,712	135,366	9,551

**5.15 Tourism**


Tourism is a multi-disciplinary feature, and includes both traditional tourism such as hospitality as well as events within conferences, music and sports. Tourists in Denmark are primarily Danish and German, and to a minor extent tourists from Sweden, Norway and the Netherlands.

Based on recent report with 2012 data from VisitDenmark /38/, tourism creates 122,500 FTEE (full time employee equivalent), which corresponds to ~4 % of the total FTEE in Denmark. These jobs are typically within hospitality, transport and trade. Tourism creates a direct economic added value of 24 billion DKK.

Tourism is associated with land and the coast, and no tourism is present in the central North Sea.

**5.16 Employment**

According to Statistics Denmark /39/, the largest employment sectors in 2013 are the public sector and trade/transport.



**Figure 5-10 Employment per sector in Denmark in 2013 /39/.**

Oil & gas activities in the North Sea create a significant number of workplaces both on-and offshore /35/. The oil and gas sector employs approx. 15,000 persons in Denmark /53/. Of these, approx. 1,700 employees are directly employed at the oil companies. This means that when one employee is employed in the oil and gas companies, approx. 8 jobs are created in related industries. A large part of the indirect activities lies in e.g. the engineering consultancy and other consulting assistance. Employment in the sector ranges widely across types of job, but generally a high level of education is seen and approx. 60 % of the jobs are located around Esbjerg.

There is no specific statistics available for the west coast of Jutland.

### **5.17 Tax revenue**

Tax revenue and the profits made by the oil and gas sector have a positive impact on the danish economy. The state's total revenue is estimated to range from DKK 20 to DKK 25 billion per year for the period from 2014 to 2018 /35/.

The sector's impact in relation to taxes and dues are also substantial, as is the business sector, which by far contributes to the largest share of taxes and dues. In 2010, the total contribution of direct taxes and dues was approx. DKK 24 billion /53/.

### **5.18 Oil & gas dependency**

Denmark has been supplied with gas from its North Sea fields since the 1980s and has also exported natural gas, primarily to Sweden and Germany. This production has significantly impacted the security of supply and balance of trade. Denmark is expected to continue being a net exporter of natural gas up to and including 2025 and Maersk Oil has license to operate until 2042 /35/.

As part of a long-term Danish energy strategy, the oil and gas production is instrumental in maintaining high security of supply, at the same time as renewable energy represents an increasing share of the Danish energy mix /53/.

## 6. IMPACT ASSESSMENT: PLANNED ACTIVITIES

### 6.1 Impact mechanisms and relevant receptors

#### 6.1.1 Potential impact mechanisms

Potential impact mechanisms associated to the planned activities at the DAN project are summarised based on the project description (section 3) and the technical sections (appendix 1).

Potential impact mechanisms include:

- Underwater noise
- Physical disturbance of seabed
- Suspended sediment
- Discharges (physical and chemical)
- Solid waste
- Emissions
- Light
- Resource use
- Restricted zones
- Employment and tax revenue
- Oil and gas dependency

The source of the potential impact mechanisms is provided in Table 6-1. The sources of impacts are related to the activities described in the seven technical sections (appendix 1).

**Table 6-1 Sources of potential impact mechanisms for the DAN project. "X" marks relevance, while "0" marks no relevance.**

Potential impact mechanism	Sesimic	Pipelines & structures	Production	Drilling	Well stimulation	Transport	Decommissioning
<b>Underwater noise</b>	X	X	X	X	X	X	X
<b>Physical disturbance of seabed</b>	X	X	0	X	0	0	X
<b>Suspended sediment</b>	X	X	0	X	0	0	X
<b>Discharges</b>	X	X	X	X	X	X	X
<b>Solid waste</b>	X	X	X	X	X	X	X
<b>Emissions</b>	X	X	X	X	X	X	X
<b>Light</b>	X	X	X	X	X	X	X
<b>Presence/removal of of structures</b>	0	X	X	0	0	0	X
<b>Resource use</b>	X	X	X	X	X	X	X
<b>Restricted zones</b>	X	X	X	X	0	0	X
<b>Employment and tax revenue</b>	X	X	X	X	X	X	X
<b>Oil and gas dependency</b>	X	X	X	X	X	X	X



### 6.1.3 Marine strategy frameworks directive - descriptors

The list of receptors and impact mechanisms described in the ESIS can be directly related to the descriptors set within the Marine Strategy Framework Directive (MSFD; section 2.1.5). The MSFD outlines 11 descriptors used to assess the good environmental status of the marine environment. The environmental status of the Danish North Sea waters is described in details in /154/.

1. Biological diversity is maintained. The quality and occurrence of habitats and the distribution and abundance of species are in line with prevailing physiographic, geographic and climatic conditions.
2. Non-indigenous species introduced by human activities are at levels that do not adversely alter the ecosystems.
3. Populations of commercially exploited fish and shellfish are within safe biological limits, exhibiting a population age and size distribution that is indicative of a healthy stock.
4. All elements of the marine food webs, to the extent that they are known, occur at normal abundance and diversity and levels capable of ensuring the long-term abundance of the species and the retention of their full reproductive capacity.
5. Human-induced eutrophication is minimised, especially adverse effects thereof, such as losses in biodiversity, ecosystem degradation, harmful algal blooms and oxygen deficiency in bottom waters.
6. Sea-floor integrity is at a level that ensures that the structure and functions of the ecosystems are safeguarded and benthic ecosystems, in particular, are not adversely affected.
7. Permanent alteration of hydrographical conditions does not adversely affect marine ecosystems.
8. Concentrations of contaminants are at levels not giving rise to pollution effects.
9. Contaminants in fish and other seafood for human consumption do not exceed levels established by Community legislation or other relevant standards.
10. Properties and quantities of marine litter do not cause harm to the coastal and marine environment.
11. Introduction of energy, including underwater noise, is at levels that do not adversely affect the marine environment.

The receptors identified in the ESIS are related to the MSFD status indicators hydrography (D7), fish, harbour porpoise and benthic communities (D1, D6). The impact mechanisms for planned activities in the ESIS are related to the MSFD pressure indicators seabed (D6), discharges (D6, D8, D9) and underwater noise (D11). Each impact mechanism is further assessed for the relevant receptors in the following sections 6.2 and 6.3.

## 6.2 Assessment of potential environmental impacts

Impact assessment for planned activities for each relevant environmental receptor is presented in the following sections.

### 6.2.1 Climate and air quality

Impacts on climate and air quality relate to emissions.

#### 6.2.1.1 Emissions

Emissions have been estimated for the planned activities for the DAN project, and are presented in Table 6-3 for each of the activities.

**Table 6-3 Overview of estimated emissions for planned activities for the DAN project, provided per activity or per year. The maximum emissions have been used. Estimates have been calculated by Ramboll based on input from Maersk Oil."-" refers to an emission which has not been quantified.**

Activity (frequency)	Unit for which estimate is provided (duration)	Emissions					
		CO <sub>2</sub> (tonnes)	NO <sub>x</sub> (tonnes)	N <sub>2</sub> O (tonnes)	SO <sub>2</sub> (tonnes)	CH <sub>4</sub> (tonnes)	nmVOC (tonnes)
<b>Seismic</b>							
4D seismic (Every 4 years)	Per survey (~1 month)	3,330	60	0.2	2	0.3	2.5
Site survey (Every year)	Per survey (1 week)	40	0.7	0.003	0.02	0.003	0.03
Borehole seismic (Every year)	Per survey (2 days)	11	0.2	0.001	0.007	0.001	0.01
<b>Pipelines and structures</b>							
Dan area redevelopment	Four platforms	13,200	240	0.8	8	1.2	10
Dan accommodation	One platform	3,300	60	0.2	2	0.3	2.5
<b>Drilling</b>							
Drilling (43 new well, 14 re-drill)	Per well (150 days)	8,450	150	0.6	6	0.6	7
Well test, workover	Not quantified	-	-	-	-	-	-
<b>Well stimulation</b>							
Matrix acid well stimulation (2 per year)	Per well stimulation (2 weeks)	625	11.6	0.043	0.395	0.047	0.473
<b>Production</b>							
Flaring, fuel, vent	Per year	475,100	2,300	58	3.4	1,150	340
<b>Transport</b>							
Vessels, helicopters	Per year*	27.6	0.5	0.002	0.02	0.002	0.03
<b>Decommissioning</b>							
Well abandonment (126 wells)	per well (20 days)	1,125	20	0.08	0.8	0.08	0.9
Cleaning and removal of structures	Regnar, Kraka, Dan B, Dan FD and Dan E	70,460	1,310	5	45	5	53

\* Note that the calculation for vessels and helicopters are assuming 20% for each of the five ESIS projects.

Emissions are primarily caused by venting, flaring of gas and the use of fossil fuels for production.

Table 6-4 provides an overview of the estimated annual emissions from operation of the DAN project and the annual Danish emissions 2012, as well as total emissions during drilling and decommissioning.

**Table 6-4 Emissions from activities at the DAN project and national emissions numbers for Denmark /20/21/. "-" refers to an emission which has not been quantified.**

<b>Emissions</b>	<b>Annual Danish emissions 2012 (tonnes)</b>	<b>Total annual production emissions at DAN (excluding drilling, new structures and decommissioning) (tonnes)</b>	<b>Total emissions for drilling 57 wells (tonnes)</b>	<b>Total emissions for decommissioning 126 wells and existing structures (tonnes)</b>
<b>CO<sub>2</sub></b>	39,412,000	480,000	480,150	212,200
<b>N<sub>2</sub>O</b>	-	58	33	15
<b>NO<sub>x</sub></b>	116,071	2,385	8,665	3,830
<b>SO<sub>x</sub></b>	12,510	6	330	145
<b>CH<sub>4</sub></b>	-	1151	35	15
<b>nmVOC</b>	-	345	400	167

#### 6.2.1.2 CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> emissions

Greenhouse gases such as CO<sub>2</sub>, N<sub>2</sub>O, SO<sub>x</sub> and CH<sub>4</sub> have a direct impact on climate and air quality.

The greenhouse gasses have different warming potential /141/, as some have a longer lifetime in the atmosphere and a higher heat absorption than others. Per definition, CO<sub>2</sub> has a global warming potential (GWP) of 1, whereas the GWP is 21 of CH<sub>4</sub> and 310 of N<sub>2</sub>O /141/. By re-calculating the estimated emissions to a GWP, it is seen that CO<sub>2</sub> constitutes the largest emission of greenhouse gasses.

Both drilling and decommissioning are emissions related to specific activities, while the annual emissions occur every year until 2042. The annual emissions will therefore over the project life cycle be of the largest quantity.

The annual emissions for the DAN project (excluding drilling and decommissioning) contributes up to 1.2 % of the total annual CO<sub>2</sub> emission for Denmark until 2042 (percentile will depend on the development of annual Danish emissions). The impact is considered to be of small intensity, a transboundary extent and long-term duration. The impact on climate change from emissions for the DAN project is assessed to be of moderate negative overall significance.

#### 6.2.1.3 NO<sub>x</sub>, SO<sub>x</sub> and nmVOC emissions

NO<sub>x</sub> and SO<sub>x</sub> are air pollutants which are spread by the wind and deposited in the surroundings. The compounds have acidification effects, that can impact the environment in terms of defoliation and reduced vitality of trees, and declining fish stocks in acid-sensitive lakes and rivers. nmVOCs, can have a number of damaging impacts on human health. Some have direct toxic effects (e.g. carcinogenic), but nmVOCs can also have indirect effects on health by contributing to the formation of ground-level ozone, which causes respiratory and cardiovascular problems.

Emissions of NO<sub>x</sub> from the DAN project production corresponds to 2 % of the total annual emission of NO<sub>x</sub> and SO<sub>x</sub> corresponds 0.05 % of total annual emission of SO<sub>x</sub> in Denmark until 2042 (percentile will depend on the development of annual Danish emissions). The impact is considered an impact of small intensity, a transboundary extent and long-term duration. The impact on air pollution from emissions for the DAN project is assessed to be of moderate negative overall significance.

6.2.1.4 Overall assessment

The overall assessment of impacts on climate and air quality from planned activities at the DAN project is summarised in Table 6-5.

**Table 6-5 Potential impacts on climate and air quality from planned activities for the DAN project.**

Impact mechanism	Intensity	Extent	Duration	Overall significance	Level of concidence
<b>CO<sub>2</sub>, N<sub>2</sub>O, SO<sub>x</sub> and CH<sub>4</sub> emissions</b> (climate change)	Small	Transboundary	Long-term	Moderate negative	Medium
<b>NO<sub>x</sub>, SO<sub>x</sub> and nmVOC emissions</b> (air pollution)	Small	Transboundary	Long-term	Moderate negative	Medium

6.2.2 Hydrography

Impacts on hydrography relate to presence and removal of structures.

6.2.2.1 Presence and removal of structures

The DAN project consist of a number of structures and pipelines in the central North Sea. New structures and/or pipelines are planned for the DAN project, including up to five platforms and new pipelines. The impact to hydrography of new structures and the removal of the existing structures is assessed to be of small intensity, local extent and of a short-term duration. The overall impact to hydrography from presence of structures is assessed to be of minor negative significance.

When the DAN project is decommissioned, the existing platforms will be removed and disposed, while pipelines will be left in situ. The impact to hydrography of new structures and the removing of the existing structures is assessed to be of small intensity, local extent and of a short-term duration. The overall impact to hydrography from presence of structures is assessed to be of minor negative significance.

6.2.2.2 Overall assessment

The overall assessment of impacts on hydrogrpahy from planned activities at the DAN project is summarised in Table 6-6.

**Table 6-6 Potential impacts on climate and air quality from planned activities at the DAN project.**

Impact mechanism	Intensity	Extent	Duration	Overall significance	Level of concidence
<b>Presence of structures</b>	Small	Local	Short-term	Minor negative	Medium
<b>Removal of structures</b>	Small	Local	Short-term	Positive	Medium

6.2.3 Water quality

Potential impacts on water quality (turbidity, chemical composition etc.) from the DAN project are related to suspended sediment and chemical discharges during production, drilling, well stimulation and decommissioning.

#### 6.2.3.1 Suspended sediment

Various activities at the DAN project are expected to lead to sediment resuspension. New pipelines are trenched and buried to a depth of ca. 1.5-2 m below the seabed surface. Trenching of the pipeline into the seabed is done either by ploughing, water jetting or as mechanical cutting. During this process sediment are suspended into the water column. Based on experience from other pipeline projects, it is estimated that the suspended sediment will settle within a few hundreds meters of the disturbed area /157//131/. During decommissioning, physical disturbance will be related to removal of the existing structures.

The impact to water quality is assessed to be of small intensity, local extent and of a short-term duration. Overall, the impact to water quality from suspended sediment at the DAN project is considered to be of minor negative significance.

#### 6.2.3.2 Discharges

Maersk Oil uses chemicals in its operations, and is constantly optimising the use and discharge of chemicals to reduce its impact on the environment. Before any chemicals can be permitted for use and discharge offshore, an application must be submitted to the Danish authorities.

Maersk Oil discharges a number of chemicals. These chemicals are primarily classified as OSPAR category 'green', which pose little or no risk to the environment, or 'yellow', which does not bioaccumulate and degrade relatively rapidly (section 8.1.3). The discharge of red chemicals is not expected, but may occur in a very limited amount. Red chemicals are only used if safety, technological and environmental considerations cannot be met by alternative products. Maersk Oil has since 2008 been phasing out the use of red chemicals which contains components that bioaccumulate or degrade slowly (section 8.1.3).

Chemicals use and discharge to sea is only permitted after authorisation from the DEPA.

#### **Discharges during production**

The forecasted volume of discharged produced water at the DAN project is shown in Figure 3-8.

Traces of production chemicals may be present in the produced water. The production chemicals are categorized as 'green' or 'yellow' chemicals, which can usually be discharged without significant impact to the environment (section 8.1.3, appendix 1). Under special circumstances, red chemicals may also be used. A list of production chemicals, their function and their partitioning in oil/water phase is presented in appendix 1. Maersk Oil has flowmeters measuring the volume of discharged produced water, and water samples are regularly obtained for analysis of oil and chemical content.

Produced water may have toxic effects to the marine environment. Results from laboratory experiments suggest that the existing discharge of production water to be diluted from 10 to 10,000 times to reach a concentration where no acute toxic effects are expected. The toxicity of the water produced is determined, inter alia, the content of dispersed oil, BTEX, PAH and residues from chemicals used. Emissions of substances that are persistent or bioaccumulative, will in principle increase the general background level of the substance, but due to the relatively small amounts expected not to be measured in practice /1/.

Based on field monitoring, environmental impacts of produced water discharges on the environment are generally local, confined to within 1-2 km from the outlet, and the risk of impact from the operational discharges is low /46/. Hydrodynamic modeling of produced water at the DAN project suggests that the discharges are diluted relatively rapidly /42/. However, there is a risk of affecting the water quality from the current discharges of produced water at Dan F up to 31 km to the North East of the outlet /42/. The calculation of risk is based on the ratio between the Predicted Effect Concentration (PEC) and a Predicted No Effect Concentration (PNEC) of 1

where the expected concentration in the marine environment is the same as the concentration without expected effects of chemicals. The calculation is conservative and only applies to the most sensitive organisms.

The impact to water quality is assessed to be of small intensity, with a local to regional extent and of a short-term duration due to rapid dilution of the discharged water. Overall, the impact to water quality from discharge of produced water at the DAN project on is assessed to be of minor negative overall significance.

During production other minor negative discharges take place, these include discharges from vessels, and cooling water from production platforms. These discharges are considered negligible in comparison with the produced water, and are not assessed further.

**Discharges during drilling**

There are currently 3 free well slots at Dan F and 14 re-drilling are expected at the existing wells. In addition, a planned development project includes the construction of up to 40 new wells. Typically, it takes between 60 and 150 days to drill a well. Water-based mud and cuttings will be discharged to the sea, whereas oil-based mud and cuttings will be brought onshore to be dried and incinerated.

Cuttings from the formation collected in the water-based mud section of the well will be discharged to the sea, along with the drilling mud and material used for cementing (mostly cement and chemicals).

Discharges of cuttings can amount to 1800 tons of cuttings per well (appendix 1). When discharged to the sea water-based mud and cuttings, which are slurries of particles of different sizes and densities in water containing dissolved salts and organic chemicals, form a plume that dilutes rapidly as it drifts away from the discharge point with the prevailing water currents. Field studies of the concentration of suspended solids in plumes of drilling mud and cuttings at different distances from the drilling activity have confirmed this pattern, concluding that the concentration of suspended drill cuttings and mud in the water column drops very quickly due to sedimentation and dilution of the material /45//46/.

Expected discharges of water-based drilling mud and cement are shown in Table 6-7. The discharges shown are based on the worst case - defined as the well that leads to the largest amount of discharges. Chemicals expected to be used are categorized as 'green' or 'yellow' chemicals, which can be discharged without significant impact to the environment.

**Table 6-7 Use and discharge of drilling mud and cement per well – worst case discharge scenario. The classification colour code is explained above.**

		Usage per well	Discharge per well
	Classification	Tons	Tons
Drilling mud		2421	2421
		994	994
Cement		631	76
		14	1.7

Based on a review of results of modeling and field studies of drilling mud and cuttings it has been concluded, that offshore discharges of water-based mud and associated cuttings will have little or no harmful effects on water column organisms. This conclusion is based on the rapid dilution in the water column and low toxicity to marine organisms of water-based mud and cuttings /45/. Environmental impacts of drilling discharges are local, in general confined to within 1 - 2 km from the point of discharge /46/. The typical chemicals discharged to sea during Maersk Oil drilling have been modelled in the EIA for Adda and Tyra /2/. The modelling was performed

for the water column and showed that the predicted effect concentration extended up to 7 km in the dominating current from the platform /2/.

The impact to water quality is assessed to be of small intensity, local extent and of a short-term duration due to dilution. Overall, the impact to water quality from discharge of drilling mud and cuttings at the DAN project is assessed to be of minor negative overall significance.

### Discharges during well stimulation

The potential 43 new wells and 14 re-drill/slot recover for the DAN project may be subjected to matrix acid stimulation or acid fracturing (no sand fracturing is expected). In addition to stimulation of the new wells, it is anticipated that approximately two well stimulations of existing wells may take place per year at the DAN project.

Expected discharges of chemicals during well stimulation at the DAN project include chemicals categorized as 'green' or 'yellow' chemicals which can usually be discharged without significant impact to the environment. Typical discharges during well stimulation are presented in Table 6-8.

**Table 6-8 Use and discharge of chemicals per well stimulation. The classification colour code is explained above.**

	Classification	Usage per well	Discharge per well
		Tons	Tons
Matrix well stimulation	Green	220	140
	Yellow	2603	522
Acid fracturing well stimulation	Green	194	134
	Yellow	2816	564

The amount of discharge per well stimulation (Table 6-8) is significantly less than discharges during drilling (Table 6-7). The impact to water quality is assessed to be of small intensity, local extent and of a short-term duration due to dilution. Overall, the impact to water quality from discharges during well stimulation at the DAN project on is assessed to be of minor negative significance.

### Discharges during testing of new pipelines

During pre-commissioning the pipeline is flooded with treated seawater that contains low concentration of corrosion inhibitor to prevent pipeline damage. Thereafter, the pipeline is cleaned and impurities are removed by pigging. After cleaning, the pipeline is pressure-tested using treated seawater. During those operations a total volume of treated seawater, corresponding to about 305 % of the pipeline volume, is expected to be discharged to the sea. The largest expected pipeline that will be installed in the redevelopment at the DAN project is expected to be up to 7 km long with a diameter of 16".

Hydrodynamic modelling indicates that the treatment water will dilute relatively rapidly. Modelling which was carried for a similar size pipeline for the Adda project suggests that there is an unlikely risk of impact to the water column up to approximately 3 km./2/.

The impact to water quality is assessed to be of small intensity, local extent and of a short-term duration due to rapid dilution of the discharged water. Overall, the impact to water quality from discharges during pre-commissioning at the DAN project is assessed to be of minor negative overall significance.

### Discharges during decommissioning

Minor negative discharges are expected during decommissioning activities. In general, all structures (jacket and topside) will be cleaned, before transport to shore. Waste will be brought onshore for disposal. The impact to water quality is assessed to be of small intensity, local extent

and of a short-term duration due to dilution. Overall, the impact to water quality from discharges during decommissioning at the DAN project is assessed to be of minor negative overall significance.

#### 6.2.3.3 Overall assessment

The overall assessment of impacts on water quality from planned activities at the DAN project is summarised in Table 6-9.

**Table 6-9 Potential impacts on water quality from planned activities at the DAN project.**

Impact mechanism	Intensity	Extent	Duration	Overall significance	Level of confidence
<b>Suspended sediment</b>	Small	Local	Short-term	Minor negative	High
<b>Discharges</b>	Small	Local/Regional	Short-term	Minor negative	High

Minor cumulative effects on the water quality between the various discharges cannot be ruled out, however due to the low toxicity of the discharges and the rapid dilution rate any effect is estimated to be very local and short-term. A review of all discharges from the Norwegian offshore petroleum industry suggests that the environmental impacts of discharges are local, in general confined to few kilometres from the platforms /46/.

#### 6.2.4 Sediment type and quality

Potential impacts on the sediment type and quality are related to physical disturbance on the seabed and discharges settling on the seabed that may affect the chemical and physical composition.

##### 6.2.4.1 Physical disturbance on the seabed

Physical disturbance on the seabed happens during site surveys, 4D seismic, drilling, installation of platforms and pipelines and decommissioning.

During site surveys, which are expected to occur annually, seabed coring will be undertaken and disturbance of seabed will occur where the sample is acquired, typically with an area of 0.1-0.25 m<sup>2</sup>. During 4D seismic surveys, presence of bottom nodes and cables may impact the seabed. The impacts on the seabed from nodes and cables is expected to be minor negative (each node is 40-50 cm). During drilling (up to 31 wells), a drilling rig will be present. The rig legs will be placed on the seabed, and are expected to sink 1-2 m into the seabed. The rig legs typically covers a few hundred m<sup>2</sup>.

New pipelines are trenched and buried to a depth of ca. 1.5-2.0 m below the seabed surface. Minor amounts of sediment will be resuspended during installation of new platforms. When the supporting jackets are placed on the seabed. During decommissioning, physical disturbance will be related to removal of the existing structures.

Disturbance to small areas of sandy sediments is expected to be short term, as sand will naturally re-settle in the disturbed areas. The impact to sediment type and quality is assessed to be of small intensity, local extent and of a short-term duration. The impact to sediment type and quality from physical disturbance is therefore assessed to be of minor negative overall significance.

##### 6.2.4.2 Discharges during drilling

Water-based drilling mud and drill cuttings are expected to be discharged to sea from up to 57 (43 new well and 14 re-drill) wells, and the cuttings and associated chemicals may settle on the seabed and impact the sediment quality.

Several field studies that have measured the concentration of suspended solids in plumes of drilling mud and cuttings at different distances from the drill rigs have confirmed this pattern. The measurements have shown that the concentration of suspended drill cuttings and mud in the water column drops very quickly due to sedimentation and dilution of the material /45//46/. Modelling of drilling mud and cuttings sedimentation for a typical Maersk Oil well shows that drilling mud will settle on the seabed at a thickness of less than 1 mm. Most of the drilling mud will settle in vicinity of the discharge location (1 - 2 km), depending on the current (Figure 6-1). Drill cuttings are heavier than the drilling mud and will settle rapidly. Model data shows that for a similar well discharge, a 50 mm-layer of cuttings could be expected within 50 m of the well. The thickness of the layer is expected to decrease to <1 mm within 200 m of the discharge (Figure 6-2).




Figure 6-1 Sedimentation of discharged water based drilling mud modelled for a typical well /1/.




Figure 6-2 Sedimentation of water based drill cuttings modelled for a typical well /1/.

Worst case scenario discharges from drilling of one well is approximately 1,800 tons of cuttings and approximately 3,500 tons of water based drill mud and cement. If all 57 wells at the DAN project (43 existing and 14 re-drill) are drilled, it would result in a total discharge of 102,600 tons of cuttings and 200,000 tons of water-based drill mud. The worst case scenario is estimated to be the case where the wells are drilled consecutively. The wells are located at five locations: Dan F has 3 existing well slots, a planned platform near Kraka has 10 well slots and 3 platforms each having 10 well slots are likely to be located approximately 4-7 km from Dan F.

The chemicals which are discharged with the mud and cuttings are categorized as 'green' or 'yellow' chemicals that are expected to have no significant impact on the environment (section 6.2.3). The mud usually contains barite or trace of heavy metals, while the cuttings may contain small quantities of oil. Chemical and biological seabed monitoring around Dan F shows that elevated concentrations of metals, THC, PAH and NPD in the sediment are local and rapidly decreasing in all directions with increasing distance from the platform /6/.

Sedimentation of drilling mud and cuttings may change the sediment grain size. However, seabed monitoring shows that the median grain size variation close to the Dan F platform fall within the natural range /6/.

The water based drilling mud may contain biodegradable organic additives, which may stimulate growth of microbial communities leading to depletion of oxygen in the sediments. Anaerobic, sulphate-reducing bacteria may further degrade the organic matter, producing hydrogen sulphide /45/. A monitoring campaign of the seabed around the Dan F platform reported that sulphide was present at 2 of the station closest to the platform /6/.

Based on the modelling results, the type of chemicals in the drilling mud and cuttings and the results from the monitoring campaign, the impact is assessed to be of small intensity, local extent and of a long-term duration. In conclusion, the impact to sediment type and quality from discharges is assessed to be of minor negative overall significance.

#### 6.2.4.3 Presence/removal of structures

Existing pipelines and structures (topsides, jacket) are present in the area. Areas which were previously sand has been altered to contain a hard substrate in the water column. Four new platforms and connecting pipelines are planned for the DAN project.

The existing structures will be removed as the DAN project is decommissioned, with planned decommissioning of Regnar within the next three years. During decommissioning buried pipelines will be cleaned, and left in situ, filled with seawater. Platform facilities and jackets will be cleaned, removed and brought to shore for dismantling.

The impact to sediment type and quality is assessed to be of small intensity, local extent and of a medium-term duration. The impact to sediment type and quality from presence of pipelines and platforms is assessed to be of minor negative overall significance.

#### 6.2.4.4 Overall assessment

The overall assessment of impacts on sediment type and quality from planned activities for the DAN project is summarised in Table 6-10.

**Table 6-10 Potential impacts on sediment type and quality from planned activities for the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Physical disturbance on seabed</b>	Small	Local	Short-term	Minor negative	Medium
<b>Discharges of drill cuttings</b>	Small	Local	Long-term	Minor negative	High
<b>Presence/removal of structures</b>	Small	Local	Short-term	Minor negative	High

Minor cumulative effects between the various discharges for the DAN project can be expected e.g. from sedimentation of drilling mud and cuttings, which will settle on top of the recently discharged drilling mud and drill cuttings (when drilling in the same area), however effects are estimated to be local. It is not possible to predict where the acid stimulation in the existing wells will be carried out simultaneously.

#### 6.2.5 Plankton

Potential impacts on plankton (phyto- and zooplankton) are related to underwater noise, suspended sediment, discharges and light.

##### 6.2.5.1 Underwater noise

Underwater noise is a form of energy which may impact plankton, due to e.g. disruption of cells (cell lysis). Underwater noise is expected to be generated from expected seismic activities (airguns, multibeam and sidescan), pile-driving during construction of new platforms, driving of conductors, drilling and various vessels in the DAN project area. Table 6-11 shows typical frequency and noise levels for these activities.

Little research has been conducted in relation to impacts of underwater noise to plankton from underwater noise, primarily focussed on emitted energy from airguns during seismic surveys. Mortality of plankton has been observed at close range (within 5 m) of the source of the seismic gun /54//55/. Behavioural and physiological effects are only expected to impact organisms close to (within a few metres) powerful noise sources e.g. seismic surveys and pile-driving /64/ /65/. A study found that close range seismic sound emission (2 m range) on snow crab eggs had impacts on larval development and settlement /66/.

Based on the abundance, productivity and size of planktonic populations and their high reproductive rate, plankton is expected to recover after disturbance. The impact is assessed to be of small intensity, local extent and short-term duration. The impact on plankton from underwater noise is assessed to be of negligible negative overall significance.

**Table 6-11 Typical frequency and noise levels of activities expected for the DAN project (based on appendix 1, /2//48/). n/a is not available.**

Activity	Frequency	Unit	Noise level at increasing distance from source				
			1m	1-500 m	3 km	5 km	10 km
<b>Seismic</b>							
Airgun (2D/3D/4D seismic)	0.005-0.200 kHz	Peak-to-peak (dB re 1 $\mu$ Pa <sup>2</sup> )	244	n.a.	n.a.	n.a.	n.a.
	0.005-80 kHz	RMS (dB re 1 $\mu$ Pa <sup>2</sup> )	179-266	n.a.	167	129	n.a.
		SEL (dB re 1 $\mu$ Pa <sup>2</sup> )	202-216	n.a.	n.a.	n.a.	n.a.
Multibeam echosounder	70-100 kHz	RMS (dB re 1 $\mu$ Pa <sup>2</sup> )	225-232	n.a.	n.a.	n.a.	n.a.
Sidescan sonar	100-900 kHz	RMS (dB re 1 $\mu$ Pa <sup>2</sup> )	220-226	n.a.	n.a.	n.a.	n.a.
<b>Pipelines and structures</b>							
Pile-driving*	0.03-20 kHz	Not specified (dB re 1 $\mu$ Pa <sup>2</sup> )	228	179.5	n.a.	n.a.	n.a.
<b>Production</b>							
Production platform	0.01-10 kHz	RMS (dB re 1 $\mu$ Pa <sup>2</sup> )	162	n.a.	n.a.	n.a.	n.a.
<b>Drilling and well stimulation</b>							
Driving of conductors	0.03-20 kHz	Not specified (dB re 1 $\mu$ Pa <sup>2</sup> )	228	179.5	n.a.	n.a.	n.a.
Drilling rig	0.002-1.2 kHz	Not specified (dB re 1 $\mu$ Pa <sup>2</sup> )	163	123	n.a.	n.a.	77
<b>Transport</b>							
Support vessel	0.01-20 kHz	RMS (dB re 1 $\mu$ Pa <sup>2</sup> )	122-192	120	n.a.	n.a.	n.a.
<b>Decommissioning**</b>							
Not available			n.a.	n.a.	n.a.	n.a.	n.a.

\*\*Noise levels for decommissioning are not provided, as activities are not specified for the DAN project. It is anticipated that no blasting will occur.

#### 6.2.5.2 Suspended sediment

Suspended sediments could potentially reduce light availability to phytoplankton and affect the ingestion rate, egg production, egg-hatching success and survival rate of zooplankton. However, only small amounts of sediment is expected to be suspended for the DAN project as part of the construction of four new platforms and pipelines as well as from decommissioning (see section 6.2.3.1).

Many larval species use their vision for feeding. Larvae of species like plaice, sole, turbot and cod sight their prey at a distance of only a few millimeters - the more turbid the water is, the harder it is for the larvae to locate and catch their prey. However, they can live a few days without food /86//119/, and a short-term increase in suspended sediment is expected to be of negligible impact. The feeding activity of herring fry is shown to be affected by concentrations of suspended matter above 20 mg/l /85/.

Pelagic fish eggs may be affected if suspended matter adheres to eggs, causing them to sink to the bottom, where there is a risk of oxygen depletion /85/. A laboratory study on herring eggs has shown that short term exposure to relatively high concentration of suspended matter do not affect the development of fish eggs /87/. Benthic eggs are associated with soft sediments, and are assessed to be less sensitive than pelagic eggs.

In addition, as the regenerating capacity of plankton is large, and the DAN project area is a small area in the central North Sea, no effect at population level is anticipated.

The sediment spread will be local, temporary and spatially distributed along new pipelines and structures during trenching and will mainly be limited to the lower parts of the water column. It is assessed that any impact on phytoplankton and zooplankton from suspended sediment will be negligible.

6.2.5.3 Discharges

Potential impacts on plankton from discharges are indirectly related to the impacts of different activities on the water quality, which are described in section 6.2.3.

Studies show that discharges of water based drilling chemicals may have short-term effects on phyto- and zooplankton communities /45//46//58/. The discharges of chemicals associated with production, drilling and stimulation may affect the phyto- and zooplankton communities. In general, it is expected that offshore discharges dilute rapidly and that only plankton found in the water column close to the discharge may be affected. Laboratory and field data confirms that the risk of significant biological impact is limited to 1 – 2 km from the discharges /46/. The effect of chemical discharges is expected to be minor negative, local and acute/non-persistent; therefore, the overall effect is assessed to be of minor negative significance.

6.2.5.4 Light

Vertical migration in the water column by some phytoplankton and zooplankton species may be influenced by light. Only Dan F and Dan B are permanently illuminated, while unmanned platform are only lit by safety lightings.

Light has been reported as a fundamental factor controlling the daily vertical migration of zooplankton /60/. Plankton migrate closer to the surface on dark nights than they do on clear, moonlit nights /62/. Some species of plankton have been reported foraging in darkness to avoid predation, only to be intensively predated when illuminated by a rising full moon /63/. However, planktonic organisms are per definition carried around with the prevailing currents, and light is expected to be detectable for planktonic organisms only in the vicinity of the platforms.

The potential affects are expected to be local, and may impact individuals but will not impact plankton populations in the North Sea. The impact on plankton from light at the DAN project is assessed to have negligible negative overall significance.

6.2.5.5 Overall assessment

The overall assessment of impacts on plankton from planned activities at the DAN project is summarised in Table 6-12.

**Table 6-12 Potential impacts on plankton from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Underwater noise</b>	Small	Local	Short-term	Negligible negative	Medium
<b>Suspended sediment</b>	Small	Local	Short-term	Negligible negative	High
<b>Discharges</b>	Small	Local	Short-term	Minor negative	Medium
<b>Light</b>	Small	Local	Immediate	Negligible negative	Low

The cumulative impact to plankton is not well known. However, little geographical overlap between the various impacts is expected. Due to the high reproductive capacity of plankton, it is expected that any cumulative impacts will be negligible negative.

#### 6.2.6 Benthic communities

Potential impacts on the benthic community are related to underwater noise, physical disturbance on seabed, suspended sediment, discharges and physical presence of structures.

##### 6.2.6.1 Underwater noise

Underwater noise may potentially impact benthic communities through e.g. behavioural and physiological effects.

Most invertebrates typically do not have delicate organs or tissues whose acoustic impedance is significantly different from water. Unlike e.g. fish with swim bladders, the general consensus regarding underwater noise effects on invertebrates and planktonic larvae under field conditions, is, that very few behavioral or physiological effects are expected unless the organisms are within a few metres of noise sources around 240 dB re 1  $\mu$ Pa /61/.

The impact is assessed to be of small intensity, local extent and short-term duration. The impact on benthic communities from underwater noise is assessed to be of negligible negative overall significance.

##### 6.2.6.2 Physical disturbance on the seabed

Physical disturbance on the seabed may physically impact the benthic fauna in the disturbed area. For the DAN project, the disturbed area is related to pipelaying operations, the presence of cables from seismic surveys and legs from the drilling rig and new platforms. During decommissioning, physical disturbance will be related to removal of the existing structures.

At the DAN project, the disturbed area is small (mostly related to cables from seismic surveys and legs from a drilling rig). In the disturbed area, benthic fauna will be displaced. Re-establishment will depend on the species present and their life cycle. Studies from the North Sea shows that benthic faunal communities on a sandy seabed generally re-establish during a period of 2-5 years /67//156/.

The intensity of the impact from physical disturbance on the seabed is assessed to be small with a local extent and of medium-term duration. Overall, the impact is assessed to be minor negative.

##### 6.2.6.3 Suspended sediment

Following the installation of new platforms and pipelines, sediments will be dispersed to the water column. The impact will depend on the concentration of suspended sediments and the duration. Increased concentrations of suspended sediment in the water column may negatively affect benthic fauna by e.g. clogging of respiratory system or feeding apparatus. Smothering by re-suspended sediment as it settles may have direct mechanical effects on fauna and may result in modification of the substratum. The maximum impact may not be immediate, as some indirect ecological changes, such as exposed organisms becoming more vulnerable to predation, may take place /68/. For this reason, the number of scavengers generally increases at recently dredged sites /69//70//70/. For these types of animals, seabed intervention works will likely have a positive effect.

The impact on suspension feeders 1-1.5 km due to suspended sediment and sedimentation from a sand-extraction site has been found to be short-term, resulting in lower activities and a reduced feeding rate, while no longterm effects were found at this distance. The study concluded that any considerable effects on benthic fauna occurred only in the vicinity of the extraction sites /75/.

A study of the effects of pipeline construction on benthic invertebrates showed that six months after disturbance there was no significant difference between the mean number of total individuals in the impacted site and in the reference areas /75/. Other studies have shown that benthic invertebrate communities began to re-colonise a disturbed seabed area very quickly, but that it could take up to five years for the structure of the community to recover /72/.

Due to the limited amount of dispersed sediment and sedimentation during construction of new platforms and pipelines impacts are assessed to be very local. Any measurable impacts to the benthic community are estimated to be within 0-10 meters from the activity.

The intensity of the impact from suspended sediment is assessed to be small with a local extent and short-term duration. Overall, the impact is assessed to be of minor negative overall significance.

#### 6.2.6.4 Discharges

Potential impacts on the benthic communities are related to discharges which can lead to changes in water (section 6.2.3) and sediment quality (section 6.2.4).

Studies show that the effects of drilling discharges on the benthic fauna communities are minor and nearly always restricted to a zone within about 100 meter of the discharge of water based drilling mud and cuttings /45//46/. There is no evidence of ecologically significant bioaccumulation of metals or petroleum hydrocarbons by benthic fauna residing or deployed in cages near water based drilling mud and cuttings discharges. The lack of bioaccumulation or toxicity of drilling waste components indicates that effects of water based drilling mud cuttings piles are highly localized and will not be exported to the local food web /45//46/. Monitoring at Dan show some effect to benthic fauna within a distance of 750 meters from the platform /6/.

Sedimentation of water-based drilling mud and drill cuttings on the seabed may bury some of the sessile benthic fauna. Changes in the sediment grain size and texture may render the sediment unsuitable for settling and growth of some species, while rendering the substrate more suitable for other, opportunistic species. Organic enrichment, can cause changes in the abundance, species composition and diversity of the benthic community /45/. Grain size in the DAN project area is considered normal, and no signs of organic enrichment has been observed (section 6.2.4).

Discharges of water based drilling mud and cuttings in the water column will shortly increase turbidity and then settle on the seabed. Discharges have been determined to cause impacts at concentrations above 0.5 mg/l, typically restricted to a radius of less than 1 km from the discharge point /46/. Marine benthic invertebrates generally have poor if any visual ability and are unlikely to be adversely affected by suspended matter. However, smothering by settling sediment has direct mechanical effects on epifauna and infauna and may result in the modification of the substratum. Sediment may directly clog the feeding or respiratory apparatuses of suspension feeders. The impact level depends on the grain-size distribution of the settled sediments and on species-specific tolerances to increased rates of sedimentation and accumulation. Water based drilling mud and cuttings have been found to affect the benthos at a thickness of at least 3 mm or more. Such layer thicknesses will normally be confined to a distance of 100-500 m /46/. Mud and cuttings for a Maersk Oil well has been modelled to settle at a thickness of above 1 mm only within 200 m of the discharge (section 6.2.4).

The risk of widespread, long-term impact from the operational discharges on benthic populations is presently considered low /46/. However, a monitoring campaign of the seabed around the Dan platform shows that measurable impacts on the benthic community are limited to the vicinity (750 m) of the discharge point, but likely with a long-term duration /6/.

The impact is assessed to be of small intensity, local extent and long-term duration. The impact on benthic communities from discharges is assessed to be of minor negative overall significance.

6.2.6.5 Presence of structures

The presence of hard substrates in the water column provide a surface that can be colonised by species that are not normally present in soft sediment environments. Structure and pipeline inspection surveys of the area show that no macrophytes are found on the seafloor and that marine growth (e.g. sea anemones, seaweed, soft corals, sea squirts and sponges) is found on the existing structures in the top 15-20 m.

The impact is assessed to be of small intensity, local extent and short-term duration. The impact on benthic communities from presence of structures is assessed to be of minor negative overall significance.

6.2.6.6 Overall assessment

The overall assessment of impacts on benthic communities from planned activities at the DAN project is summarised in Table 6-13.

**Table 6-13 Potential impacts on the benthic communities from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Underwater noise</b>	Small	Local	Short-term	Negligible negative	Low
<b>Physical disturbance on seabed</b>	Small	Local	Medium-term	Minor negative	High
<b>Suspended sediment</b>	Small	Local	Short-term	Negligible negative	High
<b>Discharges</b>	Small	Local	Long-term	Minor negative	High
<b>Presence of structures</b>	Small	Local	Short-term	Minor negative	High

The cumulative impact to benthic communities is not well known. However, little geographical overlap is expected, and it is expected that cumulative impacts on the benthic fauna will be overall minor negative.

6.2.7 Fish

Potential impacts on fish are related to underwater noise, physical disturbance on seabed, discharges, light and physical presence of structures.

Note that impacts to fish eggs and larvae are assessed as part of plankton in section 6.2.5.

#### 6.2.7.1 Underwater noise

The extent to which underwater noise may impact fish is dependent upon a number of factors including the level of noise produced at the source, the frequencies at which the sound is produced, the rate at which sound attenuates (which will vary for different frequencies and environmental conditions), the sensitivities of different species and individuals to different volumes and frequencies of noise. Noise can affect fish in several ways, including:

- Damage to non-auditory tissue
- Damage to auditory tissues (generally sensory hair cells of the ear)
- Hearing loss due to temporary threshold shift
- Masking of communication
- Behaviour (e.g. avoidance)

Fish behaviour in response to noise is not well understood. Sound pressure levels that may deter some species, may attract others. The fish may also freeze and stay in place, leaving it exposed to considerable damages. When the fish swims away, the effects could be minimal or substantial. It may lead to a fish swimming away from an important feeding ground, which is a considerable change in behaviour. The fish might also swim away from an area where it would generally reproduce. If feeding and reproduction continues to be impeded, this could lead to long term effects /46/.

There are several sources of noise emitting from the planned activities (including drilling activities, seismic surveys, construction of new platforms (pile-driving) and vessels.

#### **Underwater noise from seismic**

Noise emitted during seismic survey can have an impact either directly through harmful physiological effects or behavioral effects. The physiological effects will mainly affect younger life stages of fish such as eggs, larvae and fry /81/. These are stages in fish development where the organisms have limited ability to escape. Some injuries do not directly cause lethal conditions, but can indirectly lead to the same fatal conditions via reduced ability to assimilate food, or a change in swimming capacity which makes them more vulnerable in relation to predatory fish. The results of a Norwegian study regarding the influence of air gun shooting on the early life stages of five species of fish showed an increased mortality rate for fish eggs out to approximately 5 meters distance from the air guns /86/.

Behaviour of herring schools exposed to 3D seismic survey was observed. No changes were observed in swimming speed, swimming direction, or school size that could be attributed to the transmitting seismic vessel as it approached from a distance of 27 to 2 km, over a 6 h period /96/. The unexpected lack of a response to the seismic survey was interpreted as a combination of a strong motivation for feeding, a lack of suddenness of the air gun stimulus, and an increased level of habituation to the seismic shooting.

Some findings also indicated harmful effects on the sensory cells of adult fish /92/. The fish were kept in cages and the seismic vessel passed the cages along course lines running from 400-800 m distance at the beginning and up to 5-15 m from the cages. Since the experimental fish were so close to the air guns, one could discuss whether these types of injuries are representative for adult, free-swimming fish.

Another issue is potential disturbances that fish may be exposed to in spawning areas and during migration to the spawning grounds. This can change the areas that are used for spawning, and possibly the timing of the spawning, so that spawning conditions become less favorable. It must also be emphasised that effects must be interpreted in the light of the fact that they will be unique for each species, and that the vulnerability and effect of external stimuli depend on the life stage.

Seismic airguns may affect the behaviour of fish in the area close to the seismic vessel. However it will not lead to long term changes to the size of fish stocks in general /97/. Research has shown that injuries and increased mortality can occur at distances less than 5 m from the air guns, with fish in the early stages of life being most vulnerable. The seismic-created mortality is so low that it is not considered to have any significant negative impact on fish populations /99/.

The impact is assessed to be of small intensity, local extent and short-term duration. The impact on fish from underwater noise from seismic is assessed to be of negligible negative overall significance.

#### **Underwater noise from pile-driving and conductor driving**

There are several examples of pile driving for e.g. a monopile tower as part of a wind farm resulting in the death of fish in the immediate vicinity of the pile driving activity. Typically, mortality is reported within a distance of about 0 to 5 meters from pile driving activity /93/. Fish further away will most likely move away from the sound source. It is estimated that, for example, flounder and cod will respond within a distance of 500 m and 2 km, respectively /95/.

There is a substantial difference between pile driving for such a wind farm monopile and pile and conductor driving for an oil and gas installation. The hammering will usually be done within 2 hours for the pile used for the jacket and 6-8 hours for conductors that will be used for drilling of a new well. The total duration is expected to be short.

The effects on the fish from pile-driving are assessed to be small, local and short-term. It is concluded that the overall impacts will be minor negative. By adopting certain standard mitigation measures such as ramp-up/slow start procedures, where the first hammer blows are at reduced impact energy allowing noise sensitive fish species to escape the immediate vicinity, impact are further reduced.

#### **Underwater noise from other activities**

The sound generated from other activities expected at the DAN project will be lower than for a typical seismic survey. Field studies have shown that some species may be disturbed by noise from passing ships, while others are not affected. It is thus shown that species such as cod and haddock, which often occurs in large schools around offshore platforms, do not respond to noise from passing vessels. Other species tend to move away from a passing vessel. Reaction range varies from 100 - 200 m for many typical vessels but 400 m for noisy ones /77/. The fact that the drilling rigs and offshore platforms attract fish indicates that the noise from drilling, etc. generally does not affect fish /78//79//80//80/.

Observations from the platform and underwater inspections around the DAN installations also confirms the presence of fish schools. The impact is assessed to be of small intensity, local extent and short-term duration. The overall impact on fish from underwater noise from other activities is assessed to be of negligible negative significance.

#### **6.2.7.2 Physical disturbance on seabed**

Potential impact to fish caused by physical disturbance on seabed could be disturbance of demersal fish eggs or habitat fragmentation caused by changes to the seabed sediment.

The physical disturbance on seabed is limited to pipelay, sediment sampling and disturbance of the seabed by cables during 4D seismic surveys and legs from drilling rigs and new platforms, and thus with a very small area. It is estimated that natural processes such as storms and currents will restore the physical appearance of the seabed to its original (pre-impact) state within a few years or less.

The impact is assessed to be of small intensity, local extent and short-term duration. The overall impact on fish from physical disturbance is assessed to be of negligible negative significance.

#### 6.2.7.3 Suspended sediment

The sensitivity of fish to suspended particles varies highly between species and their life stages and depends on sediment composition, concentration and duration of exposure. High levels of suspended sediment for a short period of time may be less of a problem than a lower level that persists longer. Depending on the exposure the severity of impacts may go from change in behaviour, to sub-lethal and lethal effects.

Based on the expected dispersal of suspended sediment during pipelay and other activities for the DAN project (section 6.2.3), any impacts to fish would be limited to a local area surrounding the new pipelines.

In general, it is expected that fish temporarily will leave the areas close to the construction activity. Food sources living in the seabed e.g. polychaetes will become exposed during pipelay/trenching, and fish that are not affected by noise or higher turbidity are likely to be attracted to the construction site /88//89//90/.

Both laboratory and field investigations have showed that pelagic species such as herring and smelt began to flee areas with fine-grained suspended sediment when the concentration reached approximately 10 mg/l and 20 mg/l, respectively /82/. Flatfish are generally highly tolerant to high concentrations of suspended sediment. Studies of plaice with concentrations of 3,000 mg/l showed no effect after a 14-day exposure /83/.

The overall effects on fish from increased turbidity are assessed to be small, local and short-term. It is concluded that the overall impacts on fish and fish stocks from sediment spreading will be minor negative.

#### 6.2.7.4 Discharges

Potential impacts on fish from discharges are related to a number of discharges, which may change the water quality (section 6.2.3).

Modelling of discharges shows that impacts on sensitive fish species can occur up to a distance of 7 km downstream from the platform (section 6.2.3). However, a recent review of environmental impacts of produced water and drilling waste discharges from the Norwegian offshore petroleum industry suggests that the effects of discharges are local, and in general confined to within 1-2 km from an outlet both in the waters and on the seabed, and that the risk of widespread impact from the operational discharges is low /46/.

Studies have shown that compounds present in produced water have a potential to exert endocrine effects in fish. The experimental exposure levels studied cover a range of produced water concentrations that are typically found in close proximity to the discharge points. They might therefore elicit effects on fish standing close to platforms. However, it is concluded that widespread and long lasting effects of produced water on the population level in fish are unlikely /46/.

Modelling results shows that drilling mud generally settles on the seabed within a distance of 12 km downstream of the discharge point, with the majority settling within a few km. Drill cuttings settle within a distance of 200 m (section 6.2.4). Several field studies that have measured the concentration of suspended solids in plumes of drilling mud and cuttings at different distances from the drill rigs have confirmed this pattern. The measurements have shown that the concentration of suspended drill cuttings and mud in the water column drops very quickly due to sedimentation and dilution of the material /45//46/. A monitoring campaign of the seabed around

the Dan F platform shows that measurable impacts on the seabed sediment benthos are limited to the vicinity (750 m) of the discharge point /6/, and any impacts on fish are considered to fall within this range.

Based on the modelling results the type of chemicals discharged the intensity of the impact from discharges is assessed to be small with a local extent and of a medium-term duration. Overall, the impact is assessed to be of minor negative significance.

#### 6.2.7.5 Light

Though safely lights are present at all platforms and vessels only manned platforms (Dan F and Dan B) are illuminated. Platforms may be providing an enhanced foraging environment for larval, juvenile and adult fishes by providing sufficient light to locate and capture prey, as well as by concentrating positively phototactic prey taxa. For juvenile fish there is probably a trade-off between living and foraging in an artificially illuminated nocturnal environment. The increased illumination likely allows them to feed on zooplankton that have concentrated within the light field near the surface; however, the same light may make them more vulnerable to predators.

The potential disturbance to fish of light emissions from rigs, platforms and vessels is expected to be local, extending 90-100 m from the source /98/. As such, any impacts on fish arising from light emissions are considered to be minor and localised to a small proportion of the population. The impact is assessed to be of small intensity, local extent and short-term duration. The overall impact on fish from light at the DAN project is assessed to be of negligible negative significance.

#### 6.2.7.6 Presence of structures

The introduction of a hard substrate into the water column, provide a surface that can be colonised by species that are not normally present in soft sediment environments. Structure and pipeline inspection surveys of the area show that no macrophytes are found on the seafloor and that marine growth (e.g. sea anemones, seaweed, soft corals, sea squirts and sponges) is found on the existing structures in the top 15-20 m.

It is expected that the artificial reef will attract certain species of fish to find hiding places and food in hard bottom areas /148/. The fish are attracted to the boulders with their variety of habitats which creates a wealth of hiding places where e.g. small fish and fry can hide from predators. But also cod and whiting are attracted by the often larger food supply offered by heterogeneous structures such as boulder reefs /149/. Pelagic species are not expected to be affected by the physical presence of the structures.

The impact is assessed to be of small intensity, local extent and short-term duration as all structures are removed during decommissioning. The overall impact on fish from presence of structures is assessed to be of negligible negative significance. However, for some fish species a positive impact due to a local reef effect is expected.

#### 6.2.7.7 Overall assessment

The overall assessment of impacts on fish from planned activities for the DAN project is summarised in Table 6-14.

**Table 6-14 Potential impacts on fish from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Underwater noise</b>	Small	Local	Short-term	Minor negative	Low
<b>Physical disturbance on seabed</b>	Small	Local	Short-term	Negligible negative	Low
<b>Suspended sediment</b>	Small	Local	Short-term	Minor negative	Low
<b>Discharges</b>	Small	Local	Medium-term	Minor negative	High
<b>Light</b>	Small	Local	Short-term	Negligible negative	Low
<b>Presence of structures</b>	Small	Local	Short-term	Negligible negative/Positive	Low

The cumulative impact to fish is not well known. However, little geographical overlap is expected, and it is expected that cumulative impacts will be minor negative.

#### 6.2.8 Marine mammals

Potential impacts on marine mammals are related to underwater noise, physical disturbance, discharges, lights and presence of vessels and platforms.

All species of whales are listed in the habitats directive appendix IV, and special protective measures apply regarding deliberate capture or killing of individuals of these species in the wild; and deterioration or destruction of breeding sites or resting places. The DAN project area is not a known breeding site for whales, and no deliberate capture or killing is foreseen.

##### 6.2.8.1 Marine mammals and underwater noise

Hearing is the primary sense for many marine mammals for detection of prey or predators, communication and navigation in the environment.

Marine mammals are usually defined per functional hearing groups, based on their auditory bandwidth /41/. The hearing groups and auditory bandwidth of mammals in the North Sea are shown in Table 6-15.

**Table 6-15 Functional hearing groups and auditory bandwidth for typical species found at the DAN project /41/.**

Species in the North Sea	Functional hearing group	Auditory bandwidth
<b>Seals (pinnipeds)</b>		
Grey seal, harbour seal	Pinnipeds in water	75 Hz to 22 kHz
<b>Whales (cetaceans)</b>		
Harbour porpoise	High frequency	200 Hz to 180 kHz
White beaked dolphin	Mid-frequency	150 Hz to 160 kHz
Minke whale	Low-frequency	7 Hz to 22 kHz

The effect of underwater noise on marine mammals can generally be divided into four broad categories that largely depend on the individual's proximity to the sound source: Detection, masking, behavioural changes and physical damages /41/. The limits of each zone of impact are not distinct, and there is a large overlap between the zones.

- Detection is where the animals can hear the noise. Detection ranges depend on background noise levels as well as species specific audible threshold profiles.
- Masking is where the noise conceals other sounds, e.g. communication between individuals. The impact to e.g. communication is not well understood
- Behavioural changes are difficult to evaluate. They range from very strong reactions, such as avoidance, to more moderate negative reactions where the animal may orient itself towards the sound or move slowly away. However, the animals' reaction may vary greatly depending on season, behavioural state, age, sex, as well as the intensity, frequency and time structure of the sound causing behavioural changes /41/
- Physical damage to marine mammals relate to damage to the hearing apparatus. Physical damages to the hearing apparatus may lead to permanent changes in the animals' detection threshold (permanent threshold shift, PTS). This can be caused by the destruction of sensory cells in the inner ear, or by metabolic exhaustion of sensory cells, support cells or even auditory nerve cells. Hearing loss can also be temporary (temporary threshold shift, TTS) where the animal will regain its original detection abilities after a recovery period. For PTS and TTS the sound intensity and profile is an important factor for the degree of hearing loss, as is the frequency, the exposure duration, and the length of the recovery time /41/.

In connection with the development of offshore wind in Denmark, an expert working group for marine mammals and underwater noise have recommended thresholds for permanent hearing loss (PTS), temporary hearing loss (TTS) as well as behavioural changes for seals and harbour porpoise in Danish waters /138/. Threshold values are shown in Table 6-16. Threshold values for inflicting impact have been determined from an assessment of available data from the scientific literature, based on laboratory studies of animals. The working group was not able to recommend a threshold value for behavioural effects on seal, as there is very limited evidence on how and when seals react to underwater noise.

**Table 6-16 Values for permanent threshold shift (PTS), temporary threshold shift (TTS) and behavioural effects related to pile driving as recommended by a Danish expert working group /138/. All levels are unweighted SEL.**

Species	Behavioural response (dB re 1 µPa SEL)	TTS (dB re 1 µPa SEL cum)	PTS (dB re 1 µPa SEL cum)
<b>Grey seal and harbour seal</b>	-	176	200
<b>Harbour porpoise</b>	140 (single strike)	≥164	≥183

A recent review /142/ concluded that very few data are available for assessment of impact on other species relevant for Danish waters, primarily white-beaked dolphin and minke whale. Until further data are available, TTS thresholds from bottlenose dolphins are the best available data. These studies have shown TTS induced at sound exposure levels in the range 190-210 dB re 1 µPa<sup>2</sup>s, depending on stimulus frequency and duration. No data is available to base recommendations to define threshold regarding behavioural reactions for white-beaked dolphin and minke whale.

### **Underwater noise from seismic**

The planned seismic data acquisition in over the DAN project includes a 4D seismic with airguns (an area of a few hundred km<sup>2</sup>, with a duration of a few months), borehole seismic surveys with airguns (with a duration of some days) and shallow geophysical surveys with small airguns and electrically generated sources, side scan sonar, single and multi-beam echo sounder (typical area of 1 km<sup>2</sup>, with a duration of around 1 week). Typical noise levels and frequencies for the planned activities at the DAN project are presented in Table 6-11 (section 6.2.5).

The underwater noise levels generated during seismic activities at the DAN project can potentially be above the threshold values established for PTS, TTS and behavioural impacts. The largest noise levels are generated by the sources used for 3D, 4D and other marine seismic surveys.

An impact assessment was undertaken for a similar 4D marine seismic survey in the DUC area /144/. This concluded that:

- The probability of the survey vessel encountering any marine mammals and other marine species is small.
- Impacts on the marine species, if any, will take place at or within 30 metres from the airgun. It was assessed that no marine animals would be exposed to sound levels which could cause PTS, and that only TTS and behavioural impacts would occur.

For the DAN project, the extent of the impact will depend on the final set-up for the seismic survey. The 2012 impact assessment concluded that the effects of a seismic survey was local. As the source level is higher, the potential area where PTS, TTS or behavioural impacts may occur is assessed to be larger. A study of harbour porpoise during a 2D seismic survey in the Moray Firth found that animals showed behavioural response within 5-10 km /151/, while an assessment in the central North Sea found behavioural response to a distance of 20 km /2/. Overall, the impacts on marine mammals may be local (PTS, TTS) or regional (behavioural).

Both PTS, TTS and behavioural impacts are considered of small intensity as there will be partial impacts on individuals within the affected area. The DAN project area is not of particular importance to harbour porpoise, and few individuals are observed (sensitive periods such as breeding periods are presented in section 5.9), and it is assessed that the marine mammal populations in the North Sea will not be impacted due to seismic activities at the DAN project.

Permanent threshold shift (PTS) is considered a potential impact with a long-term duration as the impact is permanent to the affected individuals, and will persist. Temporary threshold shift (TTS) and behavioural impacts are generally considered potential impacts with a short-term duration, as the impact is not permanent. A study of harbour porpoise during a 2D seismic survey in the Moray Firth found that animals showed behavioural response, and were typically detected again within a few hours. Habituation to the underwater noise was also observed /151/. Overall, the potential impacts of seismic activities to marine mammals are considered of medium-term (TTS and behavioral) to long-term (PTS) duration.

The impact is assessed to be of small intensity, local or regional extent and medium or long-term duration. The overall impact on marine mammals from underwater noise from seismic is assessed to be of moderate negative significance. If mitigating measures (section 8.1) are implemented, the impact to marine mammals from seismic surveys can be alleviated.

### **Underwater noise from drilling and pile driving**

The planned drilling activities are associated with the drilling of up to 57 wells (43 new wells and 14 re-drills). Underwater noise is primarily associated with ramming of conductors, with a duration of approximately 6-8 hours, and ramming of piles for new structures.

The impact assessment for drilling activities at the DAN project is largely based on /140/, where underwater sound monitoring was performed for background levels, drilling operations and conductor ramming.

- Underwater drilling sound: The underwater noise from the drilling rig were masked by background sound within 500 -1000 meters from the rig. It was concluded that no harmful effects (threshold shifts or behavioural response) on marine mammals could be expected.
- Ramming of conductors: The noise levels where there is a risk of causing hearing damage to marine mammals is restricted to an area very close to the drilling rig or circumstances of prolonged exposure to continuous sounds, which is very unlikely. However, behavioural effects are most likely to be found within a few kilometres from the rig, and permanent exclusions are not expected.

DCE has recently modelled possible impacts to the North Sea harbour porpoise population by comparing the average simulated porpoise population sizes and dynamics across movement/dispersal models and pile-driving scenarios. Although the results should be considered preliminary, the patterns generated by the current version of the DEPONS model did not suggest any clear, long-lasting effects of pile-driving noise on the average porpoise population size and dynamics in the North Sea /153/.

For drilling and ramming of conductors, the extent of the impact is expected to be local /140/, of small intensity and with a short-term duration. Based on the above, the overall significance of impact caused by noise from drilling activities is assessed to be minor negative.

#### **Underwater noise from production, vessels and associated activities**

Noise will also be present from production and associated activities, and from vessels in the area, with typical noise levels and frequencies presented in Table 6-11 (section 6.2.5). Vessels such as barges and supply ships produce noise with energy content primarily below 1 kHz, as do the rigs and platforms. The marine mammals in the area (harbour porpoise, mid-frequency cetaceans, harbour seals and grey seals) are more sensitive to noise at higher frequencies.

Some of the most trafficked areas in Danish waters are also areas with a very high abundance of harbour porpoises /120/. Year-round presence of porpoise observed around the Maersk Oil platforms show that the animals make a trade-off between the noise from operational activities and likely higher prey abundance /139/. Any displacements of harbour porpoises due to this type of noise are expected to be short term, and over relatively short distances.

The impact is assessed to be of small intensity, local extent and short-term duration. The impact on marine mammals from underwater noise from production and vessels is assessed to be of minor negative overall significance.

#### **6.2.8.2 Discharges**

The main discharges are related to the production (Dan F) and drilling activities (Dan F and four new wellhead platforms near Dan F). There is a risk of affecting the water quality from the current discharges of produced water at Dan F up to 30.8 km to the North East of the outlet /42/. Any potential impacts on marine mammals are thus confined to the regional environment near the platforms and vessels. The risk of bioaccumulation will be species-specific and depend on the type of prey. Fish has been assessed not to bioaccumulate (section 6.2.7), while plankton may incorporate some substances. Maersk Oil expects to discharge only green and yellow which do not bioaccumulate (section 8.1.3).

The impact is assessed to be of small intensity, regional extent and short-term duration. The impact on marine mammals from discharges is assessed to be of minor negative overall significance.

#### **6.2.8.3 Light**

Though safety lights are present at all platforms and vessels, only manned platforms (Dan F and Dan B) are illuminated. Navigational and deck working lights used to illuminate working areas, are sources of artificial light into the environment. Light may locally attract marine mammal prey item such as plankton and fish (section 6.2.5 and 6.2.7).

A recent study at the Dan platform /139/ showed that harbour porpoises near the platform had variable diurnal activity, but a general trend showed higher acoustic activity during the night close to the platform. At further distance from the platform this pattern was not observed. The presence of marine mammals at Maersk Oil platforms indicates that marine mammals do not avoid light.

The impact is assessed to be of small intensity, local extent and short-term duration. The impact on marine mammals from light is assessed to be of negligible negative overall significance.

6.2.8.4 Presence of structures and vessels

Presence of vessels and structures may contribute to the animals' habituation to human activities. Marine mammal responses to vessels often include changes in general activity (e.g., from resting or feeding, to active avoidance), changes in surfacing-respiration-dive cycles, and changes in speed and direction of movement. Behavioural reactions tend to be reduced when animals are actively involved in a specific activity such as feeding or socializing /122/.

In addition to the Dan project existing structures, five new platforms (four new wellhead platforms at Dan F and Kraka, and a new accommodation platform at Dan F) are expected, with connecting with pipelines. Year-round presence of marine mammals observed at the Maersk Oil platforms /139/ shows that the animals are not deterred from the platforms.

The impact of the presence of vessels and structure is assessed to be of small intensity, local extent and short-term duration. The impact on marine mammals from presence of vessels is assessed to be of negligible negative overall significance.

6.2.8.5 Overall assessment

The overall assessment of impacts on marine mammals from planned activities at the DAN project is summarised in Table 6-17.

**Table 6-17 Potential impacts on marine mammals from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Underwater noise from seismic</b>	Small	Local or regional	Medium-term Long-term	Moderate negative	Medium
<b>Underwater noise from drilling</b>	Small	Local	Short-term	Minor negative	Medium
<b>Underwater noise from production, vessels etc</b>	Small	Local	Medium-term	Minor negative	Medium
<b>Discharges</b>	Small	Regional	Medium-term	Minor negative	High
<b>Light</b>	Small	Local	Short-term	Negligible negative	High
<b>Presence of vessels and structures</b>	Small	Local	Short-term	Negligible negative	High

The cumulative impact to marine mammals is not well known. However, little geographical overlap is expected (section 6.2.3 and 6.2.3), and it is expected that cumulative impacts will be minor. Note that marine mammals are frequently observed near the existing platforms /139/.

6.2.9 Seabirds

Seabirds may potentially be impacted by noise, discharges and light.

#### 6.2.9.1 Noise

Noise may negatively affect the seabirds as physical damage or behavioural response.

Little is known about underwater hearing in diving seabirds and information on effects from underwater sound on birds are sparse, but observations from seismic vessels in the Irish Sea also did not reveal any behavioural response of seabirds to seismic survey activities /146/. Birds diving very close (a few meters) to an air gun array, may potentially suffer damage to the auditory system. However, birds have the ability to regenerate the sensory cells in the inner ear and a possible hearing impairment, would thus be temporary.

Due to the highly mobile nature of birds, they are generally not considered to be sensitive to noise from surveys /147/.

The impact is assessed to be of small intensity, local extent and short-term duration. The overall impact on seabirds from noise is assessed to be of negligible negative significance.

#### 6.2.9.2 Discharges

Discharges have been described in section 6.2.3 and are assessed to have a minor negative impact on water quality.

Seabirds may be impacted if they come into contact with the discharges. The impact can include both direct impacts (contact) and indirect impacts (digestion of contaminated organisms), and will depend on the oil or chemicals encountered. Any potential impacts are thus confined to the local environment near the point of discharge.

The impact is assessed to be of small intensity, local extent and short-term duration. The overall impact on seabirds from discharges is assessed to be of minor negative significance.

#### 6.2.9.3 Light

Though safety lights are present at all platforms and vessels only manned platforms are illuminated. Light and illumination may attract seabirds when it is dark or under certain weather conditions. Birds may fly into parts of the infrastructure and get injured, killed or stranded. There is also observations at Maersk Oil showing that the platforms may function as a resting place for seabirds, and rare observations of fatalities. The potential impact is related to individuals, and is not assessed to have an effect on the North Sea population.

The impact is assessed to be of small intensity, local extent and short-term duration. The overall impact on seabirds from light is assessed to be of negligible negative significance.

#### 6.2.9.4 Overall assessment

The overall assessment of impacts on seabirds from planned activities at the DAN project is summarised in Table 6-18.

**Table 6-18 Potential impacts on seabirds from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Noise</b>	Small	Local	Short-term	Negligible negative	High
<b>Discharges</b>	Small	Local	Short-term	Minor negative	Medium
<b>Light</b>	Small	Local	Short-term	Negligible negative	High

### 6.3 Assessment of potential social impacts

Impact assessment for planned activities for each relevant social receptor is presented in the following sections.

#### 6.3.1 Cultural heritage

Potential impacts on cultural heritage relate to physical disturbance.

National authorities have laws and procedures to avoid impacts on cultural heritage from construction projects. Knowledge of cultural heritage in the North Sea is scarce, and surveys are performed prior to construction activities.

##### 6.3.1.1 Physical disturbance

Prior to drilling a site survey will be undertaken in the area around the well location and this will reveal whether any cultural heritage objects are present in the area. In case of a find, the Danish Agency for Culture which is the responsible authority for cultural heritage in Denmark will be contacted for advise. Wrecks that are more than 100 years are protected by the museum law.

At the DAN project, the planned new structures is expected to be constructed near existing platforms where surveys have been carried out. If no previous surveys have been carried out, surveys will be performed prior to construction activities of new structures. The impact from physical disturbance on cultural heritage is assessed to be of no significance.

##### 6.3.1.2 Overall assessment

The overall assessment for impacts on cultural heritage from planned activities is summarised in Table 6-19.

**Table 6-19 Potential impacts on cultural heritage from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Physical disturbance</b>	-	-	-	None	High

#### 6.3.2 Protected areas

Potential impacts on protected areas relate to discharges.

The Natura 2000 sites are assessed in a separate screening (section 10). Other protected areas include nature reserves along the west coast of Jutland, and the UNESCO reserve Wadden Sea.

##### 6.3.2.1 Discharges

As the distance between the DAN project and the Wadden Sea is more than 100 km, and the distance to the nature reserves along the west coast are more than 200 km, no impacts are anticipated from planned activities.

##### 6.3.2.2 Overall assessment

The overall assessment of impacts on protected areas (excluding Natura 2000) from planned activities at the DAN project is summarised in Table 6-20.

**Table 6-20 Potential impacts on protected areas (excluding Natura 2000) from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Discharges</b>	-	-	-	None	High

6.3.3 Marine spatial use

Potential impacts on marine spatial use are related to restricted zones. Note that impacts on fishery is addressed separately.

6.3.3.1 Restricted zones

Safety zones of 500 m surround the existing platforms (no unauthorised vessels permitted), while existing pipelines have a safety zone 200 m on each side (no anchoring and no trawling). These zones around existing structures in the North Sea cause restrictions on ship traffic.

For the DAN project, five new platforms are planned, and thereby new permanent restricted zones of approximately 4 km<sup>2</sup> are expected. New pipelines will also be subject to a safety zone of 200 m on each side. In addition, survey and drilling activities may require for safety reason a limited temporary traffic restriction during the period of seismic acquisition.

As part of a trial project, injection water supply from Dan F through a 9 km temporary pipeline to Kraka. The pipeline will not be buried and the area will be secured from boat anchoring or trawling. Information to the other sea users will be provided through relevant authorities. In addition, Maerks Oil will commissioned a guard ship that will be positioned to alert other approaching sea-user (e.g. fishing boats). Such a pipeline is assessed to be a restricted zone of short-term duration.

Once the DAN project is decommissioned, the structures will be removed. However, as pipelines are left in place and wells are plugged, there may still be limitations for use of the seabed.

The impact is assessed to be of small intensity, local extent and short-term (survey, trial project or drilling) or long-term (platform safety zones) duration. The overall impact on marine spatial use from restricted zones is assessed to be of negligible negative significance.

6.3.3.2 Overall assessment

The overall assessment of impacts on marine spatial use from planned activities at the DAN project is summarised in Table 6-21.

**Table 6-21 Potential impacts on marine spatial use from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Restricted zones</b>	Small	Local	Short-term long-term	Minor negative	High

6.3.4 Fishery

Potential impacts on fishery are related to occupation of seabed, restrictions and an indirect impact in case the target fish species are affected.

6.3.4.1 Physical disturbance on seabed

For the DAN project, five new platforms (four new wellhead platforms at Dan F and Kraka, and a new accomodation platform at Dan F) are planned, with the four wellhead platforms connected with pipelines. In addition, physical disturbance to seabed is related to site surveys and temporary placement of drilling rig legs on the seabed close to the existing platforms Dan F.

The disturbance is expected near existing structures which are already covered by a restriction zone for fishery. Overall, it is assessed that there will be no impacts on fishery.

6.3.4.2 Restricted zones

As assessed in section 6.3.3, there will be five new platforms at the DAN project, with approximately 4 km<sup>2</sup>, of new permanent restricted zones, as well as safety zones for bottom trawling and anchoring near the new pipelines.

In addition, temporary restricted zones may be imposed during survey and drilling activities. Extension of the restricted zone may pose a temporary restriction to fishery during the short period (days-months) the activities occur.

The impact is assessed to be of small intensity, local extent and short-term (survey or drilling) or long-term (platform safety zones) duration. The impact on fishery from restricted zones is assessed to be of minor negative overall significance.

6.3.4.3 Changes to target fish

Potential impacts on fishery could e.g. include seismic surveys resulting in target fish temporarily moving away from the sound source and returning after the activity stops. Impacts on fish have been assessed in section 6.2.7 to be negligible - minor negative. The impact is thus considered of small intensity, local extent and short-long term duration. The impact on fishery from changes to target species is assessed to be of negligible negative overall significance.

6.3.4.4 Overall assessment

The overall assessment of impacts on fishery from planned activities at the DAN project is summarised in Table 6-22.

**Table 6-22 Potential impacts on fishery from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Physical disturbance on seabed</b>	None	-	-	None	High
<b>Restricted zones</b>	Small	Local	Short-term Long-term	Minor negative	High
<b>Changes to target species</b>	None	-	-	Negligible negative	High

6.3.5 Tourism

6.3.5.1 Restricted zones

The planned activities at DAN take place offshore, at a distance of 220 km from shore. Tourism is related to the nearshore (and onshore) areas, and no impacts of restricted zones on tourism are expected.

6.3.5.2 Overall assessment

The overall assessment of impacts on tourism from planned activities at the DAN project is summarised in Table 6-23.

**Table 6-23 Potential impacts on tourism from planned activities at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Restricted zones</b>	None	-	-	None	High

### 6.3.6 Employment and tax revenue

Potential impacts on employment and tax revenue relate to employment at the DAN project.

#### 6.3.6.1 Employment

The offshore oil and gas production is important to Danish economy, as thousands of people are employed in the offshore industry (section 3.4.1 and 5). In particular, the future developments of Maersk oil's activities for the DAN project includes seismic surveys, maintenance of pipelines and structures, drilling of up to 57 new wells, establishment of up to five new platforms, as well as production at the existing facilities at DAN. All these activities will contribute positively to employment.

The impact is assessed to be of medium intensity, local or national extent and medium-term duration. The overall impact on employment from activities at the DAN project is assessed to be of positive significance.

#### 6.3.6.2 Tax revenue

The tax revenue from the DAN project has not been quantified, but the tax revenue to the state of Denmark from oil and gas activities is significant. The state's total revenue is estimated to range from DKK 20 to DKK 25 billion per year for the period from 2014 to 2018 8 (section 3.4.1 and 5). The DAN project is an important contributor to the Danish hydrocarbon production.

The impact is assessed to be of medium intensity, local or national extent and medium-term duration. The overall impact on tax revenue from activities at the DAN project is assessed to be of positive significance.

#### 6.3.6.3 Overall assessment

The overall assessment of impacts on employment from planned activities at the DAN project is summarised in Table 6-24.

**Table 6-24 Potential impacts on employment from planned activities at the DAN project.**

Impact mechanism	Intensity	Extent	Duration	Overall significance	Level of confidence
<b>Employment</b>	Medium	Local/national	Medium term	Positive	Medium
<b>Tax revenue</b>	Medium	Local/national	Medium term	Positive	Medium

### 6.3.7 Oil and gas dependency

#### 6.3.7.1 Dependency

As part of a long-term Danish energy strategy, the oil and gas production is considered instrumental in maintaining high security of supply. Denmark is expected to continue being a net exporter of natural gas up to and including 2025 and with license to operate until 2042 (section 3.4.1 and 5).

If no production is undertaken by Maersk Oil for the DAN project area in the North Sea, there will be no contribution to the Danish economy or security of supply from the DAN project.

The impact is assessed to be of medium intensity, local or national extent and medium-term duration. The overall impact on oil and gas dependency from activities at the DAN project is assessed to be of positive significance.

### 6.3.7.2 Overall assessment

The overall assessment of impacts on oil and gas dependency from planned activities at the DAN project is summarised in Table 6-24.

**Table 6-25 Potential impacts on employment from planned activities at the DAN project.**

Impact mechanism	Intensity	Extent	Duration	Overall significance	Level of confidence
<b>Oil and gas dependency</b>	Medium	Local/national	Medium term	Positive	Medium

## 6.4 Summary

The potential impacts on environmental and social receptors from planned activities at the DAN project are summarised in Table 6-26. The impact with the largest overall significance is provided for each receptor.

**Table 6-26 Summary of potential impacts on environmental and social receptors from planned activities at the DAN project. The impact with the largest overall significance is provided for each receptor.**

Receptor	Worst case potential impact
<b>Hydrography</b>	Minor negative
<b>Climate and air quality</b>	Moderate negative
<b>Water quality</b>	Minor negative
<b>Sediment type and quality</b>	Minor negative
<b>Plankton</b>	Minor negative
<b>Benthic communities</b>	Minor negative
<b>Fish</b>	Minor negative
<b>Marine mammals</b>	Moderate negative
<b>Seabirds</b>	Minor negative
<b>Cultural heritage</b>	None
<b>Protected areas (excluding Natura 2000)</b>	None
<b>Marine spatial use</b>	Minor negative
<b>Fishery</b>	Minor negative
<b>Tourism</b>	None
<b>Employment and tax revenue</b>	Positive
<b>Oil and gas dependency</b>	Positive

## 7. IMPACT ASSESSMENT: ACCIDENTAL EVENTS

### 7.1 Impact mechanisms and relevant receptors

#### 7.1.1 Potential impact mechanisms

Potential impact mechanisms associated to the accidental events for the DAN project are screened based on the project description (section 3) and the technical sections (appendix 1).

Potential impact mechanisms include:

- Minor accidental events (gas release, spill of chemical, diesel or oil)
- Major accidental events (oil spill or gas release)

The source of the potential impact mechanisms is provided in Table 7-1.

**Table 7-1 Sources of potential impact mechanisms for the DAN project. "X" marks relevance, while "0" marks no relevance.**

Potential impact mechanism	Sesimic	Pipelines & structures	Production	Drilling	Well stimulation	Transport	Decommissioning
<b>Minor accidental events</b> (gas, chemical, diesel or oil)	X	X	X	X	X	X	X
<b>Major accidental events</b> (oil or gas)	0	0	X	X	X	0	0

Figure Figure 7-1 presents an overview of the accidental spills over Maersk oil facilities from the period 2010 to 2014.

The number of yearly reported spills ranged from 15 to 94 from 2010-2014 and on average were less than 100 litres. During 2014, there were two large diesel spills at Harald and on an accommodation rig which contributed to an increase in the volume of oil and diesel spill. In 2013 and 2014, methanol spills at Tyra and Harald contributed to more than three quarter of the total volume of chemical spills during those years. Methanol is classified as a green chemical (section 8.1.3).

Action has been taken to eliminate the risk of such spills occurring again: replacing parts of a pump and reinforcing the need to take the utmost care when bunkering diesel.

Since 2011, all accidental discharges of oil and chemicals, regardless of volume, are reported. During 2014, the company introduced a more systematic way of reporting spills which may partly contribute to the observed increase in the number of spills being reported.




Figure 7-1 Accidental oil, diesel and chemical spills from Maersk Oil platforms in the North Sea.

7.1.2 Relevant receptors (environmental and social)

The environmental and social receptors described in the baseline are listed below.

- Environmental receptors: Climate and air quality, hydrographic conditions, water quality, sediment type and quality, plankton, benthic communities (flora and fauna), fish, marine mammals, seabirds
- Social receptors: Cultural heritage, protected areas, marine spatial use, fishery, tourism, employment, tax revenue, oil & gas dependency

The relevant receptors have been assessed based on the project description (section 3) and the potential impact mechanisms (section 7.1). Relevant receptors are summarized in Table 7-2.

**Table 7-2 Relevant receptors for the impact assessment of accidental events for the DAN project. "X" marks relevance, while "0" marks no relevance.**

Potential impact mechanism – accidental events	Environmental Receptors									Social Receptors							
	Climate & air quality	Hydrographic conditions	Water quality	Sediment type and quality	Plankton	Benthic communities	Fish	Marine mammals	Seabirds	Cultural heritage	Protected areas	Marine spatial use	Fishery	Tourism	Employment	Tax revenue	O&G dependency
<b>Gas release</b>	X	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>Chemical spill*</b>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>Oil spill</b>	0	0	X	X	X	X	X	X	X	X	X	X	X	X	0	0	0

\*a worst case chemical spill is very local, and not assessed further.

7.1.3 Marine strategy frameworks directive - descriptors

The list of receptors and impact mechanisms described in the ESIS can be directly related to the descriptors set within the Marine Strategy Framework Directive (MSFD; section 2.1.5). The MSFD outlines 11 descriptors used to assess the good environmental status of the marine environment (see presentation of descriptors in section 6.1.3).

The receptors identified in the ESIS are related to the MSFD status indicators hydrography (D7), harbour porpoise and benthic communities (D1, D6). The impact mechanisms for accidental events in the ESIS are related to the MSFD pressure indicators discharges (D6, D8, D9). Each impact mechanism is further assessed for the relevant receptors in the following sections 7.2 and 7.3.

7.1.4 Minor accidental events

A minor accidental event is a spill where the spilled volume is finite.

Minor spill could be chemical or diesel, and occur following e.g. vessel collision, pipeline leakage or rupture of a chemical container. Statistical analysis shows that collisions between vessels, platforms, riser etc. are very unlikely, typically in the range of  $1.4 \cdot 10^{-7}$  to  $6.5 \cdot 10^{-4}$  per year. Minor gas release of several m<sup>3</sup> may also occur during venting.

7.1.4.1 Minor chemical spill (rupture of chemical container)

A chemical spill was modelled for biocide at the DONG operated Hejre platform /43/. The spill was defined for loss of biocide from a container, which was considered worst case regarding potential impact. The modelled spill was for 4,500 l of biocide to the sea. Results showed that the distance, to which impacts may occur (PEC/PNEC of 1), was 500 m /43/. A minor chemical spill is thus very confined, with impacts within 500 m. A minor chemical spill is not assessed further.

7.1.4.2 Minor oil spill (vessel collision)

A diesel spill following a vessel collision has been modelled for a spill of 1,000 m<sup>3</sup> marine diesel during 1 hour /5//25/. The modelling results show that no shoreline impact occurs, and impacts are only expected in the local area. Most of the oil is expected to evaporate or submerge into the water column after 7 days, and by day 20 all of the released oil is no longer mobile; it has evaporated or biodegraded /5//25/.

7.1.4.3 Minor oil spill (full pipeline rupture)

A full rupture of a pipeline at the DAN project in a worst case scenario is a rupture of pipeline from Dan FB to Gorm C. Emergency valves will automatically close to isolate the pipeline, and the expected maximum volume from a ruptures pipeline is a spill of 2,200 m<sup>3</sup> multiphase (oil/water/gas).

A full bore pipeline rupture has been modelled for an oil spill of 10,000 barrels (about 1,600 m<sup>3</sup>) over 1 hour at the TYE to Gorm midpoint /152/. The results show that the oil will spread locally (Figure 7-2), and that it is unlikely that the oil will cross a marine border. The results show no risk of any shoreline being impacted by oil.




Figure 7-2 Probability that a surface a 1 km cell could be impacted by oil in case of full pipeline rupture /152/.

An oil spill from a pipeline rupture was also modelled for the DONG operated Hejre platform /43/. The modelling showed that the dispersion of the spill is local near the rupture. It is expected that the oil from a pipeline rupture will rise to the surface where a large part will evaporate. Following evaporation, the oil becomes heavier and more viscous and sinks to the seabed. The fate of oil was not modelled /43/.

#### 7.1.5 Major accidental events

A major spill results from an uncontrolled loss of a large volume of oil which often require intervention to be stopped. The main source of major spill is related to blow out events. Blow out events are highly unlikely and may occur during the drilling and completion phase or any operational phase of a well. Well blowout and well release frequencies are in the range (lowest frequency blow out – highest frequency well release)  $7.5 \times 10^{-6}$  to  $3.3 \times 10^{-4}$  per year in maintenance and operation. For development the frequencies are in the range  $3.8 \times 10^{-5}$  to  $6.6 \times 10^{-3}$  per well. As most reservoir contains a mixture of oil and gas, the blow out may results in an oil spill and a gas release. Gas will ultimately be dispersed into the atmosphere, whereas oil fate is more difficult to predict.


When the oil is spilled it goes through physical processes such as evaporation, spreading, dispersion in the water column and sedimentation to the seafloor. Eventually, the oil will be eliminated from the marine environment through biodegradation. The rate and importance of these processes will depend on the type and quantity of the oil as well as the prevailing weather and hydrodynamic conditions. Models are used to predict the fate of oil spill and assess the potential impact on relevant environmental and social receptors.

Oils are classified following the ITOPF classification to allow a prediction of their likely behaviour /158/. Group 1 oils (API>45) tend to dissipate completely through evaporation, whereas group 2 (API: 35-45) and group 3 (API: 17.5-35) can loose up to 40% volume through evaporation but tend to form emulsion. Group 4 oils (API< 17.5) are highly viscous and do not tend to evaporate and disperse. Group 4 is the most persistent oil type. For the DAN project, the oil is relatively light with an API of 27 (Type 3).

The maximum expected initial blow out flow rates from existing producing wells at the DAN project are 23,000 barrel of oil per day (about 3,700 m<sup>3</sup> per day) at Dan F, Dan B and Dan E, and 4,000 barrel of oil per day (about 650 m<sup>3</sup> per day) at Kraka. Regnar is not producing. (Table 7-3).

The oil spill model was done using the Oil Spill Contingency and Response (OSCAR) model. OSCAR is a 3D modelling tool developed by SINTEF, able to predict the movement and fate of oil both on the surface and throughout the water column /5//25//26//27//152/. The model simulates more than 150 trajectories under a wide range of weather and hydrodynamic conditions representative of the DAN area. The model prepares statistical maps based on the simulations that defines the areas most at risk to be impacted by an oil spill. Modelling is performed on the non-ignited spill without any oil spill response (e.g. mechanical recovery; section 8 and 9).

For the DAN project, oil spill modelling results are based on the most credible worst case oil spill scenario at Halfdan (Figure 7-3). The results from the model at Halfdan are considered to be representative of the DAN project due to similarity in oil type (ITOPF Group 3), expected release rate (23,000 barrel of oil per day) and the location (the HALFDAN and DAN platforms are located within 7 km from each other).



**Figure 7-3 Location of two Maersk Oil production wells Svend and Halfdan (and a pipeline midpoint), for which oil spill modelling has been undertaken. Halfdan is located close to the DAN project, and modelling for Halfdan are considered representative to the DAN project. The HALFDAN and DAN platforms are located within 7 km from each other**

The modelled exploration scenarios correspond to a continuous release for 90 days with a flow rate of 23,000 barrels of oil per day (about 3,700 m<sup>3</sup> per day) for ITOPF Group 3 oil (Halfdan) /152/. The duration of the modelled blowouts is based on the fact that the casing of a production well is designed to prevent the collapse of the well and a relief well may be necessary to stop the blow out (such intervention may require about 90 days).

For each spill scenario, the trajectory resulting in the most oiling onshore are considered to be the worst case, and were extracted to investigate the oil's behaviour in more detail /152/. The model set-up and results are summarized in Table 7-3, with selected results of the spill modelling shown in the following sections.

**Table 7-3 Results from the worst credible case scenarios for a well blowout at Halfdan /152/. Note that the modelling is performed without any mitigating measures.**

Parameter	Halfdan
<b>Model set-up</b>	
Release rate	23,000 bbls/d
Release period	90 days
Total mass spilled	288,725 MT
Model run	118 days
<b>Probability of reaching shore</b>	
% of simulations reaching shore	100 %
<b>Time to reach shore (days)</b>	
Min	13 days
Max	83 days
Mean	35
<b>Minimum arrival time to shore (days)</b>	
Denmark	13 days
Sweden	58 days
Germany	Does not impact
Norway	44 days
<b>Fate of oil at end of simulation (MT/%)</b>	
Onshore	4,235 MT (1 %)
Surface	86 MT (0.1 %)
Water column	1,150 MT (0.1 %)
Evaporated	107,100 MT (37 %)
Sedimentation	158,800 MT (55 %)
Biodegraded	17,270 MT (6 %)

#### 7.1.5.1 Halfdan spill modelling

Selected results of the spill modelling for Halfdan are presented in the following /152/:

- Figure 7-4. Danish and German surface waters are likely to be affected by oil. There is also a probability of Dutch, UK and Norwegian waters being affected.
- Figure 7-5. Danish and German water (water column) are likely to be affected by oil. There is also a probability of Dutch, UK and Norwegian waters being affected.
- Figure 7-6. There is 95 % risk of impact to Danish shoreline. Norwegian and Dutch shorelines could also be affected.
- Figure 7-7. The maximum time-averaged concentrations in the water column could reach 800 ppb, but is more likely to be less than 150 ppb.




Figure 7-4 Probability that a surface cell could be impacted in a surface blowout at Halfdan well /152/.

Note that these images DO NOT show the actual footprint of an oil spill, they present a statistical picture based on 159 independently simulated trajectories.





Figure 7-5 Probability that a water column cell could be impacted in a surface blowout at Halfdan well /152/.

Note that these images DO NOT show the actual footprint of an oil spill, they present a statistical picture based on 159 independently simulated trajectories.



Scenario	Threshold
S01	Thickness: 0.001 tonnes/km

Figure 7-6 Probability that a shoreline cell could be impacted in a surface blowout at Halfdan well /152/.

Note that these images DO NOT show the actual footprint of an oil spill, they present a statistical picture based on 159 independently simulated trajectories.




Figure 7-7 Maximum time-averaged total oil concentration for a surface blowout at Halfdan well /152/.

Note that these images DO NOT show the actual footprint of an oil spill, they present a statistical picture based on 159 independently simulated trajectories.

## 7.2 Assessment of potential environmental impacts

Impact assessment for the relevant environmental receptors is presented in this section for accidental events. The assessment is based on modelling data to evaluate the extent, while literature data is applied to assess the intensity and duration of impact.

### 7.2.1 Climate and air quality

Potential impacts on climate and air quality from accidental events are related to gas release.

#### 7.2.1.1 Major gas release

Natural gas is primarily composed of methane, but also often contains related organic compounds, as well as carbon dioxide, hydrogen sulfide, and other components. In case of an uncontrolled gas release, gas will be released to the atmosphere, if the gas is not ignited. Methane is a greenhouse gas and is known to influence the climate with a warming effect (see section 6.2.1).

The impact to climate and air quality from an uncontrolled gas release at the DAN project is assessed to be of medium intensity, with a transboundary extent and a short term duration. The overall significance is assessed to be moderate negative.

#### 7.2.1.2 Overall assessment

The potential impacts are summarised in Table 7-4.

**Table 7-4 Potential impacts on climate and air quality related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
Major gas release	Medium	Transboundary	Short-term	Moderate negative	Low

### 7.2.2 Water quality

Potential impact mechanisms from accidental spill on the water quality is separated into minor and major oil spills.

#### 7.2.2.1 Minor spill

Modelling results for a marine diesel spill from a vessel show that after 20 days all of the released oil is no longer mobile; it has evaporated or is biodegraded (section 7.1.4). Modelling results for a pipeline rupture (ITOPF group 3) show that the dispersion is local near the rupture.

The physical presence of an oil slick may change the physical and chemical parameters of the water by reducing light or oxygen levels. In addition, the increased concentration of oil substances (THC, PAH etc) will alter the water quality.

Based on the modelling results the extent of the impact on the water quality is assessed to be local. The intensity is considered small with a short-term duration, as the oil will evaporate, settle or biodegrade. Overall, the impact on the water quality from an oil spill will be of minor negative significance.

#### 7.2.2.2 Major oil spill

Based on the modelling of a major oil spill (section 7.1.5) oil components concentrations are likely to reach over 800 ppb around the release site, while concentrations are likely to be generally below 150 ppb in the water column. At the end of the model simulation, most of the oil has either drifted onshore, evaporated, sedimented or biodegraded (section 7.1.5).

The physical presence of a large oil slick will cause considerable changes to physical and chemical parameters of marine water quality, such as reduced light or oxygen levels. In addition, the increased concentration of oil substances (THC, PAH etc) will alter the water quality. The extent of the impact depends to a large extent on the prevailing meteorological conditions.

Based on the modelling results the impact is assessed to be of medium intensity, transboundary extent and a medium duration. Overall, the impact on water quality from a major oil spill will be of moderate negative significance.

### 7.2.2.3 Overall assessment

The potential impacts are summarised in Table 7.5.

**Table 7.5 Potential impacts on water quality related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
Minor oil spill	Small	Regional	Short-term	Minor negative	Medium
Major oil spill	Medium	Transboundary	Medium-term	Moderate negative	Medium

### 7.2.3 Sediment type and quality

Potential impact mechanisms to sediment type and quality are related to minor and major oil spill.

#### 7.2.3.1 Minor oil spill

Modelling results for a marine diesel spill from a vessel show that after 20 days all of the released oil is no longer mobile; it has evaporated or biodegraded (section 7.1.4). Modelling results for a pipeline rupture show that the dispersion is local near the rupture.

Based on the modelling results the intensity of the impact is assessed to be small with a potential regional extent and a medium-term duration. Overall, the impact on sediment type and quality from a minor oil spill will be of minor negative significance.

#### 7.2.3.2 Major spill

Based on the modelling of a major oil spill, significant impacts on the sediment type and quality may occur. Modelling shows that around half of the oil will end up on the seabed, corresponding to up to 150,000 MT over a large area in the North Sea. The rest will either drift onshore, evaporate or biodegrade (section 7.1.5).

Full recovery will require degradation or burial of contaminants in combination with naturally slow successional processes. Oil degradation in the marine environment is limited by temperature, nutrient availability (especially nitrogen and phosphorous), biodegradability of the petroleum hydrocarbons, presence of organic carbon, and the presence of microorganisms with oil degrading enzymes /123//124/.

Based on the modelling results the intensity of the impact from a major oil spill is assessed to be medium with a transboundary extent and a medium duration. Overall, the impact on the sediment type and quality will be of moderate negative significance.

### 7.2.3.3 Overall assessment

The potential impacts are summarised in Table 7.6.

**Table 7.6 Potential impacts on sediment type and quality related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	Small	Regional	Medium-term	Minor negative	Medium
<b>Major oil spill</b>	Medium	Transboundary	Medium-term	Moderate negative	Medium

### 7.2.4 Plankton

Potential impact mechanisms to plankton are related to minor and major oil spill.

#### 7.2.4.1 Minor oil spill

Based on the assessed impact to the water quality (section 7.2.2) a minor oil spill is assessed to have a limited impact on plankton community. Though planktonic organisms may be affected, the high reproductive potential of plankton is considered able to compensate.

The intensity of the impact is assessed to be small with a local extent and a short-term duration. Overall, the impact on plankton is assessed to be of minor negative significance.

#### 7.2.4.2 Major oil spill

Laboratory toxicity studies have demonstrated great variation amongst planktonic organisms in response to the effects of spilled oil, with phytoplankton generally considered less sensitive to effects than zooplankton /125/.

Off the coast of western Scotland, tests in containers of naturally occurring algae and water soluble fraction of North Sea oil concentrations of 0.1 mg/l (=100 ppb) showed no significant effects on the total primary production /126/. Toxic effects including decreases in growth rate and inhibition of photosynthesis have been observed in phytoplankton exposed to water soluble fractions of oil concentrations ranging from 1,000 ppb to 10,000 ppb /127/.

Acute lethal effects to zooplankton have been observed from contact with water soluble fractions in concentrations greater than 200 ppb /125/. Sub-lethal effects to zooplankton, including physiological, biochemical and behavioural effects have been observed at one-tenth of lethal concentrations /125/. However, such laboratory toxicity studies have been shown to be of little relevance for predicting long-term effects on natural populations. Such studies are typically short-term and use robust, easily handled species not representative of the wide variety of planktonic organisms that exist naturally. Although such experiments demonstrate oil spill effects to plankton, field observations have typically showed minimal or transient effects /125/.

There are no examples of long-term effects on plankton stocks after oil spills. This is due to plankton reproductive capacity and the water circulation bringing new plankton from outside the affected area /128//129/. Plankton populations are thus not particularly vulnerable to oil spill, and may compensate for any impact through a high reproductive potential.

Based on the assessed impact to the water quality (section 7.2.2.2) the duration of the impact on plankton is short-term. The intensity of the impact is assessed to be medium with a transboundary extent and a short-term duration. Overall, the impact of major oil spill on the plankton community will be of minor negative significance.

### 7.2.4.3 Overall assessment

The potential impacts are summarised in Table 7.7.

**Table 7.7 Potential impacts on plankton related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	Small	Local	Short-term	Minor negative	Medium
<b>Major oil spill</b>	Small	Transboundary	Short-term	Minor negative	Medium

### 7.2.5 Benthic communities

Potential impact mechanisms on the benthic communities are related to minor and major oil spill.

#### 7.2.5.1 Minor oil spill

Based on the assessed impact to the sediment type and quality (section 7.2.3) any significant impacts on the benthic communities are estimated to be limited. The intensity of the impact is assessed to be none/small with a regional extent and a medium-term duration. Overall, the impact on sediment type and quality from a minor oil spill will be of minor negative significance.

#### 7.2.5.2 Major oil spill

Lethal and sub-lethal effects to the benthos may include mortality, alterations in recruitment, growth and reproduction, as well as changes in community structure, including species richness. Nonselective deposit feeders such as polychaetes and nematodes have demonstrated resilience to the adverse effects of spilled oil /130/. Conversely, the density of crustaceans such as amphipods and copepods would be expected to decline due to their known sensitivity to the effects of oil /130/.

The biological effects of oil on the seabed and benthos depend largely on the fate of the spilled oil and the additive toxicity of aromatic hydrocarbons.

Model calculations show that around half of the oil will end at the seafloor, i.e. a large mass over a large area (section 7.1.5). It cannot be excluded that oil components could affect bottom fauna to some extent in the affected area. Recovery of soft-bottom benthos after previous shallow-water oil spills has been documented to take years to decades /123//124/.

The intensity of the impact is assessed to be medium with an transboundary extent and a medium-term duration. In conclusion, the impact on the benthic community from a major oil spill will be of major negative overall significance.

#### 7.2.5.3 Overall assessment

The potential impacts are summarised in Table 7.8.

**Table 7.8 Potential impacts on benthic communities related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	None/Small	Regional	Medium-term	Minor negative	Medium
<b>Major oil spill</b>	Medium	Transboundary	Long-term	Major negative	Medium

## 7.2.6 Fish

Potential impact mechanisms on fish are related to minor and major oil spill.

### 7.2.6.1 Minor oil spill

Based on the assessed impact to the water quality (section 7.2.2) and the sediment type and quality (section 7.2.3) a minor oil spill is assessed to have a limited impact on the fish communities. The impact of a minor oil spill is confined to impacts on individuals, and not populations. The intensity of the impact is assessed to be small with a regional extent and a short/medium-term duration. Overall, the impact on fish from a minor oil spill is assessed to be of minor negative significance.

### 7.2.6.2 Major oil spill

Although laboratory studies have shown a range of lethal and sub-lethal effects of oil on fish /131/ the hydrocarbon concentrations at which these have occurred have generally been considerably higher than those occurring during oil spills /125/. Fish appear to be more sensitive to short-term acute toxicity from the lighter aromatic components, which is probably because they possess the enzymes necessary to metabolise sub-lethal concentrations of hydrocarbons /125//131/.

Laboratory studies have shown that adult fish are able to detect oil in water at very low concentrations, and large numbers of dead fish have rarely been reported after oil spills /132/ /133/. This suggests that juvenile and adult fish are capable of avoiding water contaminated with high concentrations of oil.

Fish are most susceptible to the effects of spilled oil in their early life stages, particularly during egg and planktonic larval stages, which can become entrained in spilled oil. Contact with oil droplets can mechanically damage feeding and breathing apparatus of embryos and larvae /134/. The toxic compounds of oil in water can result in genetic damage, physical deformities and altered developmental timing for larvae and eggs exposed to even low concentrations over prolonged timeframes (days to weeks) /134/. More subtle, chronic effects on the life history of fish as a result of exposure of early life stages to oil include disruption of complex behaviours such as predator avoidance, reproductive and social behaviour /132/. Prolonged exposure of eggs and larvae to weathered concentrations of oil in water has also been shown to cause immunosuppression and allows expression of viral diseases /132/. However, the effect of an oil spill on a population of fish in an area with fish larvae and/or eggs, and the extent to which any of the adverse impacts may occur, depends greatly on prevailing oceanographic and ecological conditions at the time of the spill and its contact with fish eggs or larvae.

Concentrations of 100 ppb THC (total hydrocarbons) has been found to cause acute death of fish eggs and larvae /135/. According to the model results, concentrations of 150-300 ppb in the water column can be found with high likelihood out to a distance of 25 km. At this concentration, the eggs and larvae of fish are likely to be affected. Lethal concentrations can also occur further away from the point of discharge (25-250 km) the likelihood is, however, relatively small (1-25%).

Based on the modelling results, and the above information, the impact is assessed to be of medium intensity with an transboundary extent and a short to medium-term duration. Overall, the impact on the fish community from a major oil spill will be of major negative significance.

### 7.2.6.3 Overall assessment

The potential impacts on fish are summarised in Table 7.9.

**Table 7.9 Potential impacts on fish related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	Small	Regional	Short/medium-term	Minor negative	Medium
<b>Major oil spill</b>	Medium	Transboundary	Short/medium-term	Major negative	Medium

### 7.2.7 Marine mammals

Potential impact mechanisms to marine mammals are related to minor and major oil spill.

#### 7.2.7.1 Minor oil spill

Oil spill spill from collisions or pipeline rupture may impact marine mammals which come into contact with the spill. Marine mammals generally avoid oil slicks, but impacts on individuals may occur through ingestion, inhalation or consumption of contaminated organisms. The extent of a minor oil spill is local (section 7.1.4). The intensity of the impacts is assessed to be small with a short-term duration. The overall significance of impacts on marine mammals at the DAN project is assessed to be minor negative.

#### 7.2.7.2 Major oil spill

A major oil spill may impact marine mammals which come into contact with the spill. Impacts are related to direct contact with the oil, where smothering of seals may occur leading to inflammation, infection, suffocation, hypothermia and reduced buoyancy /25/. Whales and dolphins do not have hair, and are not susceptible to smothering. Both whales and seals may accumulate toxins through ingestion (which can lead to digestive complications), inhalation (which can lead to respiratory damage, paralysis, death) or consumption of contaminated marine organisms.

The sensitive months for marine mammals in relation to a major oil spill have been determined based on the months where the species are present the North Sea /25/. Grey seal, harbour seal and harbour porpoise are sensitive year-round, while minke whale and white-beaked dolping are sensitive in summer (May-September).

Modelling results show that oil may impact both Danish, Swedish, German, Dutch, UK or Norwegian sectors of the North Sea, and the extent is thus considered transboundary. The intensity of the impact is considered to be large, as there may be an impact to the individuals, and also to populations.

Seals can also loose their shoreline habitat if oil washes up on their haul-out sites. Oil spill modelling has identified Denmark, Sweden and Norway as most vulnerable to oil beaching, although Germany, UK and the Netherlands could also be affected.

The intensity of the impact from a major oil spill is large, and may affect the ecosystem structure of marine mammals in the North Sea. The duration of the impact is long-term, and the overall significance on marine mammals from a major oil spill is assessed to be major negative.

### 7.2.7.3 Overall assessment

The potential impacts are summarised in Table 7-10.

**Table 7-10 Potential impacts on marine mammals related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	Small	Local	Short-term	Minor negative	Medium
<b>Major oil spill</b>	Large	Transboundary	Long-term	Major negative	High

### 7.2.8 Seabirds

Potential impact mechanisms to seabirds are related to minor or major oil spill.

#### 7.2.8.1 Minor oil spill

A minor oil spill may impact seabirds, if they come into contact with the oil (see description of vulnerability below). The extent of a minor oil spill is considered local, and of medium-term duration. The intensity is considered small, as the impact of a minor oil spill will affect individuals and not populations. The overall significance of impacts on seabirds from a minor oil spill is assessed to be moderate negative.

#### 7.2.8.2 Major oil spill

Seabirds are very vulnerable to oil spills in the marine environment. Oil may destroy the insulating and water-resistant properties and affecting the buoyancy of the plumage causing the bird to die from hypothermia, starvation or drowning. In addition, birds may get intoxicated from ingestion or inhalation of fuels when they are cleaning their plumages or are feeding on contaminated food. Intoxication may cause irritation of the digestive organs, damages to liver, kidneys and salt glands and leading to anaemia. The intensity of the impacts is therefore assessed to be large /25/.

Birds tend to nest in late spring and summer, which means juveniles are most vulnerable to oil spills in the spring and summer months, although adults of many species may be found in the North Sea all year. The vulnerability for migratory birds depends on the season of the spill, and whether the birds summer or winter along the North Sea coasts.

A major oil spill is assessed to have a transboundary extent and long-term duration. The overall significance of impacts on seabirds from a major oil spill is assessed to be major negative.

#### 7.2.8.3 Overall assessment

The potential impacts are summarised in Table 7-11.

**Table 7-11 Potential impacts on seabirds related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	Large	Local-Regional	Medium-term	Moderate negative	Medium
<b>Major oil spill</b>	Large	Transboundary	Long-term	Major negative	High

### 7.3 Assessment of potential social impacts

Impact assessment for the relevant social receptors is presented in this section for accidental events. The assessment is based on modelling data to evaluate the extent, while literature data is applied to assess the intensity and duration of impact.

#### 7.3.1 Cultural heritage

Potential impacts on cultural heritage are related to oil spill.

Cultural heritage as wrecks or submerged settlements can be impacted by smothering of oil in connection with minor or major oil spills.

The impact will depend on the type of cultural heritage, and the type of oil spilled. The intensity of potential impacts is assessed to be medium, with a transboundary extent and medium-term duration. The overall significance of impacts on cultural heritage from oil spill at the DAN project is assessed to be moderate negative.

##### 7.3.1.1 Overall assessment

The potential impacts are summarised in Table 7-12.

**Table 7-12 Potential impacts on cultural heritage related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Oil spill</b>	Medium	National	Medium-term	Moderate negative	Low

#### 7.3.2 Protected areas

Potential impact mechanisms are related to minor or major spill. Potential impacts on protected areas concerns nature reserves along the west coast of Jutland, and the UNESCO reserve Wadden Sea.

##### 7.3.2.1 Minor oil spill

A chemical spill and an minor oil spill are all events which are considered of local extent, based on the presented modelling (section 7.1). As the DAN project activities are located 200 km from shore, minor oil spills are assessed to have no impact on protected areas.

##### 7.3.2.2 Major oil spill

Major oil spill has been modelled (section 7.1). The potentially impacted area include the Wadden Sea and the nature reserves along the west coast of Jutland. The intensity of the impacts is assessed to be large, with transboundary extent and long-term duration. The overall significance of impacts on protected areas from major oil spill is assessed to be major negative.

##### 7.3.2.3 Overall assessment

The potential impacts are summarised in Table 7-13.

**Table 7-13 Potential impacts on protected areas related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	None	Local	Short-term	Negligible negative	High
<b>Major oil spill</b>	Large	Transboundary	Long-term	Major negative	Medium

### 7.3.3 Marine spatial use

Potential impact mechanisms are related to minor and major oil spill and major gas release.

#### 7.3.3.1 Minor spill

Minor oil spill from e.g. collisions will impact ship traffic in terms of risk of fire, contamination of vessels and restriction areas where emergency handling is taken place. The intensity of the impacts is assessed to be small, with national extent and short-term duration. The overall significance of impacts on ship traffic from minor oil spill is assessed to be minor negative.

#### 7.3.3.2 Major oil spill

A major oil spill is assessed to impact ship traffic as risk of fire and contamination of vessels and as restriction areas where ship traffic is prohibited due to emergency handling. The impact will have a medium intensity with transboundary extent and medium-term duration. The overall significance of impacts on ship traffic from minor oil spill is assessed to be moderate negative.

#### 7.3.3.3 Major gas release

An uncontrolled gas release will likely impact the ship traffic indirectly as spatial restrictions in connection with safety distance to blow out point and danger of fire. The impact is assessed to be of medium intensity, transboundary extent and short term. The overall significance of impacts on ship traffic from major gas release at the DAN project is assessed to be minor negative.

#### 7.3.3.4 Overall assessment

The overall assessment of impacts on ship traffic from accidental events at the DAN project is summarised in Table 7-14.

**Table 7-14 Potential impacts on marine spatial use related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Minor oil spill</b>	Small	National	Short-term	Minor negative	Medium
<b>Major oil spill</b>	Medium	Transboundary	Medium-term	Moderate negative	Medium
<b>Major gas release</b>	Medium	Transboundary	Short-term	Minor negative	Medium

### 7.3.4 Fishery

Potential impact mechanisms related to oil spill and gas release.

#### 7.3.4.1 Major gas release

An uncontrolled gas release will likely impact the ship traffic indirectly as spatial restrictions in connection with safety distance to blow out point and danger of fire. The impact is assessed to be of medium intensity, transboundary extent and short term. The overall significance of impacts on fishery from gas release at DAN is assessed to be minor negative.

#### 7.3.4.2 Major oil spill

A major oil spill may impact fishery in terms of risk of contamination of vessels and gear and target species and restriction areas where emergency handling is taking place. The intensity of the impacts is assessed to be medium, with regional extent and short-term duration. The overall significance of impacts on fishery from major oil spill is assessed to be minor negative.

Physical effects to target species for fishery may have other consequences for fishery. As impacts on fish and marine invertebrates from an oil spill are expected to be major negative, it is assessed that impacts on fisheries will also occur. Further impacts on fisheries may arise due to market perceptions of poor product quality (no buyers or reduced prices, etc.). A major oil spill in the North Sea may significantly decrease buyers interest in fish and shellfish from the area. This can lead to loss of business and affect local economy. Perceptions are difficult to predict, since the actual (physical) impacts of the spill might have little to do with these perceptions. As a precautionary approach, the intensity of the impacts is assessed to be large, with transboundary extent and long-term duration. The overall significance of impacts on fishery from major oil spill from is assessed to be major.

7.3.4.3 Overall assessment

The potential impacts are summarised in Table 7-15.

**Table 7-15 Potential impacts on fisheries related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Gas release</b>	Medium	Transboundary	Short-term	Minor negative	Medium
<b>Major oil spill</b> Impacts on target species	Medium	Transboundary	Medium-term	Major negative	Low
<b>Major oil spill</b> Perception/reputation	Large	Transboundary	Long-term	Major negative	Low

7.3.5 Tourism

Potential impact mechanisms for tourism are related to major oil spill.

7.3.5.1 Major oil spill

The oil spill modelling shows, that Danish, Norwegian, German, Dutch and UK shorelines could be affected by oil. The Danish shoreline and more particularly the beaches of the west coast of Jutland and the Wadden Sea national parks are most likely to be affected. This may have a negative reputational impact for the affected region and may deter tourists from returning for years. This may lead to loss of business and impact local economy. An oil spill can thus result in long term effects on tourism.

The intensity of the impacts is assessed to be large, with transboundary extent and long-term duration. The overall significance of impacts on tourism from a major oil spill at the DAN project is assessed to be major negative.

7.3.5.2 Overall assessment

The potential impacts are summarised in Table 7-16.

**Table 7-16 Potential impacts on tourism related to accidental events at the DAN project.**

Potential impact mechanism	Intensity of impact	Extent of impact	Duration of impact	Overall significance of impact	Level of confidence
<b>Major oil spill</b>	Large	Transboundary	Long-term	Major negative	Medium

### 7.4 Summary

The potential impacts on environmental and social receptors from accidental events at the DAN project are summarised in Table 6-26. The impact with the largest overall significance is provided for each receptor.

**Table 7-17 Summary of potential impacts on environmental and social receptors for accidental events at the DAN project. The impact with the largest overall significance is provided for each receptor.**

Receptor	Worst case potential impact
Climate and air quality	Moderate negative
Water quality	Moderate negative
Sediment type and quality	Moderate negative
Plankton	Minor negative
Benthic communities	Major negative
Fish	Major negative
Marine mammals	Major negative
Seabirds	Major negative
Cultural heritage	Moderate negative
Protected areas	Major negative
Marine spatial use	Moderate negative
Fishery	Major negative
Tourism	Major negative

## 8. MITIGATING MEASURES

Maersk Oil have identified several mitigating measures for planned activities and accidental events with likely significant impacts on environmental or social receptors. The mitigating measures are in place to eliminate or reduce the environmental and social risks as low as reasonably practicable (ALARP). In addition to the mitigating measures, several monitoring campaigns are conducted around Maersk Oil platforms (section 9.5).

### 8.1 Mitigating for planned activities

#### 8.1.1 Measures to reduce emissions

Maersk Oil plans to implement a structured energy efficiency process and conduct a comprehensive review to identify ways to improve energy efficiency offshore. The production has become more energy efficient over the years, and in 2013 the energy management at Maersk Oil was ISO-14001 certified. Annual audits of performance and environmental action plans are part of this. The system is to be certified every third year.

#### 8.1.2 Underwater noise mitigating measures

The risks of underwater noise in geophysical acquisition and construction projects on marine mammals are mitigated by:

- Planning and efficient execution of the geophysical data acquisition and construction projects to minimise the duration of the operations.
- Monitoring the presence of marine mammals before the onset of noise creating activities, and throughout the geophysical data acquisition or construction.
- In areas where impacts on marine mammals are anticipated, best available technology will be assessed.
- An exclusion zone is implemented and operations will be delayed when the presence of marine mammal is detected before start-up of the operations.
- Soft-start procedures, also called ramp-up, should be used in areas of known marine mammal activity. This involves a gradual increase in sound signal level to full operational levels.

Maersk Oil evaluates the applicability or feasibility of relevant technological development in its activities (e.g. narrow frequency band air gun, bucket foundation) that could reduce the risk of underwater noise on the marine environment.

#### 8.1.3 Discharge mitigating measures

Maersk Oil uses chemicals in its operations, and is constantly examining the use and discharge of chemicals. Before any chemicals can be permitted for use and discharge offshore, an application must be submitted to the Danish authorities. Part of the application is an environmental classification of each chemical carried out in accordance with the OSPAR Recommendation 2010/4 on a harmonised pre-screening scheme for offshore chemicals. The classification applies a colour coding system used by Maersk Oil based on the criteria outlined in OSPAR, 2010 /44/:

- **Black:** Black chemicals contain one or more components registered in OSPAR's 'List of Chemicals for Priority Action'. The use of black chemicals is prohibited except in special circumstances. Maersk Oil has not used them since 2005 but has dispensation in 2015 to use black pipe dope in part of the casing in the drilling of a high-pressure, high temperature exploration well.
- **Red:** These are environmentally hazardous and contain one or more components that, for example, accumulate in living organisms or degrade slowly. OSPAR recommendation is that the discharge of these chemicals must end by 1 January 2017. Since 2008, Maersk Oil has been phasing out red chemicals, using them only if safety, technological and environmental arguments require use. Discharges have decreased sharply since 2010.

- **Green:** These contain environmentally acceptable components recorded on OSPAR's PLONOR list that 'pose little or no risk' to the environment.
- **Yellow:** These are chemicals not covered by the other classifications and can normally be discharged.

The risks of impact on the environment of operational discharges associated with production are mitigated through management of produced water through Risk Based Approach (RBA) in accordance with the OSPAR Guidelines and Recommendation /4/.

The RBA is used to review management options, evaluate measures and develop and implement site-specific actions to reduce environmental risks of production chemicals discharges which are not adequately controlled. Risk reduction measures may comprise some of the following:

- Technical measures, such as abatement at the source by redesign of the applied processes (water shut off in the well);
- Substitution of chemicals;
- Application of closed systems (e.g. injection of produced water);
- End-of-pipe techniques such as separation or clarification techniques to treat produced water prior to discharge, and;
- Organisational measures such as management systems in place (training, instructions, procedures and reporting).

An important tool within the RBA is the use of hydrodynamic models to predict the dispersion of the produced water outflow with a substance based approach /155/. This allows to identify the most important contributors to the risk and evaluate chemical substitution options while ensuring the application of BAT/BEP.

## 8.2 Mitigating of accidental events

Maersk Oil acts according to the zero tolerance for spills policy. This prescribes that all accidental discharges of oil and chemicals, regardless of volume, must be reported. Measures are introduced to reduce the volume and number of spillage through e.g. inspections and training. Maersk Oil follows industrial best practices for prevention of major accidents based on identification of major hazards assessed through risk assessment /136/.

Maersk Oil strives to reduce the risk of major accidents to as low as reasonably practicable (ALARP) through the identification of major hazards in risk analyses and the development of barriers (e.g. procedure, training, and design). For example, facilities are protected against collision by installing boat fenders to jackets. Processing facilities, wells and pipelines are protected against large release by safety valves. A safety zone around pipelines and platforms is implemented to prevent collisions from bottom trawling equipment or anchoring. Procedures are in place to restricted supply vessel traffic and hose handling in case of rough weather (see also Appendix 1).

The risk assessment and reduction measures are regularly updated in case of significant new knowledge or technology development.

Emergency response and contingency planning are also developed to limit the consequence in case of a major accident related to its projects. Maersk Oil's oil spill contingency plan is summarised in section 9.

## 9. ENVIRONMENTAL STANDARDS AND PROCEDURES IN MAERSK OIL

### 9.1 Environmental management system

Maersk Oil operates with a ISO 14001 certified environmental management system /121/. The objectives of the environmental management system is to minimise the impact on the environment by continually improving the environmental performance.

The objective shall be achieved by:

- Maintaining a complete and effective environmental management system
- Providing timely and effective innovative actions to reduce environmental impact
- Promoting the awareness of environmental matters at all organisational levels
- Minimising environmental impact through principles of best available technology (BAT) and best environmental practice (BEP).

### 9.2 Environmental and social impact in project maturation

An Environmental and Social Impact Assessment standard /159/that lays out the process for managing risk of environmental and social impacts of new large projects has recently been implemented in Maersk Oil. The standard provides a framework embedded within the Maersk Oil project maturation process which will be used from start and throughout the different development phases of future development projects.


### 9.3 Demonstration of BAT/BEP

The OSPAR Convention of 1992 requires contracting parties to apply best available techniques (BAT) and best environmental practice (BEP) including, where appropriate, clean technology, in their efforts to prevent and eliminate marine pollution.

*As defined the OSPAR convention BAT means the latest stage of development (state of the art) of processes, of facilities or of methods of operation which indicate the practical suitability of a particular measure for limiting discharges, emissions and waste. BEP is defined as the application of the most appropriate combination of environmental control measures and strategies.*

It follows that BAT and BEP for a particular source will change with time in the light of technological advances, economic and social factors, as well as changes in scientific knowledge and understanding.

BAT has also been implemented in the EU IPPC directive 96/61/EC, and the IE directive (2010/75/EU). What constitutes BAT is identical in the two directives, but Articles 13 to 16 of the IE directive require that BAT reference documents are prepared as a reference for setting permit conditions. The BAT principle is illustrated in Figure 9-1.



**Figure 9-1 Illustration of best available technique**

It is a Maersk Oil objective to implement the principles of BAT and BEP in an effort to minimize the potential environmental impacts of activities in the North Sea. This entails that environmental concerns are addressed and encompassed in the planning phase. The BAT/BEP principle has been used in the design and operation of the installations and process equipment of Maersk Oil as well as for the selection of materials and substances.

Examples of how Maersk Oil applies BAT and BEP include measures to:

- Improving energy efficiency
- Monitoring and minimising emissions
- Optimising the use and discharge of chemicals
- Supporting the development of chemicals with less environmental impact
- Use of efficient equipment during well test
- Continuous review and assessment of projects and applied equipment

For example, Maersk Oil and gas use several technologies such as hydrocyclones, induced gas flotation units, compact flotation units for treatment of produced water, which are included in the OSPAR background document concerning techniques for the management of produced water from offshore installations, an overview from 2002 of BAT for handling produced water.

#### **9.4 Oil spill contingency plan**

Maersk Oil's emergency preparedness in connection with serious incidents offshore on and around Maersk Oil's installations and in Danish concession areas held by A.P. Møller-Mærsk is centred around and coordinated by permanently established emergency committees.

Maersk Oil has developed an oil spill contingency plan /121/, which describes how to combat possible oil spills. Oil spill scenarios up to and including the worst credible case discharge scenario for Maersk Oil facilities and wells have been considered to ensure that an appropriate tiered capability is established.

- Tier 1: e.g. small operational spills
  - Mobilise oil spill monitoring/surveillance vessel.
  - Oil spill drift modelling.
  - Use in-field vessel with boom/skimmer equipment mobilised within 8 hours.

- Tier 2: medium spill volume
  - Tier 1 measures.
  - Use of additional resources (boom, skimmer and transfer pump/hoses) mobilised from Esbjerg or from the Danish National stockpile within 20 hour.
  - Waste removal is done by dedicated tanker.
- Tier 3: e.g. blow out
  - Tier 2 measures.
  - Mobilise additional vessel with 1200m boom, skimmer and transfer pump/hoses within 30 hour. Mobilise trained personnel and additional equipment from Oil Spill Response Ltd (OSRL).
  - Waste removal is done by dedicated tanker.
  - Mobilise relief well contractor.
  - Consult NGOs regarding wildlife response.

Maersk Oil has access to oil spill equipment offshore and in Esbjerg that can be mobilised to an oil spill location immediately. If necessary, additional equipment will be mobilised from the Danish stock pile and OSRL. Maersk Oil is a participant member of OSRL and has access to their world-wide pool of personnel and equipment. OSRL's main equipment stockpile in Europe is based in Southampton in the UK but additional equipment is also available in Stavanger.

The use of dispersant chemicals to increase oil dispersion, dilution and natural breakdown will be evaluated when relevant. The use of dispersant chemicals is regulated and dispersant may only be used after approval by DEPA.

Regular emergency exercises (oil spills) are carried out as a minimum every three years to train and motivate personnel, test the equipment and to ensure plans as described are effective. Relevant authorities participate in the exercise.

## 9.5 Ongoing monitoring

Maersk Oil has flowmeters measuring the volume of discharged produced water, and water samples are regularly obtained for analysis of oil and chemical content. The nature, type and quantities chemical used and chemicals and oil discharged to sea are reported to the Environmental Agency (DEPA).

Monitoring of sediment quality and benthic fauna is undertaken at regular intervals around Maersk Oil platforms /6/.

- The physical and chemical analyses included grain size analysis, dry matter (DM), loss on ignition (LOI), total organic carbon (TOC), metals (barium (Ba), cadmium (Cd), chromium (Cr), copper (Cu), lead (Pb), zinc (Zn), mercury (Hg) and aluminium (Al)), Total hydrocarbons (THC), Polycyclic aromatic hydrocarbons (PAH) and oil specific group of alkylated aromatic hydrocarbons (NPD).
- Samples obtained for identification and quantification of the benthic fauna

In addition, Maersk Oil monitors underwater noise and marine mammals through passive acoustic monitoring and an offshore sighting program in which offshore staff reports sightings of marine mammals near platforms.



**Figure 9-2 Acoustic monitoring of marine mammals (Photo: Aarhus University, DCE).**

## 10. NATURA 2000 SCREENING

### 10.1 Introduction

The Natura 2000 network comprises:

- Habitats Directive Sites (Sites of Community Importance and Special Areas of Conservation) designated by Member States for the conservation of habitat types and animal and plant species listed in the Habitats Directive
- Bird Directive Sites (Special Protection Areas) for the conservation of bird species listed in the Birds Directive as well as migratory birds

This section constitutes the Natura 2000 screening in accordance with the EC habits Directive and Order 408/2007, § 7.

### 10.2 Designated species and habitats

The designated Natura 2000 sites are shown in Figure 10-1.




Figure 10-1 Natura 2000 sites in the North Sea.

Natura 2000 sites in the central North Sea are detailed in Table 10-1.

**Table 10-1 Natura 2000 sites in the central North Sea.**

Natura 2000 Site code	Name	Designated marine species and habitattypes
UK0030352	Dogger Bank	<ul style="list-style-type: none"> <li>• 1110 Sandbanks which are slightly covered by sea water all the time</li> <li>• 1351 <i>Phocoena phocoena</i></li> <li>• 1364 <i>Halichoerus grypus</i></li> <li>• 1365 <i>Phoca vitulina</i></li> </ul>
NL2008002	Klaverbank	<ul style="list-style-type: none"> <li>• 1170 Reefs</li> <li>• 1351 <i>Phocoena phocoena</i></li> <li>• 1364 <i>Halichoerus grypus</i></li> <li>• 1365 <i>Phoca vitulina</i></li> </ul>
NL2008001	Doggersbank	<ul style="list-style-type: none"> <li>• 1110 Sandbanks which are slightly covered by sea water all the time</li> <li>• 1351 <i>Phocoena phocoena</i></li> <li>• 1364 <i>Halichoerus grypus</i></li> <li>• 1365 <i>Phoca vitulina</i></li> </ul>
DE1003301	Doggerbank	<ul style="list-style-type: none"> <li>• 1110 Sandbanks which are slightly covered by seawater all the time</li> <li>• 1351 <i>Phocoena phocoena</i></li> <li>• 1365 <i>Phoca vitulina</i></li> <li>• <i>Fulmarus glacialis, Larus fuscus, Morus bassanus, Rissa tridactyla, Uria aalge</i></li> </ul>
DE1209301	Sylter Außenriff	<ul style="list-style-type: none"> <li>• 1110 Sandbanks which are slightly covered by sea water all the time</li> <li>• 1170 Reefs</li> <li>• 1351 <i>Phocoena phocoena</i></li> <li>• 1364 <i>Halichoerus grypus</i></li> <li>• 1365 <i>Phoca vitulina</i></li> <li>• 1103 <i>Alosa fallax</i></li> <li>• <i>Gavia arctica, Gavia stellata, Lampetra fluviatilis, Larus canus, Larus fuscus, Larus marinus, Larus minutus, Morus bassanus, Rissa tridactyla, Sterna hirundo, Sterna paradisaea, Sterna sandvicensis, Uria aalge</i></li> </ul>
DK00VA347	Sydlig Nordsø	<ul style="list-style-type: none"> <li>• 1110 Sandbanks which are slightly covered by sea water all the time</li> <li>• 1351 <i>Phocoena phocoena</i></li> <li>• 1364 <i>Halichoerus grypus</i></li> <li>• 1365 <i>Phoca vitulina</i></li> <li>• <i>Gavia stellata, Gavia arctica, Larus minutus, Sula bassana, Somateria mollissima, Melanitta nigra, Stercorarius skua, Uria alge, Alca torda, Alle alle</i></li> </ul>
DK00VA257	Jyske Rev	<ul style="list-style-type: none"> <li>• 1170 Reefs</li> <li>• 1351 <i>Phocoena phocoena</i></li> </ul>

### 10.3 Screening

The screening is carried out to identify all those elements of the DAN project, alone or in combination with other projects or plans, that may have significant impacts on the Natura 2000 site.

No activities associated with the DAN project are planned to occur within a designated Natura 2000 sites. The closest Natura 2000 area from the DAN project is the German Dogger Bank located 25 km from Kraka.

Potential impact expected for activities for the DAN project are generally local (section 6). It is assessed that the only potential sources of impacts on the Natura 2000 sites include underwater noise and discharges.

#### 10.3.1 Underwater noise

A number of activities that are expected to be conducted for the DAN project may generate underwater noise. Activities include seismic data acquisition, construction, production and drilling operations, as well as transport. There is no geographical overlap between the DAN project area and Natura 2000 sites. However, underwater noise from airguns used in seismic acquisition, pile driving during the construction of the new platforms for the DAN project and ramming of conductors for the new wells may spread into nearby Natura 2000 sites.

Based on the distance between the DAN project and Natura 2000 sites (25 km), it is assessed that planned activities will not have significant environmental effects on the conservation objectives of the Natura 2000 sites.

#### 10.3.2 Discharges

The main discharges are related to production and drilling activities, though other minor discharges may also occur (e.g. from vessels).

- Discharges of water-based drilling mud and drill cuttings are expected to occur from a drilling rig at Dan F (3 wells and 14 re-drilling wells) or at the new wellhead platforms (40 wells). The area that will be impacted by the discharges will depend on the currents, and likely follow the prevailing northward currents in the Dan project area i.e. away from the closest Natura 2000 site. The results from hydrodynamic modelling for the discharge associated with a typical well indicate that there is a risk of affecting the marine environment with a PEC/PNEC ratio of 1 up to 7 km from the discharge point (section 6). Similar modelling indicates that there is a risk of affecting the sediment quality up to a few hundred meters from the drilling rig (section 6.2.4). The distance from the point of discharge (Dan F, Kraka) to the nearest Natura 2000 site is at least 25 km. It is therefore assessed that drilling activities will not have a significant environmental impact on the conservation objectives of the habitat types or species in the Natura 2000 sites.
- Discharges from production at Dan F can be expected to continue until end of the production for the DAN project (expected 2042). The area that will be impacted by the discharges will depend on the currents, and likely follow the prevailing northward currents in the Dan area i.e. away from the closest Natura 2000 site. The results from hydrodynamic modelling for the discharge associated with production for the DAN project indicate that there is a risk of affecting the water quality from the current discharges of produced water up to 31 km from Dan F /42/ (section 6). The area where impacts to the marine environment cannot be ruled out (PEC/PNEC ratio of 1) will depend on the currents, and will likely follow the prevailing northward currents, in the Dan area i.e. away from the closest Natura 2000 site. However, production discharges at Dan F may have an impact on the water quality in the Natura 2000 site Dogger Bank (26 km from Dan F), if southwestern currents dominate. It is assessed that any potential impact to the water quality in the Natura 2000 site is local and short-term, and without significant environmental effects on habitat types or species in the Natura 2000 sites.

### 10.4 Conclusion

It is assessed that planned activities for the DAN project will not have a significant environmental impacts on the conservation objectives of the habitat types or species in the Natura 2000 sites.

## 11. TRANSBOUNDARY IMPACTS

### 11.1 Introduction

The DAN project refers to the platform Dan F, Dan B, Dan E, Kraka and Regnar. An environmental and social impact assessment (EISA-16) is undertaken for the remaining lifetime of the ongoing projects, and the entire life time from exploration to decommissioning for planned projects. The ESIA-16 shall replace the EIA conducted in 2010 "Environmental impact assessment from additional oil and gas activities in the North Sea, July 2011" which is valid for the period 1st January 2010 to 31st December 2015.

Notifications for the DAN project were forwarded to the relevant German authorities in accordance with article 3.1 of the Espoo convention. The German authorities have expressed the wish to participate in the hearing for the ESIS DAN.

In this section, a summary of the DAN project and its likely significant transboundary impacts is provided. The section is focused on providing sufficient information to facilitate the identification of possible transboundary impacts. The rationale and support for the attributed level of significance and spatial extent can be found in detail in the relevant sections of the ESIS (section 6 and 7).

### 11.2 ESPOO convention

The ESPOO convention states that the concerned parties likely to be affected by transboundary adverse significant impacts are to be informed of and provided with possibilities for making comments or objections on the proposed activity.

The DAN project can be found as item 15 (offshore hydrocarbon production) on the list of activities in appendix I to the convention, that are likely to cause a significant adverse transboundary impact.

### 11.3 The DAN project

#### 11.3.1 Existing production and processing facilities

The DAN project, including the three platforms Dan F, Dan B and Dan E as well as the satellite platform Kraka and the subsea wellhead Regnar. Production was initiated at Dan B in 1972, then later at Dan Dan E (1977), Dan F (1987), Kraka (1991) and Regnar (1993). The total production peaked in 2000 and has been on a natural decline since. Maersk Oil has the license to explore for and produce oil and gas was extended until 8 July 2042.

Dan F is primarily an oil producing and oil processing platform that receives, processes and sends the Dan field production to Gorm (oil) and Tyra (gas). Regnar is an unmanned subsea wellhead, while Kraka is an unmanned satellite wellhead platform.

The processing facilities include hydrocarbon processing equipment (oil stabilisation, gas processing and processing of production water), auxiliary safety systems such as an emergency shutdown system, emergency blow-down system, fire and gas detection system, firewater system, etc.




Figure 11-1 Maersk Oil North Sea projects TYRA, HARALD, DAN, GORM and HALFDAN.

11.3.2 Planned development activities

The following main activities are planned to continue and optimise the production for the DAN project and potentially access new hydrocarbon resources:

- Seismic investigations to provide information to interpret the geological structure of the subsurface and to identify the location and volume of remaining and potential new hydrocarbon reserves. Seismic data is also acquired as part of drilling hazard site surveys to map and identify potential hazards to the installation of drilling rigs and to the drilling operation. Seismic data are also acquired as part of seabed and shallow geophysical surveys to map seabed and shallow soil conditions for the design and installation of pipelines, platforms and other structures.
- Construction of up to five new platforms (four wellhead platforms and an accommodation platform), along with connecting pipelines
- Drilling of up to 57 wells (43 new wells, and 14 redrilling from existing wells) can be expected as part of the DAN project. Drilling is performed from a drilling rig, which is placed on the seabed. Different types of drilling mud will be used based on the well and reservoir properties. Water-based mud and cuttings will be discharged to the sea, whereas oil-based mud and cuttings will be brought onshore to be dried and incinerated.
- Well stimulation will be performed to facilitate hydrocarbon extraction (for a production well) or water injection (for an injection well).
- Decommissioning will be done in accordance with technical capabilities, industry experience, relevant international conventions and under the legal frameworks at the time of decommissioning.

11.3.3 Accidental events

As part of the production, accidental spills of oil, gas or chemical may occur. There is a risk of accidents that could lead to major significant environmental and social impacts, such as vessels collisions, large pipeline rupture or a well blow out. The risk of a well blowout is very unlikely.

#### 11.3.4 Alternatives

##### Project alternative

The 0 alternative (zero alternative) is a projection of the anticipated future development without project realization, and describes the potential result if nothing is done. For the DAN project, this would mean that the production would cease. If no production is undertaken by Maersk Oil for the DAN project area in the North Sea, there will be no contribution from the DAN project to the Danish economy or security of hydrocarbon supply and employment.

##### Technical alternatives

Best environmental practice for the different type of activities planned for the DAN project (seismic, pipelines and structures, production, drilling, well stimulation, transport and decommissioning) is continuously monitored and applied when feasible.

##### Alternative location

The DAN project is a continuation of production and activities at existing facilities. As such, there is no alternative location for the project.

#### 11.4 Identified impacts – planned activities

Potential impacts to environmental and social receptors during planned activities at the DAN project have been assessed in section 6. A summary of the potential worst case impacts is presented in Table 11-1.

**Table 11-1 Summary of potential impacts on environmental and social receptors from planned activities at the DAN project. The impact with the largest overall significance is provided for each receptor (without mitigating measures).**

Receptor	Worst case potential impact	
	Extent	Overall significance of impact
<b>Hydrography</b>	Local	Minor negative
<b>Climate and air quality</b>	Transboundary	Moderate negative
<b>Water quality</b>	Local	Minor negative
<b>Sediment type and quality</b>	Local	Minor negative
<b>Plankton</b>	Local	Minor negative
<b>Benthic communities</b>	Local	Minor negative
<b>Fish</b>	Local	Minor negative
<b>Marine mammals</b>	Local or regional	Moderate negative
<b>Seabirds</b>	Local	Minor negative
<b>Cultural heritage</b>	None	None
<b>Protected areas (UNESCO, nature reserve)</b>	None	None
<b>Natura 2000</b>	No significant environmental effects	
<b>Marine spatial use</b>	Local	Negligible negative
<b>Fishery</b>	Local	Negligible negative
<b>Tourism</b>	None	None
<b>Employment and tax revenue</b>	Local or national	Positive
<b>Oil and gas dependency</b>	Local or national	Positive

Transboundary adverse impacts have been identified for climate and air quality, where the emissions from the DAN project may have a minor contribution to climate change and air pollution. Maersk Oil has implemented a structured energy efficiency process and conduct a comprehensive review to identify ways to improve energy efficiency offshore. The production has become more energy efficient over the years, and in 2013 the environmental management system at Maersk Oil was ISO-14001 certified.

No other significant adverse transboundary impacts have been identified for the planned activities at the DAN project.

A Natura 2000 screening is presented for the planned activities. It is assessed that the planned activities will have no significant environmental effects on the conservation objectives of the habitat types or species in the national and international Natura 2000 sites (section 10).

### 11.5 Identified impacts – accidental events

Potential impacts to environmental and social receptors during accidental events from the DAN project have been assessed in section 7. A summary of the worst case potential impacts (without mitigating measures) is presented in Table 11-2.

**Table 11-2 Summary of potential impacts on environmental and social receptors for accidental events at the DAN project. The impact with the largest overall significance is provided for each receptor (without mitigating measures).**

Receptor	Worst case potential impact	
	Extent	Overall significance of impact
<b>Climate and air quality</b>	Transboundary	Moderate negative
<b>Water quality</b>	Transboundary	Moderate negative
<b>Sediment type and quality</b>	Transboundary	Moderate negative
<b>Plankton</b>	Transboundary	Minor negative
<b>Benthic communities</b>	Transboundary	Major negative
<b>Fish</b>	Transboundary	Major negative
<b>Marine mammals</b>	Transboundary	Major negative
<b>Seabirds</b>	Transboundary	Major negative
<b>Cultural heritage</b>	National	Moderate negative
<b>Protected areas (UNESCO, nature reserve)</b>	Transboundary	Major negative
<b>Marine spatial use</b>	Transboundary	Moderate negative
<b>Fishery</b>	Transboundary	Major negative
<b>Tourism</b>	Transboundary	Major negative

If a major oil spill occurs, there is a risk of major negative transboundary impacts. The risk of a major oil spill is very unlikely, but could potentially have significant, adverse transboundary impacts. Oil released could cross maritime boundaries with Norway, Germany, the Netherlands and the UK. The oil spill modelling identified the north and west of Denmark and south Norway as most vulnerable to oil beaching, although Germany, UK and the Netherlands could also be affected.

Maersk Oil acts according to the zero tolerance for spills policy. Measures are introduced to reduce the volume and number of spillage, and Maersk Oil follows industrial best practices for prevention of accidents based on identification of major hazards assessed through risk assessment. Emergency response and contingency planning are also developed to limit the consequences of a major accident related to its projects

## 12. LACK OF INFORMATION AND UNCERTAINTIES

Uncertainty may be viewed as an inescapable part of assessment of impacts of plans, programmes or projects.

### 12.1 Project description

The project description has been based on input from Maersk Oil. The project description is based on a scenario with maximum activity, emissions and discharges.

For some activities, the location and/or timing has not been decided. This will be done as part of the preparation of the detailed planning of the activities. The ESIS is undertaken using a worst case approach, and therefore minor alterations to location and/or timing is assessed to be of minor influence to the assessments.

The understanding of the employment and tax revenue for the project has not been described in detail. The assessment is therefore based on the overall DUC contribution.

### 12.2 Environmental and social baseline

The central North Sea is relatively well known, and the environmental and social baseline is generally considered sufficient for the ESIS.

However, a few receptors are less well understood:

- The distribution and biology of non-commercial fish species is scarce, and knowledge of spawning areas is limited.
- The variability of distribution of marine mammals within and between years is not well known, and the breeding and moulting periods and locations are not certain.
- Fishery is mapped based on the North Sea Atlas which applies ICES data. However, the variability between years is not detailed for this ESIS.

### 12.3 Impact assessment

Predictions can be made using varying means, ranging from qualitative assessment and expert judgement to quantitative techniques like modelling. Use of these quantitative techniques allows a reasonable degree of accuracy in predicting changes to the existing environmental and social conditions. However, not all of the assessed impacts are easy to measure or quantify, and expert assumptions are needed.

Uncertainty has been addressed in this ESIS by presenting a level of confidence for each of the assessments in section 6 and 7. The level of confidence includes interactions between impact mechanisms and receptors, available baseline data as well as modelling (section 4).

Overall, impacts are assessed based on today's technological capabilities. Maersk Oil expects that technological development will lead to a reduction in emissions and discharges, which will reduce impact.

#### 12.3.1 Planned activities

The potential environmental impacts have been assessed for each receptor (e.g. plankton, employment). The impact assessment is based on empirical studies, scientific literature, modelling results as well as previous EIAs.

Previous modelling results have been applied in this ESIS, with no site-specific modelling. Similar activities have previously been assessed for the same area, and modelling has been undertaken for e.g. dispersion of drill mud and cutting, dilution of produced water as well as propagation of underwater noise. In addition, Maersk Oil prepares EIF and the PEC/PNEC

calculations for each of the five projects, within the "CHARM" means the Chemical Hazard Assessment and Risk Management model developed by authorities and offshore industry. The calculations have some weaknesses (as reviewed in /1/), but is considered valid for the impact assessment.

The project which is assessed is at or near existing platforms, where monitoring of chemical and biological conditions have been undertaken for many years. These surveys contribute to a solid baseline, as well as an understanding of the environmental impacts.

Impacts of underwater noise is not well understood, and there is ongoing debate regarding thresholds for potential impact.

#### 12.3.2 Accidental events

Oil spill modelling has been undertaken for a number of spill scenarios. However, the spill rates for blowouts are not directly comparable, but considered applicable as a worst case scenario.

#### 12.3.3 Cumulative impacts

The North Sea is one of the most heavily trafficked in the world, and there are intensive fisheries. The Greater North Sea is surrounded by densely populated and highly industrialised countries, and regional and global changes to oceanic, atmospheric, and climate regulation processes pose additional threats. A number of responses and measures have been implemented to reduce pressures on the environment and resulting impacts, but despite this, the cumulative environmental effects on the area are causing concern.

There is no general method for combining impacts across different geographical scales and as a result of different pressures. It is therefore difficult to assess the severity of the cumulative environmental effects on the ecosystem. Uncertainty and lack of knowledge about the population status of species, the range and ecological status of habitat types, and the impacts of environmental pressures also add to the uncertainty of assessments of environmental impacts.

Assessment of the impact of oil and gas activities in isolation may thus understate overall impacts by excluding potential impacts of past, present, or future impacts of other human activities.

## 13. REFERENCES

- /1/ Maersk Oil. 2011. Vurdering af virkningen på miljøet fra yderligere olie og gas aktiviteter i Nordsøen.
- /2/ Maersk Oil. 2014. Vurdering af virkninger på miljøet (VVM) for etablering og drift af Adda og Tyra N.
- /3/ DHI/Ramboll. 2014. Maersk Oil Atlas of Environment and Industrial Activities in the North Sea.
- /4/ IRIS. 2014. Risk based approach to produces water management. EIF calculations Dan 2013.
- /5/ OSRL. 2014. Oil spill modelling report: Siah NE-1X. Doc CONS0874, rev 02.
- /6/ DHI, 2009. Chemical and biological monitoring of the seabed around the Dan F platform in May 2009.
- /7/ Statens Forurensning Tilsyn. 2007. Veileder for klassifisering av miljøkvalitet i fjorder og kystfarvann. Revidering av klassifisering av metaller og organiske miljøgifter i vann og sedimenter . TA-2229/2007.
- /8/ Vejledning fra By – og Landskabsstyrelsen. 2008. Dumpning af optaget havbundsmateriale – klapping. Vejl. nr. 9702 af 20/10 2008.
- /9/ OSPAR Commission. 2012. CEMP 2011 assessment report.
- /10/ OSPAR. 2000. Quality Status Report 2000, Region II - Greater North Sea. OSPAR Commission, London, 136 pages.
- /11/ UK Offshore Energy - Department of Energy and Climate Change. 2009. Strategic Environmental Assessment Future Leasing for Offshore Wind Farms and Licensing for Offshore Oil & Gas and Gas Storage. Appendix 3a Biodiversity, habitats, flora and fauna.
- /12/ SAHFOS. 2001. Technical report produced for Strategic Environmental Assessment – SEA2. An overview of plankton ecology in the North Sea.
- /13/ Callaway, R., Alsvåg, J., de Boois, I., Cotter, J., Ford, A., Hinz, H., Jennings, S., Kröncke, I., Lancaster, J., Piet, G., and Prince, P. 2002. Diversity and community structure of epibenthic invertebrates and fish in the North Sea. ICES Journal of Marine Science, 59: 1199–1214.
- /14/ Künitzer, A., Basford, D., Craeymeersch, J. A., Dewarumez, J. M., Dorjes, J., Duineveld, G. C. A., Eleftheriou, A., Heip, C., Herman, P., Kingston, P., Niermann, U., Rachor, E., Rumohr, H., and de Wilde, P. A. J. 1992. The benthic infauna of the North Sea: species distribution and assemblages. ICES Journal of Marine Science, 49: 127–143.
- /15/ ICES Advice. 2008. Book 6 North Sea.
- /16/ VKI. 1999. Analysis of Impact on Sediment and Bottom Fauna in Relation to Offshore Activities in the Danish Sector of the North Sea 1989-1998. North Sea Operators Committee – Denmark.
- /17/ DHI - Water Environment Health. 2008. Analyses and assessment of biological and chemical monitoring data from offshore platforms in the Danish sector of the North Sea in 1989-2006. Report to Danish Operators.
- /18/ Olsgard, F. and J.S. Gray. 1995. A comprehensive analysis of the effects of offshore oil and gas exploration and production on benthic communities of the Norwegian continental shelf. Marine Ecology Progress Series Vol.122: 277-306.
- /19/ Mærsk Olie og Gas AS. 2003. Registrering af fiskesamfund og fiskeæg omkring Halfdan-Feltet. Hovedrapport. Udarbejdet af Carl Bro, Bio/consult og Simråd.
- /20/ Mærsk Oil, Miljøstatusrapport 2013, Den danske Nordsø
- /21/ DCE Emission Inventories, <http://envs.au.dk/en/knowledge/air/emission-inventories/emissioninventory/>.
- /22/ Worsøe, L.A, Horsten, M.B and Hoffmann, E. 2002. Gyde og opvækstpladser for kommercielle fiskearter i Nordsøen, Skagerrak og Kattegat. DFU rapport nr. 118-02.
- /23/ ICES FishMap <http://www.ices.dk/marine-data/maps/Pages/ICES-FishMap.aspx>.
- /24/ Muus, B.J., Nielsen, J.G., Dahlstrøm, P. and Nyström, B.O. 1998. Havfisk og fiskeri. Gads Forlag.
- /25/ OSRL. 2014. Oil spill risk assessment: Siah NE-1X. Doc CONS0874.
- /26/ OSRL. 2014. Oil spill modelling report: Xana-1X. Doc CONS0896.
- /27/ OSRL. 2014. Oil spill risk assessment: Xana-1X. Doc CONS0848.
- /28/ Naturstyrelsen. 2012. Danmarks Havstrategi – Basisanalyse. [www.nst.dk](http://www.nst.dk).
- /29/ OSPAR. 2014. OSPAR/ICES Workshop on evaluation and update of BRCs and EACs. OSPAR report 214.
- /30/ Thompson, D. & Härkönen, T. (IUCN SSC Pinniped Specialist Group) 2008. Phoca vitulina. The IUCN Red List of Threatened Species. Version 2014.1. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 14 July 2014.

- /31/ Hammond, P.S., Bearzi, G., Bjørge, A., Forney, K., Karczmarski, L., Kasuya, T., Perrin, W.F., Scott, M.D., Wang, J.Y., Wells, R.S. & Wilson, B. 2008. *Phocoena phocoena*. The IUCN Red List of Threatened Species. Version 2014.1. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 14 July 2014.
- /32/ Thompson, D. & Härkönen, T. (IUCN SSC Pinniped Specialist Group) 2008. *Halichoerus grypus*. The IUCN Red List of Threatened Species. Version 2014.1. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 14 July 2014.
- /33/ Hammond, P.S., Bearzi, G., Bjørge, A., Forney, K.A., Karczmarski, L., Kasuya, T., Perrin, W.F., Scott, M.D., Wang, J.Y., Wells, R.S. & Wilson, B. 2012. *Lagenorhynchus albirostris*. The IUCN Red List of Threatened Species. Version 2014.1. <[www.iucnredlist.org](http://www.iucnredlist.org)>. Downloaded on 15 July 2014.
- /34/ Birdlife: Marine IBAs. <http://maps.birdlife.org/marineIBAs/default.html>. Accessed February 2nd 2015.
- /35/ Danish Energy Agency. 2014. Oil and gas production in Denmark 2013.
- /36/ FAO. 2013. Fishery Statistics and Information Unit in May 2013. <http://www.fao.org/fishery/facp/DNK/en>
- /37/ NaturErhvervstyrelsen. 2014. Dataudtræk fra landingsdatabasen. Accessed February 10<sup>th</sup> 2015.
- /38/ VisitDenmark. 2014. Turismens økonomiske betydning i Danmark 2012. ISBN: 978-87-93227-00-2
- /39/ Statistics Denmark. 2015. Beskæftigede efter branche og tid – RAS150. Accessed February 10<sup>th</sup> 2015.
- /40/ Skov, H. & Piper, W. 2009. Kortlægning af havfugle og havpattedyr i Nordsøen 2006-2008. Rapport over flytællingsresultater. Mærsk Olie & Gas AS.
- /41/ Southall, B., Bowles, A. E., Ellison, W. T., Finneran, J. J., Gentry, R. L., Greene, C. R. Jr., Kastak, D., Ketten, D. R., Miller, J. H., Richardson, W. J., Thomas, J. A., Tyack, P. L. 2007. Marine mammal noise exposure criteria: initial scientific recommendations. *Aquatic mammals* 33(4).
- /42/ IRIS. 2014. Risk based approach to produced water management: EIF calculations Dan 2013. Ref 2014/419
- /43/ DONG Energy. 2010. Vurdering af virkninger på miljøet for udbygning af Hejre-felter, licens 05/98. Ref CO86-COWI-S-DG-0002.
- /44/ OSPAR, 2010. OSPAR Recommendation 2010/4 on a Harmonised Pre-screening Scheme for Offshore Chemicals.
- /45/ Neff, J.M. 2010. Fate and effects of water based drilling muds and cuttings in cold water environments. Review prepared for Shell Exploration and Production Company Houston Texas. May 25, 2010.
- /46/ Bakke, T., Klungsøyr, J. & Sanni, S. 2013. Environmental impacts of produced water and drilling waste discharges from the Norwegian offshore petroleum industry. *Marine Environmental Research* 92 (2013) 154-169.
- /47/ Popper, A.N. & Hastings, M.C. 2009. The effects of human generated sound on fish. *Integrative Zoology*. 4: 43-52.
- /48/ Genesis. 2011. Review and assessment of underwater sound produced from oil and gas sound activities and potential reporting requirements under the marine strategy framework directive. Doc J71656-Final Report\_G2.
- /49/ OSPAR. 2014. OSPAR inventory of measures to mitigate the emission and environmental impact of underwater noise
- /50/ JNCC. 2010. JNCC guidelines for minimising the risk of injury and disturbance to marine mammals from seismic surveys
- /51/ Expert Working Group 2014. Memorandum prepared for Energinet.dk. »Marine mammals and underwater noise in relation to pile driving. « 21. 01 2015.
- /52/ Centre for Energy, Environment and Health. 2011. Assessment of Health-Cost Externalities of Air Pollution at the National Level using the EVA Model System. CEEH Scientific Report No 3
- /53/ QUARTZ+CO. 2012 The Danish oil and gas sector's development and social impact (1992-2022)
- /54/ Boertmann, D. and Mosbech, A. (eds.). 2011. Eastern Baffin Bay - A strategic environmental impact assessment of hydrocarbon activities. Scientific Report no 9, from Danish Centre for Environment and Energy. 270 pp.
- /55/ Christian, J.R., Mathieu, A., Thomson, D.H., White, D., and Buchanan, R.A. 2003. Effects of Seismic Energy on Snow Crab (*Chionoecetes opilio*). Report from LGL Ltd. and Oceans Ltd. for the National Energy Board, File No.: CAL-1-00364, 11 April 2003.

- /56/ Sherk, J.A., O'Connor, J.M., Neumann, D.A., Prince, R.D. and Wood, K.V. 1974. Effects of suspended sediment on feeding activity of the copepods *Eurytemora affinis* and *Acartia tonsa*. In: Effects of suspended and deposited sediments on estuarine organisms, phase 11. Final Report, chapter 8. Reference nos. 74-20. Natural Resources; Institute, University of Maryland, Prince Frederick, p.166-200.
- /57/ Paffenhöfer, G.A. 1972. The effects of suspended "Red mud" on mortality, body weight and growth of the marine planktonic copepod *Calanus helgolandicus*. *Water, Air and Soil Pollution* 1 (1972) 314-321
- /58/ Alldredge, A.L., Elias, M. and Gotschalk, C.C. 1986. Effects of drilling muds and mud additives on the primary production of natural assemblages of marine phytoplankton. *Marine Environmental Research* 19: 157 to 176.
- /59/ Neff, J.M. 2005. Composition, environmental fates and biological effects of water based drilling muds and cuttings discharged to the marine environment: A synthesis and annotated bibliography. Prepared for Petroleum Environmental Research Forum (PERF) and American Petroleum Institute.
- /60/ Haney, J.F. 1993. Environmental control of diel vertical migration behaviour, *Arch. Hydrobiol. Beih. Ergebn. Limnol* 39, pp. 1-17.
- /61/ Vella, G., Rushforth, I., Mason, E., Hough, A., England, R., Styles, P., Holt, T. & Thorne, P. 2001. Assessment of the effects of noise and vibration from offshore wind farms on marine wildlife. ETSU W/13/00566/REP – DTI/Pub URN 01/1341. 107 pages.
- /62/ Dietz, R.S. 1962. The sea's deep scattering layers. *Scientific American*.
- /63/ Gliwicz, Z.M. 1986. A lunar cycle in zooplankton. *Ecology* 67, pp. 883-97.
- /64/ U.S. Geological Survey. 2011. Final programmatic environmental impact statement for marine seismic research funded by the national science foundation or conducted by the U.S. Geological Survey.
- /65/ Vella, G., Rushforth, I., Mason, E., Hough, A., England, R., Styles, P., Holt, T. and Thorne, P. 2001. Assessment of the effects of noise and vibration from offshore wind farms on marine wildlife. ETSU W/13/00566/REP – DTI/Pub URN 01/1341.
- /66/ Christian, J.R., Mathieu, A., Thomson, D.H., White, D. and Buchanan, R.A. 2003. Effects of seismic energy on snow crab (*Chionoecetes opilio*). Report from LGL Ltd. and Oceans Ltd. for the National Energy Board, File No.: CAL-1-00364, 11 April 2003.
- /67/ de Groot, S. J. 1986. Marine sand and gravel extraction in the North Atlantic and its potential environmental impact, with emphasis on the North Sea, *Ocean Management*, Vol. 10, pp. 21- 36.
- /68/ Currie, D.R. and Parry, G.D. 1996. Effects of scallop dredging on a soft sediment community: a large scale experimental study. *Marine Ecology Progress Series*, Vol. 134, pp. 131- 150.
- /69/ Caddy, J.F. 1973. Underwater observations on tracks of dredges and trawls and some effects of dredging on a scallop ground. *Journal of the Fisheries Research Board of Canada*, Vol. 30, pp. 173-180.
- /70/ Chapman, C.J., Mason, J. and Drinkwater, J.A.M. 1977. Diving observations on the efficiency of dredges used in the Scottish fishery for the scallop, *Pecten maximus* (L.). *Scottish Fisheries Research*, Vol. 10.
- /71/ Ramsay, K., Kaiser, M.J. and Hughes, R.N. 1998. Responses of benthic scavengers to fishing disturbance by towed gears in different habitats. *Journal of Experimental Marine Biology and Ecology*, Vol. 224, pp. 73- 89.
- /72/ Boyd, S.E., Cooper, K.M., Limpenny, D.S., Kilbride, R., Rees, H.L., Dearnaley, M.P., Stevenson, J., Meadows, W.J. and Morris, C.D. 2004. Assessment of the rehabilitation of the seabed marine aggregate dredging.
- /73/ Turk, T.R. and Risk, M.J. 1981. Effect of sedimentation on infaunal invertebrate populations of Cobequid Bay, Bay of Fund. *Canadian Journal of Fisheries & Aquatic Sciences*, Vol. 38, pp. 642- 648.
- /74/ Lewis, L.J., Davenport, J. and Kelly, T.C. 2002. A study of the impact of a pipeline construction on estuarine benthic invertebrate communities, *Estuarine, Coastal and Shelf Science*, Vol. 55.
- /75/ Lisbjerg, D., Petersen, J.K. and Dahl, K. 2002. Biologiske effekter af råstofindvinding på epifauna. Faglig rapport fra DMU, nr. 391.
- /76/ Currie, D.R. and Parry, G.D. 1996. Effects of scallop dredging on a soft sediment community: a large scale experimental study, *Marine Ecology Progress Series*, Vol. 134, pp. 131- 150.
- /77/ Mitson, R.B. Underwater Noise of Research Vessels: Review and Recommendations. ICES Cooperative Research Report 1995;209:61.

- /78/ Fabi, G., Grati, F., Lucchetti, A. and Trovarelli, L. 2002. Evolution of the fish assemblages around a gas platform in the northern Adriatic Sea. ICES Journal of Marine Science. Vol. 59, Supplement 1, October 2002 pp. S309-S315.
- /79/ Løkkeborg, S., Humborstad, O.B., Jørgensen, T. and Soldal, A.V. 2002. Spatiotemporal variations in gillnet catch rates in the vicinity of North Sea oil platforms. ICES Journal of Marine Science Vol 59, Supplement 1 October 2002 pp S294- S299.
- /80/ Soldal, A.V., Svellingen, I., Jørgensen, T. and Løkkeborg, S. 2002. Rigs to reefs in the North Sea: hydroacoustic quantification of fish in the vicinity of a "semi-cold" platform. ICES Journal of Marine Science 59: S281-S287.
- /81/ Whomersley, P. and G.B. Picken. 2003. Long-term dynamics of fouling communities on offshore installations in the North Sea. Journal of the Marine Biological Association of the UK. 83: 897-901. Cambridge University Press.
- /82/ COWI/VKI Joint Venture. 1992. Öresund impact assessment. Sub-report nr. 2. The Öresundskonsortiet. Environmental impact assessment for the fixed link across the Öresund.
- /83/ Moore, P.G. 1991. Inorganic particulate suspensions in the sea and their effects on marine animals. - Oceanogr. Mar. Biol. Ann. Rev. 15: 335-363.
- /84/ Popper, A.N., Smith, M.E., Cott, P.A., Hanna, B.W., MacGillivray, A.O., Austin, M.E. and Mann, D.A. 2005. Effects of exposure to seismic airgun use on hearing of three fish species. Journal of the Acoustical Society of America 117 (6): 3958-3971.
- /85/ Engell-Sørensen, K. and Skytt, P.H. 2001. Evaluation of the effect of Sediment Spill from Offshore Wind Farm Construction on Marine Fish. Report to SEAS, Denmark.
- /86/ Johnston, D.D. & Wildish, D.J. 1982. Effect of suspended sediment on feeding by larval herring (*Clupea harengus harengus* L.). Bulletin of Environmental Contamination and Toxicology, vol. 29, 261-267.
- /87/ Rönbäck, P. & Westerberg, H. 1996. Sedimenteffekter på pelagiska fiskägg och gulesäckslarver. Fiskeriverket, Kustlaboratoriet, Frölunda, Sweden.
- /88/ Caddy, J.F. 1973. Underwater observations on tracks of dredges and trawls and some effects of dredging on a scallop ground. Journal of the Fisheries Research Board of Canada, Vol. 30, pp. 173-180.
- /89/ Chapman, C.J., Mason, J. and Drinkwater, J.A.M. 1977. Diving observations on the efficiency of dredges used in the Scottish fishery for the scallop, *Pecten maximus* (L.), Scottish Fisheries Research, Vol. 10.
- /90/ Ramsay, K., Kaiser, M.J. and Hughes, R.N. 1998. Responses of benthic scavengers to fishing disturbance by towed gears in different habitats. Journal of Experimental Marine Biology and Ecology, Vol. 224, pp. 73- 89.
- /91/ Kioerboe, T., Frantsen, E., Jensen, C. and Nohr, O. 1981. Effects of suspended-sediment on development and hatching of herring (*Clupea harengus*) eggs. Estuarine, Coastal and Shelf Science, vol. 13, 107-111.
- /92/ Booman, C., Dalen, J., Leivestad, H., Levsen, A., van der Meeren, T. and Toklum, K. 1996. Effekter av luftkanonskyting på egg, larver og yngel. Undersøkelser ved Havforskningsinstituttet og Zoologisk Laboratorium, UiB. (Engelsk sammendrag og figurtekster). Havforskningsinstituttet, Bergen. Fisken og Havet, nr. 3 (1996). 83 s.
- /93/ McCauley, R.D., Fewtrell, J., and Popper, A.N. 2003. High intensity anthropogenic sound damages fish ears. Journal of the Acoustical Society of America 113: 638-642.
- /94/ Thomsen, F., Lüdemann, K., Kafemann, R. & Piper, W. 2006. Effects of offshore wind farm noise on marine mammals and fish. COWRIE Ltd.
- /95/ Parvin, S.J., Nedwell, J.R. 2006. Underwater noise survey during impact piling to construct the Barrow offshore windfarm. Report Reference: 544R0602.
- /96/ Peña, H., Handegard, N.O. and Ona, E. 2013. Feeding herring schools do not react to seismic air gun surveys. ICES Journal of Marine Science, 70: 1174-1180.
- /97/ Norwegian Oil Industry Association (OLF). 2003. Seismic surveys impact on fish and fisheries by Ingebret Gausland.

- /98/ Browse FLNG Development 2010. Environmental Resources Management 2010. Browse Upstream LNG Development: Light Impact. Appendix F16. Assessment, Report produced for Woodside Energy Limited. Draft Environmental Impact Statement EP BC 2013/7079
- /99/ DNV ENERGY. 2007. Effects of seismic surveys on fish, fish catches and sea mammals. Report for the Cooperation group - Fishery Industry and Petroleum Industry. Report no.: 2007-0512.
- /100/ Forteath, G.N.R., Picken, G.B., Ralph, R. and Williams, J. 1982. Marine growth studies on the North Sea Oil Platform Montrose Alpha. Mar. Ecol. Prog. Ser Vol 8: 61- 68
- /101/ Love, M.S. and Nishimoto, M.M. 2012. Completion of fish assemblage surveys around manmade structures and natural reefs off California. Marine Science Institute, University of California, Santa Barbara, California. BOEM Cooperative Agreement No.: M10AC2001.
- /102/ Scarcella, G., Grati, F. and Fabi, G. 2011. Temporal and spatial Variation of the fish assemblage around a gas platform in the Northern Adriatic Sea, Italy, Turkish Journal of Fisheries and Aquatic Sciences 11: 433-444.
- /103/ Jørgensen, T., Løkkeborg, S. and Soldal, A.V. 2002. Residence of fish in the vicinity of a decommissioned oil platform in the North Sea. ICES Journal of Marine Science. Vol 59, Supplement 1, October 2002, pp S288-S293.
- /104/ Love, M.S., Saiki, M.K., May, T.W. and Yee, J.L. 2013. Whole-body concentrations of elements in three fish species from offshore oil platforms and natural areas in the Southern California Bight. USA Bulletin of Marine Science, Volume 89, Number 3, July 2013 , pp. 717-734(18).
- /105/ Oceans inc. 2013. Monday April 22, 2013. US geological Survey study helps inform rigs-to-reef plans (www. Oceans inc.org).
- /106/ Clarke, D.G and Wilber, D.H. 2000. Assessment of potential impacts of dredging operations due to sediment resuspension. DOER Technical Notes Collection (ERDC TN-DOER-E9), U.S. Army Engineer Research and Development Center, Vicksburg, MS.
- /107/ Moore, P.G. 1977. Inorganic particulate suspensions in the sea and their effects on marine animals. Oceanogr. Mar. Biol. Ann. Rev. 15: 225-363.
- /108/ Levings, C.D. 1982. The ecological consequences of dredging and dredge spoil disposal in Canadian waters. National research council of Canada. NRCC Associate committee on scientific criteria for environmental quality. Publication NRCC No. 18130.
- /109/ Redding, J.M and Schreck, C.B. 1987. Physiological effects in Coho salmon and steelhead of exposures to suspended solids. Trans. Am. Fish. Soc., 116: 737-744.
- /110/ Noggle, C.C. 1978. Behavioural, physiological and lethal effects of suspended sediments on juvenile salmonids. Master Thesis. University of Washington, Seattle, USA.
- /111/ Auld, A.H. and Schubel, J.R. 1978. Effects of suspended sediment on fish eggs and larvae: A laboratory assessment. Estuarine and Coastal Marine Science, 6. Page 153-164.
- /112/ Birklund, J. and Wijsman, j.W.M. 2005. Agregate Extraction: A Review on the effect on ecological funktions. - Prepared for: EC Fifth Framework Programme Project SANDPIT: 54
- /113/ Rönbäck, P. and Westerberg, H. 1996. Sedimenteffekter på pelagiska fiskägg och gulesäckslarver. Fiskeriverket, Kustlaboratoriet, Frölunda, Sweden.
- /114/ Kioerboe, T., Frantsen, E., Jensen, C. and Nohr, O. 1981. Effects of suspended-sediment on development and hatching of herring (*Clupea harengus*) eggs. Estuarine, Coastal and Shelf Science, vol. 13, 107-111.
- /115/ Caddy, J.F. 1973. Underwater observations on tracks of dredges and trawls and some effects of dredging on a scallop ground. Journal of the Fisheries Research Board of Canada, Vol. 30, pp. 173-180.
- /116/ Chapman, C.J., Mason, J. and Drinkwater, J.A.M. 1977. Diving observations on the efficiency of dredges used in the Scottish fishery for the scallop, *Pecten maximus* (L.), Scottish Fisheries Research, Vol. 10
- /117/ Ramsay, K., Kaiser, M.J. and Hughes, R.N. 1998. Responses of benthic scavengers to fishing disturbance by towed gears in different habitats. Journal of Experimental Marine Biology and Ecology, Vol. 224, pp. 73- 89.
- /118/ Moles, A. and Norcross, B.L. 1995. Sediment preference in juvenile pacific flatfishes. Netherlands Journal of Sea Research 34 (1-3). Page 177-182.

- /119/ Groot, S.J. De. 1980. The consequences of marine gravel extraction on the spawning of herring, *Clupea harengus*. Linne. Journal of Fish Biology, 16: 605–611.
- /120/ Sveegaard S., Teilmann J., Tougaard J., & Dietz R. (2011) High-density areas for harbor porpoises (*Phocoena phocoena*) identified by satellite tracking. Marine Mammal Science, 27, 230–246.
- /121/ Maersk Oil. 2014. Environmental Management System: Danish Business Units. DK-HSE-PRO-0026 Rev 3.0.
- /122/ Richardson, W.J., C.R. Greene Jr., C.I. Malme, and D.H. Thomson. (1995). Marine Mammals and Noise. Academic Press, San Diego. 576 p.
- /123/ Boucher, G. 1985. Long term monitoring of meiofauna densities after the Amoco Cadiz oil spill. Mar Pollut Bull 16: 328–333.
- /124/ Dauvin, J.C. 1998. The fine sand *Abra alba* community of the Bay of Morlaix twenty years after the Amoco Cadiz oil spill. Mar Pollut Bull 36: 669–676.
- /125/ Volkman, J.K., Miller, G.J., Revill, A.T. and Connell, D. W. 1994. Oil spills. Part 6. In: Environmental Implications of Offshore Oil and Gas Development in Australia – The Findings of an Independent Scientific Review. Eds. Swan, J. M., Neff, J. M. and Young, C. P. Australian Petroleum Exploration Association, Sydney. pp. 509-695.
- /126/ Davies, J.M. 1980. Some effects of oil-derived hydrocarbons on a pelagic food web from observations in an enclosed ecosystem and a consideration of their implications for monitoring. Rapp. P.-v. Reün. Cons. int. Explor. Mer. vol. 179, pp. 201-211.
- /127/ Neff, J.M. 1991. Water quality in Prince William Sound and the Gulf of Alaska.
- /128/ Anon (1985). Oil in the sea. Inputs, fates and effects. National Academy Press, Washington D.C 1985.
- /129/ Varelaa, M. et al. 2006. The effect of the "Prestige" oil spill on the plankton of the N-NW Spanish coast. Marine Pollution Bulletin Volume 53, Issues 5-7, 2006, Pages 272-286.
- /130/ Wei, C.L., Rowe, G.T., Esobar-Briones, E., Nunnally, C., Soliman, Y. and Ellis, N. 2012. Standing stocks and body size of deep-sea macrofauna: Predicting the baseline of 2010 Deepwater Horizon oil spill in the northern Gulf of Mexico. Deep-Sea Research I, 69, 82-99.
- /131/ Neff, J.M., Anderson, J.W. 1981. Response of marine animals to petroleum and specific petroleum hydrocarbons. Halsted Press. New York.
- /132/ Hjermmann, D.O., Melsom, A., Dingsor, G.E., Durant, J.M., Eikeset, A.M., Roed, L.P., Ottersen, G., Storvik, G. and Stenseth, N.C. 2007. Fish and oil in the Lofoten-Barents Sea system: synoptic review of the effect of oil spills on fish populations. Marine Ecology Progress Series, 339, 283-289.
- /133/ Edwards, R. and White, I. 1999. The Sea Empress Oil Spill: Environmental Impact and recovery. In International Oil Spill Conference (IOSC) Conference Proceedings 199, 97-102).
- /134/ Fodrie, F.J. and Heck, K.L.Jr. 2011. Response of Coastal Fishes to the Gulf of Mexico Oil Disaster. PLoS ONE 6(7).
- /135/ OLF. 2008. Metodikk for miljørisiko på fisk ved akutte oljeutslipp. Oljeindustriens Landsforening. DNV Energy.
- /136/ Ramboll, 2012. Dan Quantitative Risk Assessment (QRA), Prepared for Maersk Oil.
- /137/ Ramboll, 2012. Kraka Quantitative Risk Assessment, Prepared for Maersk Oil.
- /138/ Marine mammal working group. 2015. Marine mammals and underwater noise in relation to pile driving – Working Group 2014. Revision 2, 21.01.2015. For Energinet.dk.
- /139/ Balle, J.D., Clausen, K.T., Mikkelsen, L., Wisniewska, D.M. and Teilmann, J. 2014. Harbour porpoises and noise around an operating oil and gas production platform in the North Sea – Status report. DCE.
- /140/ Henriksen, O.D., Maxon, C. and Degn, U. 2005. Underwater sound from offshore drilling activities. Potential effects on marine mammals. DDH consulting 362 05 124, rev4.
- /141/ DCE. 2015. <http://envs.au.dk/videnudveksling/luft/emissioner/>. Accessed April 29<sup>th</sup> 2015.
- /142/ Tougaard, J. 2014. Vurdering af effekter af undervandsstøj på marine organismer. Del 2 – Påvirkninger. Aarhus Universitet, DCE – Nationalt Center for Miljø og Energi, 51 s. -Teknisk rapport fra DCE - Nationalt Center for Miljø og Energi nr. 45 <http://dce2.au.dk/pub/TR45.pdf>
- /143/ Teilmann J. & Carstensen J. (2012) Negative long term effects on harbour porpoises from a large scale offshore wind farm in the Baltic - evidence of slow recovery. Environmental Research Letters, 7, doi:10.1088/1748-9326/7/4/045101.
- /144/ COWI. 2012. Assessment of impacts on marine mammals of seismic survey in DUC area in 2012. AU28981.

- /145/ OSPAR. 2009. Overview of the impacts of anthropogenic underwater sound in the marine environment.
- /146/ Stemp, R. (1985): Observations on the effects of seismic exploration on seabirds. p. 217-233 In: G.D. Greene, F.R. Engelhardt, and R.J. Peterson (eds.), Proceedings of workshop on effects of explosives use in the marine environment. Cdn. Oil and Gas Admin., Env. Prot. Branch, Tech. Rep. No. 5. Ottawa.
- /147/ Evans, P.G.H., Lewis, E.J. and Fisher, P. (1993): A study of the possible effects of seismic testing upon cetaceans in the Irish Sea. Rep. by Sea Watch Foundation, Oxford, to Marathon Oil UK Ltd. Aberdeen. 35 p.
- /148/ Santos, M.N., Monteiro, C.C. & Lassère, G. (1996). Finfish attraction and fisheries enhancement on artificial reefs: a review. In: Jensen, A.C. (Ed.) European artificial reef re-search. Proceedings of the 1st EARRN conference, Ancona, Italy, March 1996. Pub. South-ampton Oceanography Centre: 97-114.
- /149/ Valdemarsen, J.W. (1979). Behaviour aspects of fish in relation to oil platforms in the North Sea. ICES C.M., B:27
- /150/ Callaway, R.J., Alsvåg, I. de Boois, J., Cotter, A., Ford, G.H., Hinz, S. Jennings, I. Kröncke, J. Lancaster, G. Piet, P. Prince and As. Ehrich (2002). Diversity and community structure of epibenthic invertebrates and fish in the North Sea. ICES Journal of Marine Science 59: 1199-1214.
- /151/ Thompson, P., Brookes, K., Cordes, L., Barton, T., Cheney, B. and Graham, I. 2013. Assessing the potential impact oil and gas exploration operations on cetaceans in the Moray Firth. Novembre 18<sup>th</sup> 2013.
- /152/ OSRL. 2015. Oil spill modelling report: Danish Business Units. CONS1050-R01.
- /153/ DCE. 2015. Disturbance Effects on the Harbour Porpoise Population in the North Sea (DEPONS): Status report on model development.
- /154/ Naturstyrelsen. Danmarks Havstrategi. Miljømålsrapport.
- /155/ OSPAR. 2012. Guidelines in support of Recommendation 2012/5 for a Risk-based Approach to the Management of Produced Water Discharges from Offshore Installations.
- /156/ COWI/DHI joint venture. 2001. The great belt link. The monitoring programme 1987-2000.
- /157/ Nord Stream. 2009. Environmental Impact Assessment: Documentation for Consultation under the Espoo Convention Nord Stream Espoo Report: Key Issue Paper Seabed Intervention: Works and Anchor Handling
- /158/ ITOPF. 2014. Members Handbook.
- /159/ Maersk Oil. 2015. Environmental and Social Impact Management in Projects Standard. MOG-HSE-ENV-STD-0042, rev1

## **APPENDIX 1 TECHNICAL SECTIONS**