



Impacts of offshore wind farms on waves and sediment transport

Environmental mapping and screening of areas for offshore wind in Denmark

Danish Energy Agency

Date: 28. November 2025

Rev.no.	Date	Description	Prepared by	Verified by	Approved by
01	20250509	Rev 1	FRGR/RUGA/TEB	TEB	RHO
02	20250918	Rev 2	ANSL/TEB	TEB	LRM
03	20251128	Rev 3	FRGR/TEB	TEB	LRM

Contents

1.	Preface.....	6
2.	Summary.....	7
3.	Introduction.....	8
3.1.	Scope of work.....	8
3.2.	Coordinate System.....	8
4.	Background data.....	9
4.1.	Bathymetry.....	9
4.2.	Wind forcing.....	10
4.3.	Wave observations.....	10
5.	Methodology.....	15
5.1.1.	Spectral wave model.....	15
6.	Model setup and calibration.....	17
6.1.	Computational grid.....	17
6.2.	Model set-up.....	17
6.3.	Model calibration.....	18
7.	Results baseline.....	26
8.	Impact on waves.....	28
8.1.	Wave impact from wind farm scenarios.....	28
8.2.	Annual and monthly variation.....	31
8.3.	Conclusion on wave impact.....	37
9.	Impact on sediment transport.....	38
9.1.	Introduction.....	38
9.1.1.	Løkken to Hirtshals.....	39
9.1.2.	Ferring to Hvide Sande.....	40
9.1.3.	Hjelm Bugt.....	40
9.2.	Data.....	41
9.2.1.	Profile orientation.....	41
9.2.2.	Sediment grain size.....	42
9.2.3.	Waves.....	42
9.3.	Methods.....	42
9.3.1.	Wave height conversion.....	42
9.3.2.	Limitations.....	43
9.4.	Results.....	43
9.5.	Uncertainty analysis.....	50

9.6.	Conclusion on sediment transport impact.....	50
10.	References.....	51

Appendix:

Appendix A	Available measurements of waves
Appendix B	Time-series of significant wave heights
Appendix C	Calibration of significant wave height
Appendix D	Wave roses at selected locations
Appendix E	Wind farms spatial impact on significant wave height
Appendix F	Wind farms spatial impact on significant wave height, Danish waters
Appendix G	Waves, annual and monthly variations at selected locations

NOMENCLATURE

Abbreviation	Description
BHS	German Federal Maritime and Hydrographic Agency
DEM	Digital Elevation Model
DTM	Digital Terrain Model
H_{m0}	Modelled Significant Wave Height
H_s	Measured Significant Wave Height
MEA	Mean Absolute Error
MSL	Mean Sea Level
MWD	Mean Wave Direction
NCCS	Norwegian Center for Climate Services
RMSE	Root Mean Squared Error
SI	Scatter Index
SMHI	Swedish Meteorological and Hydrological Institute
T_p	Peak Wave Period
UTM	Universal Transverse Mercator
VD	Vertical Datum
WGS84	World Geodetic System 1984

1. Preface

This report contributes to the project “*Environmental mapping and screening of areas for offshore wind in Denmark*” initiated in 2022 by the Danish Energy Agency. The project aims to support the long-term planning of offshore wind farms by providing a comprehensive overview of the combined offshore wind potential in Denmark. It is funded under the Finance Act 2022 through the programme “Investeringer i et fortsat grønnere Danmark” (Investing in the continuing greening of Denmark). The project is carried out by NIRAS, Aarhus University (Department of Ecoscience) and DTU Wind.

The overall project consists of four tasks defined by the Danish Energy Agency (<https://ens.dk/energikilder/planlaegning-af-fremtidens-havvindmoelleparker>):

1. Sensitivity mapping of nature, environmental, wind and hydrodynamic conditions.
2. Technical fine-screening of areas for offshore wind based on the sensitivity mapping and relevant technical parameters.
3. Assessment of potential cumulative effects from large-scale offshore wind development in Denmark and neighbouring countries.
4. Assessment of barriers and potentials in relation to coexistence.

This report addresses one component of each of the first three tasks. Specifically, it provides an overview of areas within Danish offshore regions that are likely to be particularly vulnerable to offshore wind farm development regarding waves based on wind data provided by DTU Wind, considering the wind in a scenario without wind farms as well as two scenarios with wind farms in the years 2021 and 2030, respectively.

The spectral wave modelling was conducted based on wind data from a single year (2019), which hinders inter-annual comparisons. However, in order to address the impact of wind farms on waves, the natural annual/monthly variation have been analyzed with the use of existing wave data for the years 1988 to 2024 at several locations. Section 4 specifies the data sources used in the sensitivity mapping for waves and outlines additional existing data. Similar for the sediment transport capacity analysis is listed in Section 9.2.

It is important to recognize that sensitivity mapping serves as a dynamic tool, which can be updated as new data becomes available.

2. Summary

The Danish Energy Agency commissioned this report to support the long-term planning for offshore wind development in Denmark. The report focuses on the potential impact on waves and sediment transport due to current and future offshore wind development in the North Sea, Kattegat, inner Danish Waters and western Baltic Sea.

Large-scale development of offshore wind can affect hydrodynamics both within and beyond the wind farms and may affect factors such as water transport, current conditions, waves, resuspension, sediment transport, salinity, and temperature. These hydrodynamic alterations have the potential to impact environmental conditions in areas within and around wind farms. Thus, investigating the potential impact on hydrodynamic factors such as waves and sediment transport due to offshore wind development is an important part in the assessment of potential environmental and cumulative impacts.

In order to address this, the report presents the results from both wave modelling and sediment transport modelling of three scenarios: 1) a baseline scenario with no wind farms present, 2) a current (2021) wind farm scenario and 3) a future 2030 wind farm scenario. The wave modelling was conducted for a large area covering the North Sea and the western Baltic Sea, whereas the sediment transport modelling was conducted for three stretches along the Danish coastline. The input data for the wave model are based on simulated wind data from the year 2019 and are calibrated using data from buoy recordings for the same period. The impact on waves is assessed through the impact on the mean wave direction and the significant wave height.

The results from the wave modelling indicate that current wind farms (in operation in 2021) exert minimal influence on wave conditions. Compared to the results from the current wind farm scenario, the future large-scale offshore wind farm development (2030 scenario) generally causes larger reductions in significant wave heights. However, upon examining the annual variation in wave climate, it is observed that the changes in mean significant wave height attributable to the wind farms are significantly smaller than natural fluctuations for the west coast of Jutland and in Kattegat. In contrast, the reduction in significant wave height observed in the Baltic Sea due to wind farms approaches the annual and monthly standard deviation. Generally, for the North Sea, Kattegat and the Baltic Sea, the natural mean wave direction varies by several degrees, and the impact of the wind farms is considered negligible in comparison to these inherent variations.

The sediment transport modelling results show, similarly to the impact on waves, that the impact on sediment transport is larger for the future wind farm scenario (2030) compared to the current wind farm scenario. Comparison between the selected coastal stretches indicate that the impact at Hjelm Bugt in the Baltic Sea is larger compared to the impact along the west coast of Jutland bordering the North Sea, which may indicate that the wave climate in the Baltic Sea is more sensitive to impacts from future offshore wind farm development. It is important to note, however, that the annual variation in the wave climate indicates a larger change compared to the impact introduced by wind farms.

3. Introduction

Large-scale offshore wind farms have been documented to affect hydrodynamics both within and beyond the wind farms, extending over many kilometres (van Duren, et al., 2021; Arneborg, et al., 2024). These effects include a reduction in wind resource efficiency behind large wind farms due to the wake effect, along with localized impacts from turbine foundations.

Water transport, current conditions, waves, resuspension, sediment transport, salinity, and temperature may also be affected, causing changes in mixing and stratification patterns. These hydrodynamic alterations have the potential to impact environmental conditions and exacerbate oxygen depletion in areas within and around wind farms (Daewel, et al., 2022; Maar, et al., 2025).

3.1. Scope of work

The aim of this report is to produce a sensitivity map showing how the presence of wind farms in the North Sea, Kattegat and Baltic Sea affects the wave climate and for three coastal sections the sediment transport capacity. For this purpose, a spectral wave model was used to simulate the following three cases:

- 1) **Baseline scenario:** a scenario without any wind farm installed based on the wind in 2019.
- 2) **Current scenario, 2021:** current situation including existing Danish offshore wind turbines utilising the wind in 2019.
- 3) **Future scenario, 2030:** scenario following the future large-scale offshore wind development in Danish sea areas: the North Sea, Kattegat, Belts and the Baltic Sea utilising the wind in 2019.

The scenarios are executed based on calculation from DTU Wind (Hahmann, et al., 2025a; 2025b) and using the 2D MIKE Spectral Wave model by DHI.

Furthermore, to assess the impact of changes in wave climate on coastal longshore sediment transport, the sediment transport capacity calculated with use of LITPACK was evaluated along three distinct coastal sections for the Baseline scenario, as well as for the Current (2021) and Future (2030) wind farm scenarios.

3.2. Coordinate System

Unless stated otherwise, the Vertical Datum (VD) throughout the report is Mean Sea Level (MSL), while the geographic coordinate system is WGS84 EPSG:7416 ETRS89 UTM32N.

4. Background data

The upcoming sections provide the background information utilized for numerical modelling. This chapter encompasses descriptions of metocean and meteorological conditions, such as wave observations and bathymetric data. The wind data is described in Hahmann, et al., (2025a; 2025b).

4.1. Bathymetry

One of the main inputs to the spectral wave model is the Digital Terrain Model (DTM). A DTM is a digital representation of the terrain or ground surface of a specific area or location. It is a three-dimensional model that provides information about the elevation, slope, and other topographic features of the terrain.

DTMs are created using Digital Elevation Models (DEMs), which are datasets that provide information about the elevation of the ground surface.

The DTM used for the model originates from the combination of two different sources of data: the European Marine Observation and Data Network 2023 (EMODnet, 2023), reported in Figure 4.1, and the digital nautical charts obtained from the application MIKE C-Map from DHI, shown in Figure 4.2.




Figure 4.1: Extent of the model area (black box) with EMODnet Bathymetry used for the interpolation of the mesh for the spectral wave model.




Figure 4.2: MIKE C-Map scatter data of water depths for the North Sea and Baltic Sea used for the interpolation of the mesh for the spectral wave model.

4.2. Wind forcing

Wind and pressure used as forcings for the simulations have been developed by DTU Wind and are described in detail in Hahmann, et al. (2025a; 2025b).

4.3. Wave observations

To validate/calibrate the spectral wave model, wave buoy recordings related to the simulated period (January–December 2019) are key for an accurate description of the physical processes. Four sources of data provided wave measurements within the area of study (see Figure 4.3 for buoy locations):

- The Copernicus Marine In Situ TAC: The In Situ TAC component of the Copernicus Marine Service ensures consistent and reliable access to a range of in situ data for the purpose of service production and validation;
- The Swedish Meteorological and Hydrological Institute database (SMHI);
- The Norwegian Center for Climate Services (NCCS);

- The German Federal Maritime and Hydrographic Agency (BSH) which provides in the “Report on the oceanographic conditions at site O-1.3” (German Federal Maritime and Hydrographic Agency (BSH), 2021) a description of the marine monitoring instrument that records at a location close to the Arkona Basin.




Figure 4.3: Location of the retrieved wave measurements.

The retrieved observed wave data are characterized by significant wave height (H_s), peak period (T_p), and mean wave direction (MWD). However, although all stations recorded H_s , only a few provided information on period and direction.

The collected measurement data underwent a comprehensive quality check to determine which stations provided reliable time series of wave parameters and which could only be used to evaluate the general patterns of the simulated wave data. A description of each recovered measurement station, i.e. location, time-zone and coverage (%) is reported in Appendix A.

Three representative stations for each side were selected to be following reported, while the time-series of H_s and T_p for the other stations are placed in Appendix B.

The stations selected for comparison and reported in the main body of this report are listed in Table 4.1 and shown in Figure 4.4.

Table 4.1: Stations selected for calibration of the spectral wave model.

Station ID	Longitude [deg]	Latitude [deg]	Time	Coverage (%)
NO_TS_MO_F3platform	4.727	54.853	UTC 0	95
NO_TS_MO_A122	3.816	55.417	UTC 0	85
NO_TS_MO_6202112	7.15	55.200	UTC 0	100




Figure 4.4: Location of the stations selected for calibration of the SW model.

The time series of significant wave height and peak period (when available) for each of the selected stations is reported in Figure 4.5 to Figure 4.7.




Figure 4.5: Significant wave height (top) and peak period (bottom) recorded at the station **NO_TS_MO_F3platform**.




Figure 4.6: Significant wave height (top) and peak period (bottom) recorded at the station **NO_TS_MO_A122**.




Figure 4.7: Significant wave height (top) and peak period (bottom) recorded at the station **NO_TS_MO_6202112**.

The comparison between the wave parameters recorded at three selected stations and the modelling results are presented and discussed in Section 6.3.

It is important to highlight that in the case of the retrieved Arkona Basin station, there is a 30-minute interval between recordings, and the time steps never correspond to the simulated time step. The data was interpolated in order to be compared to the spectral wave model results.

5. Methodology

A spectral wave model was used to simulate the wave climate in Danish waters to assess the impact of existing and future offshore wind farms. For each scenario, a distinct wind forcing file was applied, all reflecting wind conditions from 2019, a “typical” year that is representative of the climatological conditions, but accounting for varying configurations of scenarios for wind farms; Current (2021 wind farms) and Future (2030 wind farms), within the study area (Hahmann, et al., 2025a; 2025b). Figure 5.1 presents an overview the locations of the wind farms includes in the current (2021) and future (2030) scenarios, respectively.




Figure 5.1: Overview of the location of the currently present wind farm parks (orange) and of the future large-scale development in the North Sea, Kattegat, and the Baltic Sea (green).

Prior to assessing potential impacts, the base model is calibrated using the collected wave measurements mentioned above, see 4.3.

5.1.1. Spectral wave model

MIKE 21 Spectral Wave Model (SW) is a computer-based simulation tool used for analyzing and predicting wave characteristics in coastal and offshore environments.

The MIKE 21 SW utilizes spectral wave theory to simulate the transformation and propagation of ocean waves. It takes into account various factors such as wind, currents, bathymetry (seafloor topography), and wave interactions to accurately predict wave height, direction, period, and other wave parameters.

The MIKE 21 SW module is capable of modelling the growth, decay, and transformation of wind-sea and swell waves in coastal and offshore areas. Its engine includes the following physical processes:

- Wave growth by wind,
- Non-linear wave-wave interaction,
- Wave-current interaction,
- Dissipation of energy due to white capping,
- Dissipation of energy due to bottom friction,
- Dissipation of energy due to depth-induced wave breaking,
- Refraction and shoaling due to variation in water depth,
- Effect of time-varying water depth and currents,
- Effect of ice coverage,
- Diffraction,
- Reflection and
- Influence of structures such as Wave/Tidal Energy Converters (WEC/TEC)

The main computational features in MIKE 21 SW are:

- Source functions based on state-of-the-art 3rd generation formulations,
- Fully spectral and directionally decoupled parametric formulations,
- In-stationary and quasi-stationary solutions,
- Optimal degree of flexibility in describing the bathymetry and the ambient flow conditions using depth-adaptive and boundary fitted unstructured mesh,
- Coupling with hydrodynamic flow module for modelling wave-current interaction and time-varying water depth,
- Flooding and drying in connection with time-varying water levels,
- Extensive range of model output parameters (e.g. wind-sea, swell, air-sea interaction parameters, radiation stresses, wave spectra, etc.) and
- Parallelization using Open MP and MPI techniques.

The model uses a flexible mesh based on unstructured triangular or quadrangular elements and applies a finite volume numerical solution technique. The 2024 version of MIKE 21 SW was used in the present study. Further details are provided in DHI (2021).

6. Model setup and calibration

This section introduces the core of the calculations in the present study: DHI's MIKE 21 SW module and the SW model built specifically for it.

6.1. Computational grid

Figure 6.1 shows the computational mesh and boundaries of the SW model. The resolution of the SW model was selected as a balance between maintaining reasonable computational time.

The resolution progressively varies from the offshore area of the North Sea to the internal region between Sweden and Denmark, where the majority of the offshore wind farms are situated. Offshore and along the coasts of Great Britain and Germany, the resolution is 2 km², which decreases to 1 km² as it approaches the Baltic Sea.




Figure 6.1: Computational mesh and boundaries defined for the spectral wave model of the North Sea, Kattegat, Belts and Baltic Sea.

6.2. Model set-up

The model setup is summarized in Table 6.1. When not otherwise specified, the configuration of the SW model follows the default setup.

Through a calibration process, the best fit between modelled data and measurements was obtained by increasing the value of the air-water interaction parameters so that the energy dissipation within the domain decreased.

Table 6.1: Summary of the SW model settings.

Setting	Value
MIKE Version	MIKE 2024
Mesh resolution	6,417 km ² (at the boundaries) to 23 km ² (at the project area)
Simulation period	Jan 1st 2019 00:00 to Dec 31th 2019 23:00
Spectral Formulation	Fully Spectral formulation, Instationary Formulation
Energy Transfer	Included quadruplet-wave interaction
Growth parameter	1.3
Wave age tuning parameter	0.011
Background Charnock parameter	0.01
Atmospheric forcing	DTU generated Wind
Boundary Conditions	ERA5

6.3. Model calibration

Time-series of three wave parameters (modelled significant wave height (H_{m0}), peak wave period (T_p) and mean wave direction (MWD)) were extracted at the wave buoys' locations and validated against the available measurements to calibrate the model and obtain an accurate description of the physical processes involved in the wave dynamics and propagation.

Three representative stations were selected to be reported in the following, while the comparison between the model results and the other stations is placed in Appendix C. The stations selected for comparison and reported in the present section are listed in Table 4.1.

The comparison between final model results and measurements is presented in terms of time-series, scatter plots, and statistical coefficients such as:

- The **Correlation Coefficient**, which quantifies the strength of a linear relationship between two variables, with values from -1 to 1. A value of -1 indicates a perfect negative correlation, where one variable increases as the other decreases, while a value of 1 represents a perfect positive correlation, where both variables move in the same direction;
- The **Bias**, which refers to a systematic error that causes an estimator to deviate from the true population value. Bias occurs when the method of collecting, measuring, or interpreting data consistently skews results in a particular direction, leading to inaccurate or misleading conclusions;
- The **Mean Absolute Error** (MAE), representing the average magnitude of errors in a model's predictions, without considering their direction (positive or negative). It ranges from 0 to infinity. A lower MAE indicates a model with more accurate predictions;
- The **Root Mean Square Error** (RMSE), that RMSE reflects the average magnitude of prediction error, with larger errors given more weight due to the squaring. The RMSE can range from 0 to infinity, and a lower RMSE indicates more accurate predictions;
- The **Scatter Index** (SI) is a standard metric for wave model intercomparison and represents a normalized measure of error. The SI can be presented as percentage. Lower values of SI correspond to a better match between observations and modelled values, and in the context of significant wave height, reports of the scatter index (SI) in the literature range between from 0 to 20% (0.20) for hindcasts with sophisticated models and high quality wind fields (Howard, et al., 2009).

The statistical coefficients are presented in Table 6.2, while the comparison of the time-series of measurements and modeled parameters is presented in Figure 6.1 to Figure 6.12 and in Appendix C.




Figure 6.1: **Station ID NO_TS_MO_F3platform** - Comparison model vs. measurements of yearly time series of the significant wave height.




Figure 6.2: **Station ID NO_TS_MO_F3platform** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-02-2019 to 15-02-2019.




Figure 6.3: **Station ID NO_TS_MO_F3platform** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 15-02-2019 to 01-03-2019.




Figure 6.4: **Station ID NO_TS_MO_F3platform** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-03-2019 to 15-03-2019.




Figure 6.5: **Station ID NO_TS_MO_A122** - Comparison model vs. measurements of yearly time series of the significant wave height.




Figure 6.6: **Station ID NO_TS_MO_A122** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-01-2019 to 15-01-2019.




Figure 6.7: **Station ID NO_TS_MO_A122** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 15-01-2019 to 01-02-2019.




Figure 6.8: **Station ID NO_TS_MO_A122** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-02-2019 to 15-02-2019




Figure 6.9: **Station ID NO_TS_MO_6202112**- Comparison model vs. measurements of yearly time series of the significant wave height.




Figure 6.10: **Station ID NO_TS_MO_6202112** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-06-2019 to 15-06-2019.




Figure 6.11: **Station ID NO_TS_MO_6202112** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-08-2019 to 15-08-2019.




Figure 6.12: **Station ID NO_TS_MO_6202112** - Comparison model vs. measurements: time series of the significant wave height (top) and peak wave period (bottom). Period: 01-09-2019 to 15-09-2019.

Table 6.2: Correlation coefficient, Bias, MAE, RMSE and SI for the comparison between simulated and measured significant wave height at three selected locations (stations).

Station ID	Parameter: Sign. Wave Height				
	Correlation Coefficient	Bias	MAE	RMSE	SI
NO_TS_MO_F3platform	0.93	+0.11	0.22	0.33	0.23
NO_TS_MO_A122	0.94	-0.05	0.28	0.37	0.20
NO_TS_MO_6202112	0.93	+0.00	0.21	0.29	0.20

Despite the model's tendency to overestimate some of the peak values of significant wave height, the comparison figures indicate that the overall pattern of model results aligns well with the measurements. This is evidenced by a scatter index (SI) close to 0.2 and a correlation of 0.93 to 0.94, which is regarded as indicative of good agreement. Since the comparison between scenarios with and without the wind farm is relative, the resulting impact of the model's tendency to overestimate peak values of significant wave height is considered negligible and will hence not influence the overall conclusions drawn from the modelled results.

7. Results baseline

To better understand the wave climate in Danish waters, the modelled significant wave height (H_{m0}) was extracted from the results of the baseline case, thus without taking into account the presence of wind farms in the domain, at three locations: one in the Baltic Sea, one in Kattegat and one offshore Jutland, in the North Sea west of Hvide Sande. The locations are shown in Figure 7.1 with details in Table 7.1.




Figure 7.1: Locations of extraction of significant wave height and mean wave direction for Baseline case assessment.

Table 7.1: Description of the locations of extraction of wave conditions.

ID	X	Y	Area
P1	762483.01	6102118.35	Baltic Sea
P2	631675.33	6312239.29	Kattegat
P3	396695.01	6208658.55	North Sea

The annual wave roses for the three locations are shown in Figure 7.2, which highlights the following observations:

- For the Baltic Sea location (P1, first rose on the left), while the predominant wave direction in 2019 was from the southwest, wave directions from west to northwest were equally significant in terms of magnitude,
- For the Kattegat area (P2, middle rose), data reflects minimal directional dependency, likely due to its sheltered conditions and the fetch, in contrast to the more defined directional patterns observed at the other two sites, and
- Like P1, the North Sea location (P3, off the Jutland coast) shows that the primary wave directions extend from north-northwest to south-southwest.

The annual wave roses for the three locations in Figure 7.1 are reported in Figure 7.2.




Figure 7.2: Annual wave rose for selected locations, from left to right, P1 (Baltic Sea), P2 (Kattegat) and P3 (North Sea).

8. Impact on waves

As previously mentioned, the assessment of the wave climate impact is based on a single year (2019) and compares the current (2021) wind farm scenario and with a future (2030) wind farm scenario with respect to the impact on waves. This comparison between the two scenarios is presented in the below section 8.1. To evaluate the significance of these impacts, the general wave climate and its variations on an annual and monthly basis have been analysed and compared to the identified effects, which is presented in section 8.2.

8.1. Wave impact from wind farm scenarios

The impact of current and future wind energy development on wave climate is evaluated by comparing the differences in significant wave heights between:

- Baseline and Current wind farms, i.e. year 2021,
- Baseline and Future wind farms within the considered domain, i.e. year 2030.

The difference is calculated for both the annual average and the monthly average, but only the annual results are included in the main body of this report (see Figure 8.2). In addition to the overall results for the studied domain, a detailed view of the area of primary interest, specifically certain waters within the Danish Exclusive Economic Zone (EEZ), is provided (see Figure 8.3 and Figure 8.4), where the results are shown without and with wind turbines marked respectively.

The monthly differences, for the whole domain and for the Danish Waters, can be found in Appendix F.

Based on the processed results, it is generally noted that:

- Referring to the whole domain (Figure 8.1 and Figure 8.2), the already installed wind farms generate little to no disturbance compared to the wave conditions calculated in the absence of offshore wind farms. Indeed, the decrease in annual averaged significant wave height ranges from -0.1 m to 0.0 m,
- With regards to the area of primary interest, the waters within the Danish (EEZ) (Figure 8.3 and Figure 8.4), the already installed wind farms do not affect the significant wave height (EEZ is not shown on figures, and it is evident from Figure 8.3 and Figure 8.4 that only very small areas categorized with a -0.05 m to -0.03 m change in the significant wave height is shown within the Danish EEZ),
- In reference to the whole domain (Figure 8.1 and Figure 8.2), the Future (2030) wind farm scenario affects the baseline wave conditions to a larger degree compared to the current wind farm scenario. The decrease in annual averaged significant wave height ranges from -0.2 m to 0.0 m, where the largest decrease is reached in the area between the Netherlands and England because of the high density of wind turbines (see figure 4 in Hahmann, et al. (2025b)) and
- For the Future (2030) wind farm scenario in the waters within the Danish (EEZ) (Figure 8.3 and Figure 8.4), the annual average of significant wave height is affected by the installed wind turbines in the North Sea, Baltic Sea and in the Kattegat area, with a maximum change in significant wave height of approximately -0.1 m (only a small area within the Danish EEZ in the Baltic Sea is modelled to experience a change in significant wave height of -0.15 m to -0.1 m). Along the coast of Jutland the maximum change in wave height is less than -0.05 m.

With regard to the observed seasonal variation, in relation to the year 2019, the following are observed:

- In January, where higher significant wave heights (H_{m0}) generally are observed compared to other months, the changes for the Future scenario exceed -0.2 meters, particularly along the coastlines of the Netherlands and Great Britain. For the Current scenario, a maximum of -0.1 m is observed in January only within the wind farms in the southern area of the North Sea and Baltic Sea,

- West of Jutland, waves predominantly come from west and northwest, except in April, June and August. In April, the prevailing directions are northeast to southeast, while in June and August the prevailing direction is Southwest (see Appendix D). This variability results in varied patterns of wave reduction in response to the presence of wind farms along the western coast of Jutland,
- For both wind farm scenarios, the prevalence of a specific direction, or a range of directions, is less pronounced in the Kattegat and the Baltic Sea than the North Sea, as it is observed in Figure 7.2 and
- Based on the monthly wave roses for P1 (located in the Baltic Sea) reported in Appendix D, a shift is observed in the prevailing wave direction from westerly to easterly directions from April to June for both wind farm scenarios. The pattern shows that reduced wind speeds are associated with lower wave heights, particularly in areas near the coastline from April to July.

Based on the monthly wave roses for P2 reported in Appendix D, the following seasonal variations are observed for both wind farm scenarios:

- The highest waves come from Northwest to Southwest between February to October, except for April which, as November, shows an important southeasterly component. This is reflected in the pattern of influence of the reduction of wind speed on the wave heights. Indeed, the reduction appears to affect the area closer to the coastline in April and November,
- In general, during the summer period (from May onwards), the reduction in wave height becomes less pronounced as an overall milder wave climate is observed, with a maximum decrease in significant wave height down to -0.2 meters (see Appendix E and Appendix F) and
- In general, the seasonal variation is more easily spotted for the Future scenario due to the larger impact.




Figure 8.1: **Whole model domain - Baseline and Baseline vs. Scenarios.** Left: annual average significant wave height (m), Mid: Scenario 2021 minus Baseline, Right: Scenario 2030 minus Baseline.




Figure 8.2: **Whole model domain - Baseline and Baseline vs. Scenarios.** Left: annual average significant wave height (m), Mid: Scenario 2021 minus Baseline, Right: Scenario 2030 minus Baseline. "Mid" and "Right" marked with the wind farms.




Figure 8.3: **Danish Waters - Baseline and Baseline vs. Scenarios.** Left: annual average significant wave height (m), Mid: Scenario 2021 minus Baseline, Right: Scenario 2030 minus Baseline.




Figure 8.4: **Danish Waters - Baseline and Baseline vs. Scenarios.** Left: annual average significant wave height (m), Mid: Scenario 2021 minus Baseline, Right: Scenario 2030 minus Baseline. "Mid" and "Right" marked with the wind farms.

8.2. Annual and monthly variation

The influence of wind farm installations in Danish waters was further evaluated by extracting and comparing wave parameters, specifically, significant wave height (H_{m0}) and mean wave direction (MWD), from the "Atlantic-European North West Shelf- Wave Physics Reanalysis" wave model, AENWS, (Copernicus, 2025) and across simulated scenarios: the baseline scenario, the Current (2021) scenario, and the Future (2030) scenario. The locations where results were extracted are illustrated in Figure 8.5 and detailed in Appendix G.

To evaluate the significance of changes in wave conditions induced by the presence of wind farms in the Current and Future wind farm scenarios, the calculated annual and monthly variations were compared with the corresponding means and standard deviations derived from AENWS wave dataset. This allows the influence of offshore wind farms on the wave climate in Danish waters to be compared to natural variations in the wave climate on annual and monthly basis.

Statistical analysis was performed on wave data covering the period from 1988 to 2024. For each year, the mean annual and mean monthly significant wave height (H_{m0}) and mean wave directions (MWD) were calculated. An interannual evaluation was then carried out to obtain the overall annual mean, the monthly mean values, and the corresponding standard deviations. Thus, the standard deviation indicates how much the mean H_{m0} and the MWD varies from year to year, or for the same month across different years. The annual and monthly calculated statistics for the wave data are reported in Table 8.1 to Table 8.4, in the first and second columns for MWD and fifth and sixth columns for mean H_{m0} .

The third and fourth as well as the seventh and eighth columns of Table 8.1 to Table 8.4 show the difference between the annual or monthly MWD and mean H_{m0} , respectively, calculated from the modelled impact for the Current and Future windfarm scenarios compared to the Baseline Scenario.

Although all 16 points were analysed, this section presents the results for four of them (2, 8, 12 and 13), shown in Figure 8.5, while the remaining results, along with detailed information for each location (Appendix G), are provided in Appendix G. The results for the four selected locations are reported in Table 8.1 to Table 8.4.




Figure 8.5: : Locations of extraction of significant wave height and mean wave direction.

Based on the results for Point 2, located west of Jutland, presented in Table 8.1, the following key observations have been identified:

- **Wave Direction Statistics:** The mean annual and mean monthly wave directions vary between 265°N and 292°N. This range is consistent with the location of the measurement point and indicates a general stability of predominant wave directions throughout the year. The lowest standard deviation for MWD corresponds to the annual mean, which shows that the variability of MWD from year to year is smaller than the variability observed for individual months across different years.

- **Impact of Wind Farm Scenarios on Direction:** Changes in wave direction due to wind farm scenarios are minimal. The Current Scenario results in an annual shift of 0.2°, while the Future Scenario shows a change of 0.4°. Monthly differences generally stay within ±1.7°, except for April under the Future Scenario (9.0°), which, although notable, remains within the bounds of natural variability.
- **Wave Height Statistics:** The annual AENWS wave statistics indicate relatively stable conditions, with an annual mean H_{m0} of 1.46 m and a low standard deviation of 0.11 m, suggesting moderate wave energy at this site. Monthly averages vary from 1.09 m in May to 1.95 m in January, and standard deviations range from 0.20 m to 0.50 m, highlighting increased variability during the winter season.
- **Impact of Wind Farm Scenarios on Wave Height:** The impacts of both wind farm scenarios on significant wave height is limited. The Current scenario shows an annual change of approximately -0.01 m, with monthly variations generally between -0.03 m and 0.00 m. The Future scenario indicates a slightly larger reduction in significant wave height, with an annual change of -0.05 m and monthly differences up to -0.06 m (notably in May, July, August, and September). Overall, these results suggest only a slight decrease in wave energy under both scenarios, with changes remaining negligible compared to natural variability.
- **Overall Interpretation:** Both wave direction and wave height impacts from wind farms are negligible compared to natural seasonal variability in the location of Point 2. While the future scenario shows slightly larger reductions in significant wave height and minor directional shifts, these changes do not exceed thresholds that would significantly alter the local wave climate.

Table 8.1: **Point 2** - Annual and monthly statistical results based on AENWS data, showing mean and standard deviation of mean wave directions (first two columns), the differences between calculated annual and monthly mean wave directions across the defined scenarios (third and fourth column), mean and standard deviation of significant wave height (fifth and sixth column), and the differences between calculated annual and monthly significant wave height across the defined scenarios (seventh and last columns).

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean H_{m0}	St. Dev.	Impact on H_{m0}	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	276.6°	5.8°	0.2°	0.4°	1.46	0.11	-0.01	-0.05
Jan	269.8°	26.8°	0.0°	-0.1°	1.95	0.50	0.00	-0.03
Feb	272.3°	21.2°	0.4°	1.5°	1.83	0.50	-0.01	-0.05
Mar	276.2°	26.7°	0.2°	0.4°	1.54	0.38	-0.01	-0.04
Apr	292.1°	20.1°	-1.0°	9.0°	1.18	0.27	-0.01	-0.03
May	292.1°	19.3°	0.0°	-0.2°	1.09	0.21	-0.01	-0.06
Jun	292.4°	14.5°	0.4°	1.7°	1.16	0.20	-0.02	-0.05
Jul	287.3°	14.2°	0.4°	0.5°	1.13	0.23	-0.01	-0.06
Aug	279.3°	16.9°	0.9°	1.5°	1.17	0.22	-0.02	-0.06
Sep	279.1°	24.4°	0.4°	0.2°	1.39	0.32	0.00	-0.06
Oct	265.5°	22.7°	0.7°	0.6°	1.61	0.27	-0.01	-0.05
Nov	268.1°	30.0°	-9.1°	-6.9°	1.71	0.37	-0.02	-0.05
Dec	270.1°	24.1°	0.0°	0.3°	1.83	0.45	-0.01	-0.04

Similar patterns and conclusions were identified across all other selected locations, with expected variations reflecting differences in local wave climates. As shown in Table 8.2, the specific observations for Point 8, located in Kattegat near the east coast of northern Jutland, are as follows:

- **Wave Direction Statistics:** The annual and monthly mean wave directions span from 104°N to 208°N, which aligns with the geographical position of the measurement point. Unlike Point 2, this wide range indicates a high variability in dominant wave directions throughout the year. The lowest standard deviation for MWD corresponds to the annual mean, confirming that interannual variability is smaller than the variability observed across individual months. Consistent with the broader range of mean directions, Point 8 exhibits generally higher annual and monthly standard deviations compared to Point 2, varying from 25° to nearly 90°.
- **Impact of Wind Farm Scenarios on Direction:** When comparing natural variability to the modeled wind farm impact on MWD, the impact remains below 5° in both scenarios, which is well within the natural directional variability simulated by AENWS.
- **Wave Height Statistics:** Due to its location, Point 8 reflects a milder wave climate than Point 2, though similar trends are observed. Annual statistics show relatively stable conditions, with a mean significant wave height of 0.55 m and a very low standard deviation (0.02 m). Monthly means range from 0.37 m to 0.70 m, with slightly higher values during winter, and monthly standard deviations between 0.06 m and 0.13 m. While monthly variability exceeds annual variability, the overall magnitude of wave heights is lower, resulting in smaller differences and reduced variability.
- **Impact of Wind Farm Scenarios on Wave Height:** The Current scenario shows almost no change (annual difference 0.00 m, monthly differences mostly -0.01 m to 0.00 m). Depending on the month considered, the Future scenario can indicate a slightly larger reduction that in no case exceeds -0.01 m. Overall, these results suggest only a slight decrease in wave energy under both scenarios for the position of this point. Despite the smaller gap between natural variability and wind farm-induced changes compared to Point 2, the differences remain negligible relative to natural variability.
- **Overall Interpretation:** Both wave direction and wave height impacts from wind farms are negligible compared to natural seasonal variability at Point 8.

Table 8.2: **Point 8** - Annual and monthly statistical results based on AENWS data, showing mean and standard deviation of mean wave directions (first two columns), the differences between calculated annual and monthly mean wave directions across the defined scenarios (third and fourth column), mean and standard deviation of significant wave height (fifth and sixth column), and the differences between calculated annual and monthly significant wave height across the defined scenarios (seventh and last columns).

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	167.9°	25.0°	-0.1°	-0.2°	0.55	0.02	0.00	-0.01
Jan	171.0°	54.8°	0.2°	-1.4°	0.70	0.13	0.00	-0.01
Feb	158.2°	79.2°	-0.3°	-0.9°	0.68	0.12	0.00	-0.01
Mar	158.1°	55.5°	-0.6°	-0.3°	0.58	0.09	0.00	0.00
Apr	104.3°	78.9°	-0.9°	-1.2°	0.48	0.07	0.00	-0.01
May	146.9°	87.9°	0.5°	2.4°	0.42	0.06	0.00	0.00
Jun	201.3°	69.7°	0.6°	-0.2°	0.39	0.07	0.00	0.00
Jul	208.3°	56.0°	3.2°	4.4°	0.37	0.07	0.00	-0.01
Aug	188.2°	62.0°	-0.1°	0.6°	0.42	0.08	-0.01	-0.01
Sep	166.7°	63.8°	2.3°	3.3°	0.55	0.10	0.00	-0.01
Oct	161.9°	43.1°	0.8°	-0.2°	0.66	0.11	0.00	-0.01
Nov	151.5°	64.7°	-0.4°	-0.7°	0.70	0.10	-0.01	-0.01
Dec	168.4°	58.4°	0.8°	0.9°	0.67	0.09	-0.01	-0.01

The following observations were drawn from Table 8.3 for Point 12, located in southern Kattegat north of Zealand:

- **Wave Direction Statistics:** The annual mean wave direction is 292°N, with a very low annual standard deviation (9.3°), indicating stable long-term directional conditions. Monthly mean directions vary significantly, from 277°N (January) to 326°N (April), and monthly standard deviations range widely (13.4° to 62.0°), confirming strong seasonal variability and directional spread.
- **Impact of Wind Farm Scenarios on Direction:** Differences between the wind farm scenarios and the baseline scenarios are minimal, generally within $\pm 2.5^\circ$, which is well below the natural monthly variability (up to 62°). The largest deviations occur in July (-2.5°) and October (-2.4°) under the Future scenario, but these remain negligible compared to the natural variability.
- **Wave Height Statistics:** Annual mean significant wave height is 0.65 m, with a low annual standard deviation (0.05 m), indicating mild wave conditions at this location. Monthly means range from 0.50 m (May) to 0.83 m (January), with standard deviations between 0.11 m and 0.23 m, showing higher variability during winter months.
- **Impact of Wind Farm Scenarios on Wave Height:** The Current scenario shows almost no change (annual difference 0.00 m, monthly differences mostly -0.01 m to 0.00 m). The Future scenario indicates slightly larger reductions in significant wave height, up to -0.03 m in several months, but these remain small relative to natural variability.
- **Overall Interpretation:** Both wave direction and wave height impacts from wind farms are negligible compared to natural seasonal variability at Point 12. While the future scenario shows slightly larger reductions in significant wave height and minor directional shifts, these changes do not exceed thresholds that would significantly alter the local wave climate.

Table 8.3: **Point 12** - Annual and monthly statistical results based on AENWS data, showing mean and standard deviation of mean wave directions (first two columns), the differences between calculated annual and monthly mean wave directions across the defined scenarios (third and fourth column), mean and standard deviation of significant wave height (fifth and sixth column), and the differences between calculated annual and monthly significant wave height across the defined scenarios (seventh and last columns).

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	292.4°	9.3°	-0.3°	-1.5°	0.65	0.05	0.00	-0.02
Jan	277.5°	42.0°	0.1°	-0.1°	0.83	0.23	0.00	-0.03
Feb	286.9°	47.9°	-0.3°	-1.0°	0.80	0.21	0.00	-0.03
Mar	291.0°	38.7°	-0.4°	-1.8°	0.66	0.18	-0.01	-0.03
Apr	325.7°	48.0°	1.0°	1.6°	0.54	0.14	-0.01	-0.01
May	318.2°	30.3°	-0.3°	-1.4°	0.50	0.11	0.00	-0.03
Jun	302.9°	13.4°	0.9°	-1.3°	0.54	0.13	-0.01	-0.03
Jul	299.5°	16.2°	-0.9°	-2.5°	0.53	0.14	0.00	-0.03
Aug	294.2°	30.6°	0.2°	-0.7°	0.55	0.14	0.00	-0.02
Sep	296.9°	35.1°	-0.4°	-1.8°	0.64	0.16	0.00	-0.03
Oct	281.1°	33.4°	-0.7°	-2.4°	0.72	0.15	0.00	-0.01
Nov	288.5°	62.0°	-1.2°	0.1°	0.74	0.16	0.00	-0.01
Dec	279.4°	40.4°	0.4°	-0.4°	0.80	0.19	0.01	-0.01

The observations for Point 13, located south of Møn in the Baltic Sea, shown in Table 8.4, are as follows:

- **Wave Direction Statistics:** The annual average wave direction is 188°N, accompanied by a standard deviation of 29.5°, indicating a high variability in dominant wave directions throughout the year. Monthly averages span from 112.7°N in April to 214.2°N in July, while monthly standard deviations range between 45.5° (January) and 74.6° (May), suggesting that Point 13 generally experiences a high variability in dominant wave directions throughout the year.
- **Impact of Wind Farm Scenarios on Direction:** Changes in wave direction due to wind farm scenarios are limited. The Current Scenario results in an annual shift of 0.7°, while the Future Scenario shows a change of 3.3°. Monthly differences generally stay within $\pm 5^\circ$, which is well within the bounds of the natural variability at this location.
- **Wave Height Statistics:** The annual mean significant wave height is 0.68 m, with a standard deviation of 0.04 m, suggesting mild wave conditions at this location. Monthly averages vary from 0.49 m in June to 0.87 m in January, and standard deviations range from 0.09 m to 0.18 m, with higher values during winter.
- **Impact of Wind Farm Scenarios on Wave Height:** the Current scenario shows negligible impact, with monthly differences typically between -0.03 m and 0.00 m. The Future scenario presents greater changes in wave height, reaching up to -0.10 m in April, and thus exceeding the standard deviation for this month. Generally, for Point 13, however, the impacts from the wind farm scenarios are limited when compared to natural variation.
- **Overall Interpretation:** The impact of wind farms on both wave direction and wave height is limited in comparison to natural variability. Although the Future scenario introduces a reduction in mean significant wave height in April exceeding the natural variation, these changes are larger than for the other points but not substantial enough to significantly affect the local wave climate.

Table 8.4: **Point 13** - Annual and monthly statistical results based on AENWS data, showing mean and standard deviation of mean wave directions (first two columns), the differences between calculated annual and monthly mean wave directions across the defined scenarios (third and fourth column), mean and standard deviation of significant wave height (fifth and sixth column), and the differences between calculated annual and monthly significant wave height across the defined scenarios (seventh and last columns).

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	188.2°	29.5°	0.7°	3.3°	0.68	0.04	-0.01	-0.04
Jan	201.7°	45.5°	-0.6°	-2.2°	0.87	0.18	-0.01	-0.02
Feb	186.9°	68.9°	-0.0°	-0.2°	0.85	0.16	-0.01	-0.03
Mar	173.2°	59.6°	-0.1°	-0.3°	0.73	0.14	0.00	-0.01
Apr	112.7°	73.5°	0.5°	0.7°	0.62	0.09	-0.03	-0.10
May	116.9°	74.6°	0.3°	5.3°	0.55	0.09	-0.01	-0.04
Jun	206.9°	72.4°	0.8°	4.4°	0.49	0.12	-0.02	-0.05
Jul	214.2°	58.6°	1.4°	2.2°	0.50	0.09	-0.01	-0.03
Aug	202.0°	55.7°	1.5°	3.6°	0.52	0.09	-0.02	-0.02
Sep	187.4°	64.3°	0.6°	0.7°	0.67	0.11	-0.01	-0.03
Oct	189.2°	49.3°	-0.0°	0.9°	0.79	0.15	-0.01	-0.03
Nov	187.8°	56.8°	1.0°	2.7°	0.80	0.10	0.01	-0.04
Dec	203.0°	47.4°	0.1°	0.0°	0.85	0.12	0.00	-0.02

8.3. Conclusion on wave impact

A comparative analysis of conditions without wind farms versus those with wind farms in 2021 and projected for 2030 indicates that wind farms lead to a reduction in significant wave height on both annual and monthly scales as described in section 8.1.

The analysis comparing conditions with and without wind farms is based on data from 2019. To account for natural annual and monthly variations, wave data—including significant wave height and mean wave direction—were examined using information from Copernicus (2025) for the years 1988 to 2024. This analysis covers 16 locations, ranging from the west coast of Jutland, through the inner Danish waters, to the Baltic Sea.

The study, described in section 8.2, indicates that natural variation exceeds the variation attributed to both the 2021 wind farm scenario and projections for 2030 along the west coast of Jutland and in the Kattegat area by approximately an order of magnitude. In contrast, the reduction observed in the Baltic Sea due to wind farms is greater and approaches the annual and monthly standard deviation, suggesting that the Baltic Sea is more sensitive towards the establishment of wind farms with relation to the wave climate, compared to the Kattegat and the North Sea.

9. Impact on sediment transport

9.1. Introduction

The establishment of wind farms in Danish and neighbouring waters results in changes to wave patterns. Such modifications have the potential to affect longshore sediment transport rates, which may consequently impact the ongoing evolution of the shoreline.

To investigate how these changes in wave climate affect the coastal longshore sediment transport, the sediment transport capacity has been calculated along three coastal stretches for the Baseline scenario, as well as for the Current (2021) and Future (2030) wind farm developments in the North Sea, Kattegat, and the Baltic Sea. The following stretches are included in the analysis:

- Løkken to Hirtshals.
- Ferring to Hvide Sande.
- Hjelm Bugt.

The locations of the three coastal stretches are presented in Figure 9.1.




Figure 9.1: Location of the three stretches investigated for impact on longshore sediment transport.

The sediment transport capacity is the upper theoretical limit for the amount of sediment that waves and currents can transport under specific hydrodynamic conditions. It reflects the potential for sediment movement, based on the assumption that there is an unlimited supply of sand available.

On the other hand, the actual sediment transport is the real amount of sediment that is moved by waves and currents in a given place and time. This depends not only on the transport capacity but also on the availability of sediment, grain size, distribution, and other environmental factors such as vegetation or man-made structures.

Changes in the net longshore sediment transport gradients along the coast may lead to sediment deposition if the gradient is negative, or erosion if the gradient is positive.

9.1.1. Løkken to Hirtshals

As illustrated in Figure 9.2, the longshore sediment transport between Løkken and Hirtshals is predominantly directed northward (Kystdirektoratet, 2022). Previous studies have estimated the magnitude of the net longshore sediment transport to be in the range of 400,000 – 900,000 m³/yr (Kystdirektoratet, 2001; DHI, 2000).




Figure 9.2: Coastal stretch from Løkken to Hirtshals. Blue arrow indicates the net sediment transport direction (Kystdirektoratet, 2022). Placements of the eight profiles are indicated by red lines perpendicular to the coast. Orthophoto: 2024.

9.1.2. Ferring to Hvide Sande

As illustrated in Figure 9.3, the longshore sediment transport between Ferring and Hvide Sande is predominantly directed southward (Kystdirektoratet, 2022). North of Ferring, the transport is directed northward, resulting in a point near Ferring where the net longshore sediment transport is approximately $0 \text{ m}^3/\text{year}$. From this point, sediment diverges both northward and southward. In a previous study, the magnitude of the net longshore sediment transport at Hvide Sande was estimated to be approximately $400,000 \text{ m}^3/\text{year}$ (DHI, 2000).




Figure 9.3: Coastal stretch from Ferring to Hvide Sande. Blue arrow indicates the net sediment transport direction (Kystdirektoratet, 2022). Placements of the ten profiles are indicated by red lines perpendicular to the coast. Orthophoto: 2024.

9.1.3. Hjelm Bugt

As illustrated in Figure 9.4, the longshore sediment transport in Hjelm Bugt is predominantly directed westward (Kystdirektoratet, 2022). However, an orthophoto from spring 2019 suggests that the sediment transport was eastward at that time. Figure 9.4 presents the orthophoto from the northern part of Hjelm Bugt, showing erosion on the east side of the groynes and sediment accumulation on the west side. This pattern indicates that sediment was transported from west to east, becoming trapped by the groynes and resulting in sediment build-up and a wider beach on their western side.




Figure 9.4: Erosion and accumulation in groyne field in the northern part of Hjelm Bugt. Blue arrow indicates the net sediment transport direction (Kystdirektoratet, 2022). Placements of the eight profiles are indicated by grey lines perpendicular to the coast. Orthophoto: 2019.

9.2. Data

The sediment transport capacity is calculated in cross-shore profiles along the selected coastal stretches. These profiles are a crucial input to the LITDRIFT model (DHI, 2024).

Along the west coast of Jutland, the Danish Coastal Authority (Kystdirektoratet) conducts regular measurements of coastal profiles at predefined locations spaced approximately 1 km apart. These profiles extend from the top of the dune to about 4 km offshore.

At Hjelm Bugt, no measured cross-shore profiles are available. Instead, elevation data for the dry part of the profile is extracted from the Danish elevation model (SDFI, 2025), while depth data is obtained from the Danish depth model (Geodatastyrelsen, 2024). These datasets are combined to form complete profiles, with the gap between land and sea data filled using linear interpolation.

9.2.1. Profile orientation

The orientation of each cross-shore profile is determined as the line perpendicular to the coastline at the profile location. To achieve this, the coastline is coarsely digitized to capture the average orientation over the distance represented by each profile, thereby excluding minor undulations. The orientation is then assigned to each profile based on this smoothed coastline.

9.2.2. Sediment grain size

Sediment characteristics are described by the mean grain size (D_{50}). For the Løkken to Hirtshals and Ferring to Hvide Sande stretches, the mean grain size is based on sediment samples from the 2 m depth contour, as presented in DHI (2000). For Hjelm Bugt, the mean sediment grain size is assumed to be 0.2 mm. Table 9.1 shows the D_{50} values for all profiles.

Table 9.1: Mean sediment grain size (D_{50}) used in the model. The grain sizes are provided for each profile at the three locations (see Figure 9.2 to Figure 9.4 for profile positions).

Location	Profile	D_{50} [mm]
Løkken to Hirtshals	All	0.17
Ferring to Hvide Sande	1	0.31
Ferring to Hvide Sande	2	0.30
Ferring to Hvide Sande	3	0.29
Ferring to Hvide Sande	4, 5	0.27
Ferring to Hvide Sande	6	0.26
Ferring to Hvide Sande	7, 8	0.25
Ferring to Hvide Sande	9, 10	0.24
Hjelm Bugt	All	0.20

9.2.3. Waves

Wave parameters (significant wave height (H_{m0}), peak wave period (T_p), and mean wave direction (MWD)) are extracted from the MIKE SW model at points located approximately 10 km offshore along the coast. This ensures that the wave time series are representative of the conditions at the corresponding cross-shore profiles.

9.3. Methods

The DHI LITDRIFT model is used to calculate the longshore sediment transport capacity. LITDRIFT simulates wave transformation across the cross-shore profile—from the offshore point to the coastline—and the resulting wave-driven currents. Based on the computed hydrodynamics, the model estimates the longshore sediment transport along each profile.

Required input for LITDRIFT includes:

- Bathymetry in cross-shore profiles
- Orientation of cross-shore profiles
- Time-series of:
 - Root-mean-square wave height
 - Peak wave period
 - Mean wave direction
- Sediment median grain size (D_{50})

9.3.1. Wave height conversion

The wave heights extracted from the wave model are significant wave heights (H_{m0}). Since the LITDRIFT model requires root-mean-square wave heights (H_{rms}), the following conversion is applied:

$$H_{rms} = \frac{H_{m0}}{\sqrt{2}}$$

9.3.2. Limitations

While the LITDRIFT model provides valuable insights into longshore sediment transport dynamics, several limitations should be noted:

- In this analysis, the model simulations are based on a single year of wave data. This makes the results sensitive to the waves occurring that year which may not be fully representative for the wave climate at the analyzed locations. As a result, the long-shore sediment transport may be different that year compared to other years.
- The model assumes a constant sediment grain size (D_{50}) across each cross-shore profile. In reality, sediment characteristics often vary across the profile due to natural sorting processes, wave action, and sediment sources. This simplification may affect the accuracy of transport estimates.
- The LITDRIFT model focuses exclusively on longshore sediment transport and does not account for cross-shore transport processes such as onshore-offshore sediment exchange, bar migration, or storm-induced erosion and recovery. These processes can significantly influence beach morphology and sediment budgets.
- The model assumes an open natural coastline without man-made structures. However, all three coastal stretches include features such as groynes and breakwaters. These structures can locally modify wave conditions and influence sediment transport patterns that are not captured by the model.
- The model does not include morphological updating, meaning that changes in the seabed or shoreline due to sediment transport are not fed back into the hydrodynamic calculations. This limits the model's ability to simulate evolving coastal systems over time.

Despite these limitations, the model output can be used for evaluating the overall impact on sediment transport from the two wind farm scenarios compared to the baseline situation as the comparison is relative. Unlike the documented effects on the significant wave height, longshore sediment transport reflects cumulative influences based on hourly data, taking into account not only significant wave height, but also changes in wave period and mean wave direction. For example, when considering sediment transport from Ferring to Hvide Sande, the movement can occur either northward or southward. By analysing the sediment transport, it is possible to determine if one direction is more impacted than the other, which may result in either deposition or erosion.

9.4. Results

Figure 9.5, Figure 9.7 and Figure 9.9 present the simulated longshore sediment transport capacity for the coastal stretches from Løkken to Hirtshals, Ferring to Hvide Sande, and Hjelm Bugt, respectively. Each figure shows the simulated longshore sediment transport capacity in both directions along the shore for the baseline scenario. Additionally, the figures illustrate the changes in transport capacity in both directions and net transport for the current and future scenarios relative to the baseline.

Figure 9.6, Figure 9.8 and Figure 9.10 present the simulated net longshore sediment transport capacity for the three scenarios together with annotation of whether the gradient in transport capacity leads to erosion or accumulation in the profile. Additionally, for the current and future scenarios, it is indicated whether more or less sediment is eroded or accumulated in the profile compared to the baseline.

At **Løkken to Hirtshals**, the simulated longshore sediment transport capacity highlights the following observations:

- **Baseline:** The sediment transport is directed northward along the entire coastal stretch, with an overall increase towards Hirtshals, as expected.
- **Current scenario (2021):** Compared to the baseline, only minor changes are observed in the transport capacity in both directions and net transport capacity across all profiles ($< \pm 0.7\%$ and $< \pm 0.4\%$, respectively). Consequently, the changes in erosion and accumulation due to the current wind farm scenario are minimal.
- **Future scenario (2030):** Compared to the baseline, all profiles exhibit an increase in northward transport. Profiles 4 and 8 show a decrease in southward transport, while the remaining profiles indicate an increase. However, the changes remain small ($< \pm 1.3\%$). The relatively larger increase in northward transport capacity results in an overall increase in net transport capacity across all profiles, particularly between profiles 4 and 8. Due to differences in the increase of net transport capacity, erosion is intensified in profiles 4, 7, and 8, while accumulation increases in profiles 5 and 6.




Figure 9.5: Simulated longshore sediment transport capacity in both directions for Løkken to Hirtshals for the baseline scenario (a). Difference in sediment transport capacity in both directions between the baseline and the current scenario (b),

between the baseline and the future scenario (c) and difference in net sediment transport capacity for both scenarios (d). Note that in panel (d), positive values indicate northward transport, and negative values indicate southward transport.




Figure 9.6: Direction and magnitude of simulated net sediment transport capacity for the Baseline, Current (2021) and Future (2030) scenario are indicated by the arrows. Based on the gradient in transport capacity, it is shown whether the profile experience erosion (E) or accumulation (A). In the current and future scenarios, it is noted whether the erosion or accumulation has increased (+) or decreased (-), and whether more (blue) or less (red) sediment is being transported to the profile.

At **Ferring to Hvide Sande**, the simulated longshore sediment transport capacity highlights the following observations:

- **Baseline:** The net sediment transport is directed southward along the entire coastal stretch, as expected. The erosion and accumulation pattern varies along the coast.
- **Current scenario (2021):** Compared to the baseline, only minor changes are observed in southward transport capacity across all profiles (<0.8%). Profiles 1–3 exhibit a slight increase (<0.5%) in northward transport capacity, while profiles 4–10 show a decrease, reaching up to -2.2% in profile 10.
- **Future scenario (2030):** All profiles show a decrease in both northward and southward transport capacity. The decrease in northward transport is relatively uniform (6%–10%), while the decrease in southward transport ranges from 1%–5%. Despite the decrease in transport capacity in both directions, the net transport capacity increases in all profiles except profile 10. This is due to a larger reduction in

northward transport compared to southward transport. The most significant changes in net transport capacity occur in profiles 1–4, where the decrease in southward transport is relatively small compared to the decrease in northward transport. These changes in net transport capacity result in increased erosion in profiles 1 and 2, due to a rising transport gradient between them. From profile 3 onwards, the remaining profiles—except profile 6—experience reduced erosion or increased accumulation, as the transport gradient decreases.





Figure 9.7: Simulated longshore sediment transport capacity in both directions for Ferring to Hvide Sande for the baseline scenario (a). Difference in sediment transport capacity in both directions between the baseline and the current scenario (b), between the baseline and the future scenario (c) and difference in net sediment transport capacity for both scenarios (d). Note that in panel (d), positive values indicate northward transport, and negative values indicate southward transport.



Impact on net sediment transport capacity




Figure 9.8: Direction and magnitude of simulated net sediment transport capacity for the baseline, Current (2021) and Future (2030) scenario are indicated by the arrows. Based on the gradient in transport capacity, it is shown whether the profile experience erosion (E) or accumulation (A). In the Current (2021) and Future (2030) scenarios, it is noted whether the erosion or accumulation has increased (+) or decreased (-), and whether more (blue) or less (red) sediment is being transported to the profile.

At **Hjelm Bugt**, the simulated longshore sediment transport capacity highlights the following observations:

- **Baseline:** In the northern part of the bay (profiles 1–5), sediment transport is directed eastward, while in the rest of the bay, it is directed westward.
- **Current (2021) scenario:** Compared to the baseline, the eastward transport capacity in the current scenario shows only minor changes across all profiles (<1.4%).
- **Future (2030) scenario:** All profiles show a decrease in eastward transport, although this is minor compared to the decrease in westward transport.
- **Current (2021) and Future (2030) scenarios:** Westward transport capacity has decreased substantially in all profiles except profiles 1 and 2, primarily due to reduced wave energy from the west. This results in an increase in net transport capacity in profiles 3–5, and a decrease in profiles 6–8, where the net sediment transport is westward.
- **Baseline:** The net sediment transport capacity increases away from the point of divergence, leading to erosion in the middle of the bay (profiles 3–7) and accumulation in profiles 1, 2, and 8.
- **Current (2021) and Future (2030) scenarios:** No clear trend is observed in the changes to erosion and accumulation patterns.

- It is noteworthy that sediment transport is typically westward throughout the bay (see section 9.1.3). Therefore, a reduction in westward transport may lower both the magnitude of net sediment transport and the associated gradients.





Figure 9.9: Simulated longshore sediment transport capacity in both directions for Hjelm Bugt for the baseline scenario (a). Difference in sediment transport capacity in both directions between the baseline and the current scenario (b), between the baseline and the future scenario (c) and difference in net sediment transport capacity for both scenarios (d). Note that in panel d positive values indicate a net northward transport and negative values indicate a net southward transport. Note that in panel (d), positive values indicate westward transport, and negative values indicate eastward transport.



Impact on net sediment transport capacity

Figure 9.10: Direction and magnitude of simulated net sediment transport capacity for the Baseline, Current (2021) and Future (2030) scenario are indicated by the arrows. Based on the gradient in transport capacity, it is shown whether the profile experience erosion (E) or accumulation (A). In the current and future scenarios, it is noted whether the erosion or accumulation has increased (+) or decreased (-), and whether more (blue) or less (red) sediment is being transported to the profile.

9.5. Uncertainty analysis

To examine the model's limitations, profile orientations along the Ferring to Hvide Sande coast were adjusted by $\pm 2^\circ$ from baseline. Longshore sediment transport capacity was then calculated using these altered orientations.

A change in profile orientation of 2° is equivalent to altering the wave climate by 2° . The comparison between the baseline and sensitivity model runs shows that the magnitude of transport capacity in both northward and southward directions vary by 6,000 to 43,000 m³/year, corresponding to a change of approximately 1% to 8%.

In section 8.2 at a location offshore Hvide Sande, the annual MWD standard deviation is found to be above 5° and the monthly between 10° to 30° , indicating that the longshore sediment transport on annual basis experience change on a larger scale than may be expected given by the wind farms.

9.6. Conclusion on sediment transport impact

Overall, the modelled impact on sediment transport is larger for the Future (2030) scenario compared to the Current (2021) scenario, and the impact at Hjelm Bugt in the Baltic Sea is larger compared to the impact along the west coast of Jutland bordering the North Sea, which may indicate that the wave climate in the Baltic Sea is more sensitive to impacts from future offshore wind farm development.

It is important to note, however, that the annual variation in the wave climate indicates a larger change compared to the impact introduced by wind farms.

Also, although the results show that offshore wind farms can influence coastal processes, the current analysis is insufficient to draw definitive conclusions about long-term coastline development or morphological changes.

10. References

- Arneborg, L., et al. 2024.** *Hydrographic effects in Swedish waters of future offshore wind power scenarios*. s.l. : REPORT OCEANOGRAPHY No 77, 2024, 2024.
- Copernicus. 2025.** Copernicus Marine Data Store. [Online] 01 07 2025.
<https://data.marine.copernicus.eu/products?facets=specificVariables%7EWave+significant+height--areas%7EAtlantic%3A+North>.
- Daewel, U, et al. 2022.** Offshore wind farms are projected to impact primary production and bottom water deoxygenation in the North Sea. *Communications Earth & Environment*. 2022.
- DHI. 2024.** *Littoral Processes FM - Scientific Documentation*. 2024.
- . **2021.** *MIKE 21 Spectral Wave Module. Scientific Documentation*. Hørsholm : s.n., 2021.
- . **2000.** *Sedimentbudget Vestkysten*. 2000.
- EMODnet. 2023.** Bathymetry. <portal.emodnet-bathymetry.eu/#>. [Online] 15 03 2023. <portal.emodnet-bathymetry.eu/#>.
- Geodatastyrelsen. 2024.** *Den danske dybdemodel*. 2024.
- German Federal Maritime and Hydrographic Agency (BSH). 2021.** *Report on the oceanographic conditions at site O-1.3*. 2021.
- Hahmann, Andrea N, et al. 2025a.** *Environmental Mapping and Screening of the Offshore Wind Potential in Denmark*. s.l. : DTU Public report number DTU E-0255 2025, 2025a.
- Hahmann, Andrea N., et al. 2025b.** *Comparison of Old and New Simulations*. 2025b.
- Howard, Katherine, et al. 2009.** *The Impact Of Atmospheric Model Resolution On A Coupled Wind/Wave Forecast System*. 2009.
- Kystdirektoratet. 2022.** *Kystdirektoratets Kystatlas*. [Online] 29 03 2022. [Cited: 29 03 2022.]
<https://kms.maps.arcgis.com/apps/webappviewer/index.html?id=8669133b3f4842b7a9a19fb24b08ffd5>.
- . **2001.** *Sedimentbudget Vestkysten*. 2001.
- Maar, M, et al. 2025.** *Spatial impacts of offshore wind farms on hydrodynamics and biogeochemical environment*. s.l. : Scientific Report from DCE – Danish Centre for Environment and Energy No. 658, 2025.
- SDFI. 2025.** *Danmarks Højdemodel*. 2025.
- van Duren, LA, et al. 2021.** *Ecosystem effects of large upscaling of offshore wind on the North Sea - Synthesis report*. s.l. : Rijkswaterstaat Water, Verkeer en Leefomgeving, 2021.

Appendix A: Available measurements of waves

Location and list of stations with measurements covering the year 2019.





Figure 10.1: Location of the retrieved wave measurements for 2019.


Station ID	Longitude [deg]	Latitude [deg]	Time	Coverage (%)
BO_TS_MO_BrofjordenWR	11.225	58.251	UTM0	100
BO_TS_MO_VaderoarnaWR	10.953	58.496	UTM0	86
NO_TS_MO_1000044	10.267	54.500	UTM0	97
NO_TS_MO_6200041	0.475	53.058	UTM0	100
NO_TS_MO_6200042	1.100	53.058	UTM0	100
NO_TS_MO_6200043	1.784	52.312	UTM0	100
NO_TS_MO_6200044	1.579	51.570	UTM0	93
NO_TS_MO_6200045	-2.504	56.188	UTM0	100
NO_TS_MO_6200170	2.000	51.240	UTM0	90
NO_TS_MO_6200288	0.754	50.746	UTM0	100
NO_TS_MO_6200289	1.054	53.532	UTM0	100
NO_TS_MO_6200305	0.000	50.400	UTM0	100
NO_TS_MO_6201010	-1.129	50.650	UTM0	100
NO_TS_MO_6201011	-0.959	50.732	UTM0	100
NO_TS_MO_6201012	-0.839	50.723	UTM0	100
NO_TS_MO_6201015	0.417	50.783	UTM0	100
NO_TS_MO_6201019	-0.067	53.917	UTM0	100
NO_TS_MO_6201050	0.447	53.245	UTM0	100
NO_TS_MO_6201058	2.109	51.953	UTM0	100
NO_TS_MO_6201059	1.817	52.476	UTM0	100
NO_TS_MO_6201067	7.783	55.017	UTM0	85
NO_TS_MO_6201068	6.839	53.985	UTM0	80
NO_TS_MO_6202112	7.150	55.200	UTM0	44
NO_TS_MO_6202600	7.818	54.219	UTM0	93
NO_TS_MO_6202601	8.114	53.996	UTM0	97
NO_TS_MO_A122	3.817	55.418	UTM0	100
NO_TS_MO_Akkaert	2.819	51.416	UTM0	100
NO_TS_MO_AWG	5.950	53.500	UTM0	100
NO_TS_MO_Deurlø	3.242	51.504	UTM0	100
NO_TS_MO_DomburgerRassen	3.399	51.620	UTM0	100
NO_TS_MO_EurogeulDWE	3.000	51.948	UTM0	100
NO_TS_MO_EurogeulE13	3.737	52.010	UTM0	100
NO_TS_MO_Europlatform3	3.276	51.999	UTM0	100
NO_TS_MO_F3platform	4.727	54.854	UTM0	96
NO_TS_MO_IJmuidenMunitiestort2	4.058	52.550	UTM0	100
NO_TS_MO_J61	2.950	53.817	UTM0	100
NO_TS_MO_K13a3	3.220	53.218	UTM0	100
NO_TS_MO_K141	3.633	53.267	UTM0	96
NO_TS_MO_Kwintebank	2.706	51.349	UTM0	100
NO_TS_MO_L91	4.961	53.614	UTM0	100
NO_TS_MO_LichteilandGoeree2	3.670	51.926	UTM0	96
NO_TS_MO_MeetboeiWEO1	6.518	53.617	UTM0	96
NO_TS_MO_MeetboeiWEW1	6.368	53.620	UTM0	96
NO_TS_MO_SchiermonnikoogNoord	6.167	53.596	UTM0	96

Station ID	Longitude [deg]	Latitude [deg]	Time	Coverage (%)
NO_TS_MO_SchiermonnikoogWestgat	6.165	53.529	UTM0	100
NO_TS_MO_Schouwenbank2	3.306	51.745	UTM0	100
NO_TS_MO_StortemelkBoei	4.989	53.323	UTM0	100
NO_TS_MO_WaddenEierlandseGat	4.662	53.277	UTM0	100
BROFJORDEN WR BOJ	11.5734	57.6234	UTM2	100
Vinga BS	11.2229	58.2513	UTM2	44
Arkona Basin	13.859	54.884	UTM1	N/A
Valhall A	3.393	56.278	UTM1	100
Ula	2.846	57.408	UTM1	100
Granefeltet	2.487	59.165	UTM1	100
Gudrunfeltet	1.744	58.845	UTM1	100


Appendix B: Time-series of significant wave heights




Arkona_Basin_30T_Hs: TS_plot_Arkona_Basin_30T_Arkona_Basin_30T




Arkona_Basin_30T_Tp: TS_plot_Arkona_Basin_30T_Arkona_Basin_30T




BO_TS_MO_BrofjordenWR_Hs: TS_plot_BO_TS_MO_BrofjordenWR_BO_TS_MO_BrofjordenWR




BO_TS_MO_BrofjordenWR_Tp: TS_plot_BO_TS_MO_BrofjordenWR_BO_TS_MO_BrofjordenWR




BO_TS_MO_VadaroarnaWR_Hs: TS_plot_BO_TS_MO_VadaroarnaWR_BO_TS_MO_VadaroarnaWR




BO_TS_MO_VadaroarnaWR_Tp: TS_plot_BO_TS_MO_VadaroarnaWR_BO_TS_MO_VadaroarnaWR




BROFJORDEN WR BOJ_Hs: TS_plot_BROFJORDEN WR BOJ_BROFJORDEN WR BOJ




BROFJORDEN WR BOJ_Tp: TS_plot_BROFJORDEN WR BOJ_BROFJORDEN WR BOJ




Granefeltet_Hs: TS_plot_Granefeltet_Granefeltet




Gudrunfeltet_Hs: TS_plot_Gudrunfeltet_Gudrunfeltet




Heimdal_Hs: TS_plot_Heimdal_Heimdal




NO_TS_MO_1000044_30T_Hs: TS_plot_NO_TS_MO_1000044_30T_NO_TS_MO_1000044_30T




NO_TS_MO_1000044_30T_Tp: TS_plot_NO_TS_MO_1000044_30T_NO_TS_MO_1000044_30T




NO_TS_MO_6200041_Hs: TS_plot_NO_TS_MO_6200041_NO_TS_MO_6200041




NO_TS_MO_6200041_Tp: TS_plot_NO_TS_MO_6200041_NO_TS_MO_6200041




NO_TS_MO_6200042_Hs: TS_plot_NO_TS_MO_6200042_NO_TS_MO_6200042




NO_TS_MO_6200042_Tp: TS_plot_NO_TS_MO_6200042_NO_TS_MO_6200042




NO_TS_MO_6200043_Hs: TS_plot_NO_TS_MO_6200043_NO_TS_MO_6200043




NO_TS_MO_6200043_Tp: TS_plot_NO_TS_MO_6200043_NO_TS_MO_6200043




NO_TS_MO_6200044_Hs: TS_plot_NO_TS_MO_6200044_NO_TS_MO_6200044




NO_TS_MO_6200044_Tp: TS_plot_NO_TS_MO_6200044_NO_TS_MO_6200044




NO_TS_MO_6200045_Hs: TS_plot_NO_TS_MO_6200045_NO_TS_MO_6200045




NO_TS_MO_6200045_Tp: TS_plot_NO_TS_MO_6200045_NO_TS_MO_6200045




NO_TS_MO_6200170_Hs: TS_plot_NO_TS_MO_6200170_NO_TS_MO_6200170




NO_TS_MO_6200288_Hs: TS_plot_NO_TS_MO_6200288_NO_TS_MO_6200288




NO_TS_MO_6200288_Tp: TS_plot_NO_TS_MO_6200288_NO_TS_MO_6200288




NO_TS_MO_6200289_Hs: TS_plot_NO_TS_MO_6200289_NO_TS_MO_6200289




NO_TS_MO_6200289_Tp: TS_plot_NO_TS_MO_6200289_NO_TS_MO_6200289




NO_TS_MO_6200305_Hs: TS_plot_NO_TS_MO_6200305_NO_TS_MO_6200305




NO_TS_MO_6201010_Hs: TS_plot_NO_TS_MO_6201010_NO_TS_MO_6201010




NO_TS_MO_6201010_Tp: TS_plot_NO_TS_MO_6201010_NO_TS_MO_6201010




NO_TS_MO_6201011_Hs: TS_plot_NO_TS_MO_6201011_NO_TS_MO_6201011




NO_TS_MO_6201011_Tp: TS_plot_NO_TS_MO_6201011_NO_TS_MO_6201011




NO_TS_MO_6201012_Hs: TS_plot_NO_TS_MO_6201012_NO_TS_MO_6201012




NO_TS_MO_6201012_Tp: TS_plot_NO_TS_MO_6201012_NO_TS_MO_6201012




NO_TS_MO_6201014_Hs: TS_plot_NO_TS_MO_6201014_NO_TS_MO_6201014




NO_TS_MO_6201014_Tp: TS_plot_NO_TS_MO_6201014_NO_TS_MO_6201014




NO_TS_MO_6201015_Hs: TS_plot_NO_TS_MO_6201015_NO_TS_MO_6201015




NO_TS_MO_6201015_Tp: TS_plot_NO_TS_MO_6201015_NO_TS_MO_6201015




NO_TS_MO_6201019_Hs: TS_plot_NO_TS_MO_6201019_NO_TS_MO_6201019




NO_TS_MO_6201019_Tp: TS_plot_NO_TS_MO_6201019_NO_TS_MO_6201019




NO_TS_MO_6201050_Hs: TS_plot_NO_TS_MO_6201050_NO_TS_MO_6201050




NO_TS_MO_6201050_Tp: TS_plot_NO_TS_MO_6201050_NO_TS_MO_6201050




NO_TS_MO_6201058_Hs: TS_plot_NO_TS_MO_6201058_NO_TS_MO_6201058




NO_TS_MO_6201058_Tp: TS_plot_NO_TS_MO_6201058_NO_TS_MO_6201058




NO_TS_MO_6201059_Hs: TS_plot_NO_TS_MO_6201059_NO_TS_MO_6201059




NO_TS_MO_6201059_Tp: TS_plot_NO_TS_MO_6201059_NO_TS_MO_6201059




NO_TS_MO_6201067_Hs: TS_plot_NO_TS_MO_6201067_NO_TS_MO_6201067




NO_TS_MO_6201067_Tp: TS_plot_NO_TS_MO_6201067_NO_TS_MO_6201067




NO_TS_MO_6201068_Hs: TS_plot_NO_TS_MO_6201068_NO_TS_MO_6201068




NO_TS_MO_6201068_Tp: TS_plot_NO_TS_MO_6201068_NO_TS_MO_6201068




NO_TS_MO_6202112_Hs: TS_plot_NO_TS_MO_6202112_NO_TS_MO_6202112




NO_TS_MO_6202112_Tp: TS_plot_NO_TS_MO_6202112_NO_TS_MO_6202112




NO_TS_MO_6202600_Hs: TS_plot_NO_TS_MO_6202600_NO_TS_MO_6202600




NO_TS_MO_6202600_Tp: TS_plot_NO_TS_MO_6202600_NO_TS_MO_6202600




NO_TS_MO_6202601_Hs: TS_plot_NO_TS_MO_6202601_NO_TS_MO_6202601




NO_TS_MO_6202601_Tp: TS_plot_NO_TS_MO_6202601_NO_TS_MO_6202601




NO_TS_MO_A122_Hs: TS_plot_NO_TS_MO_A122_NO_TS_MO_A122




NO_TS_MO_A122_Tp: TS_plot_NO_TS_MO_A122_NO_TS_MO_A122




NO_TS_MO_Akkaert_Hs: TS_plot_NO_TS_MO_Akkaert_NO_TS_MO_Akkaert




NO_TS_MO_AWG_Hs: TS_plot_NO_TS_MO_AWG_NO_TS_MO_AWG




NO_TS_MO_AWG_Tp: TS_plot_NO_TS_MO_AWG_NO_TS_MO_AWG




NO_TS_MO_Deurlo_Hs: TS_plot_NO_TS_MO_Deurlo_NO_TS_MO_Deurlo




NO_TS_MO_Deurlfo_Tp: TS_plot_NO_TS_MO_Deurlfo_NO_TS_MO_Deurlfo




NO_TS_MO_DomburgerRassen_Hs: TS_plot_NO_TS_MO_DomburgerRassen_NO_TS_MO_DomburgerRassen




NO_TS_MO_DomburgerRassen_Tp: TS_plot_NO_TS_MO_DomburgerRassen_NO_TS_MO_DomburgerRassen




NO_TS_MO_EurogeulDWE_Hs: TS_plot_NO_TS_MO_EurogeulDWE_NO_TS_MO_EurogeulDWE




NO_TS_MO_EurogeulDWE_Tp: TS_plot_NO_TS_MO_EurogeulDWE_NO_TS_MO_EurogeulDWE




NO_TS_MO_EurogeulE13_Hs: TS_plot_NO_TS_MO_EurogeulE13_NO_TS_MO_EurogeulE13




NO_TS_MO_EurogeulE13_Tp: TS_plot_NO_TS_MO_EurogeulE13_NO_TS_MO_EurogeulE13




NO_TS_MO_Europlatform3_Hs: TS_plot_NO_TS_MO_Europlatform3_NO_TS_MO_Europlatform3




NO_TS_MO_Europlatform3_Tp: TS_plot_NO_TS_MO_Europlatform3_NO_TS_MO_Europlatform3




NO_TS_MO_F3platform_Hs: TS_plot_NO_TS_MO_F3platform_NO_TS_MO_F3platform




NO_TS_MO_F3platform_Tp: TS_plot_NO_TS_MO_F3platform_NO_TS_MO_F3platform




NO_TS_MO_IJmuidenMunitiestort2_Hs: TS_plot_NO_TS_MO_IJmuidenMunitiestort2_NO_TS_MO_IJmuidenMunitiestort2




NO_TS_MO_IJmuidenMunitiestort2_Tp: TS_plot_NO_TS_MO_IJmuidenMunitiestort2_NO_TS_MO_IJmuidenMunitiestort2




NO_TS_MO_J61_Hs: TS_plot_NO_TS_MO_J61_NO_TS_MO_J61




NO_TS_MO_J61_Tp: TS_plot_NO_TS_MO_J61_NO_TS_MO_J61




NO_TS_MO_K13a3_Hs: TS_plot_NO_TS_MO_K13a3_NO_TS_MO_K13a3




NO_TS_MO_K13a3_Tp: TS_plot_NO_TS_MO_K13a3_NO_TS_MO_K13a3




NO_TS_MO_K141_Hs: TS_plot_NO_TS_MO_K141_NO_TS_MO_K141




NO_TS_MO_K141_Tp: TS_plot_NO_TS_MO_K141_NO_TS_MO_K141




NO_TS_MO_Kwintebank_Hs: TS_plot_NO_TS_MO_Kwintebank_NO_TS_MO_Kwintebank




NO_TS_MO_L91_Hs: TS_plot_NO_TS_MO_L91_NO_TS_MO_L91




NO_TS_MO_L91_Tp: TS_plot_NO_TS_MO_L91_NO_TS_MO_L91




NO_TS_MO_LichteilandGoeree2_Hs: TS_plot_NO_TS_MO_LichteilandGoeree2_NO_TS_MO_LichteilandGoeree2




NO_TS_MO_LichteilandGoeree2_Tp: TS_plot_NO_TS_MO_LichteilandGoeree2_NO_TS_MO_LichteilandGoeree2




NO_TS_MO_MeetboeiWEO1_Hs: TS_plot_NO_TS_MO_MeetboeiWEO1_NO_TS_MO_MeetboeiWEO1




NO_TS_MO_MeetboeiWEW1_Hs: TS_plot_NO_TS_MO_MeetboeiWEW1_NO_TS_MO_MeetboeiWEW1




NO_TS_MO_SchiermonnikoogNoord_Hs: TS_plot_NO_TS_MO_SchiermonnikoogNoord_NO_TS_MO_SchiermonnikoogNoord




NO_TS_MO_SchiermonnikoogNoord_Tp: TS_plot_NO_TS_MO_SchiermonnikoogNoord_NO_TS_MO_SchiermonnikoogNoord




NO_TS_MO_SchiermonnikoogWestgat_Hs: TS_plot_NO_TS_MO_SchiermonnikoogWestgat_NO_TS_MO_SchiermonnikoogWestgat




NO_TS_MO_SchiermonnikoogWestgat_Tp: TS_plot_NO_TS_MO_SchiermonnikoogWestgat_NO_TS_MO_SchiermonnikoogWestgat




NO_TS_MO_Schouwenbank2_Hs: TS_plot_NO_TS_MO_Schouwenbank2_NO_TS_MO_Schouwenbank2




NO_TS_MO_Schouwenbank2_Tp: TS_plot_NO_TS_MO_Schouwenbank2_NO_TS_MO_Schouwenbank2




NO_TS_MO_StortemelkBoei_Hs: TS_plot_NO_TS_MO_StortemelkBoei_NO_TS_MO_StortemelkBoei




NO_TS_MO_StortemelkBoei_Tp: TS_plot_NO_TS_MO_StortemelkBoei_NO_TS_MO_StortemelkBoei




NO_TS_MO_WaddenEierlandseGat_Hs: TS_plot_NO_TS_MO_WaddenEierlandseGat_NO_TS_MO_WaddenEierlandseGat




NO_TS_MO_WaddenEierlandseGat_Tp: TS_plot_NO_TS_MO_WaddenEierlandseGat_NO_TS_MO_WaddenEierlandseGat




Ula_Hs: TS_plot_Ula_Ula



Valhall A_Hs: TS_plot_Valhall A_Valhall A




VINGA BS_Hs: TS_plot_VINGA BS_VINGA BS




VINGA BS_Tp: TS_plot_VINGA BS_VINGA BS

Appendix C: Calibration of significant wave height


NO_TS_MO_6200041




NO_TS_MO_6200041




NO_TS_MO_6200289




NO_TS_MO_6201011




NO_TS_MO_EurogeulE13




NO_TS_MO_Europlatform3




NO_TS_MO_F3platform




NO_TS_MO_IJmuidenMunitiestort2




NO_TS_MO_J61




NO_TS_MO_K13a3




NO_TS_MO_K141




NO_TS_MO_L91




Appendix D: Wave Roses at Selected Locations





Point 1 - Baltic Sea






Point 2 - Kattegat





Point 3 - North Sea




Appendix E: Wind Farms Spatial Impact on Significant Wave Heights

All the figures in Appendix present:


- Left: Baseline - averaged (monthly or yearly) significant wave height, H_{m0} , simulated without wind farms within the domain,
- Mid: Difference between the averaged H_{m0} simulated with the Current scenario and the Baseline scenario,
- Right: Difference between the averaged H_{m0} simulated with the Future scenario and the Baseline scenario.

Baseline


Annual 2019
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Jan/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Feb/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Mar/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Apr/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


May/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Jun/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Jul/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Aug/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Sep/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Oct/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Nov/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Dec/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Annual 2019
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Jan/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Feb/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Mar/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Apr/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


May/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Jun/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Jul/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Aug/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Sep/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Oct/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Nov/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]




0 100 200 km




Baseline


Dec/19
Average




Current Scenario - Baseline




Future Scenario - Baseline



Significant Wave Height [m]



Difference: Year - Baseline [m]






0 100 200 km




Appendix F: Wind Farms Impact on Significant Wave Heights, Danish waters

All the figures in Appendix present:



- Left: Baseline - averaged (monthly or yearly) significant wave height, H_{m0} , simulated without wind farms within the domain,
- Mid: Difference between the averaged H_{m0} simulated with the Current scenario and the Baseline scenario,
- Right: Difference between the averaged H_{m0} simulated with the Future scenario and the Baseline scenario.






Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]










Significant Wave Height [m]




Difference: Year - Baseline [m]







Significant Wave Height [m]




Difference: Year - Baseline [m]



Appendix G: Waves, Annual and Monthly variations at selected locations

Analysis of the annual and monthly wave climate based on the Atlantic- European North West Shelf- Wave Physics Reanalysis” wave model, AENWS.



Locations of extracted AENWS significant wave height (H_{m0}) and mean wave direction (MWD).

Location of the extracted AENWS significant wave height (H_{m0}) and mean wave direction (MWD).

Location	Easterly coordinate [m]	Northerly coordinate [m]	Coordinate System
1	53878.5	6178532.8	UTM32
2	59055.8	6234090.4	UTM32
3	69412.3	6301994.4	UTM32
4	110637.5	6362581.3	UTM32
5	194011.0	6406269.6	UTM32
6	265128.3	6335758.0	UTM32
7	263095.0	6375780.9	UTM32
8	226193.4	6318429.9	UTM32
9	246624.0	6284199.1	UTM32
10	289605.8	6303538.8	UTM32
11	291627.6	6277955.9	UTM32
12	315748.9	6229539.8	UTM32
13	341136.4	6080754.1	UTM32
14	318478.2	6062920.4	UTM32
15	348974.3	6123958.3	UTM32
16	264207.7	6222170.6	UTM32

P1

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	275.3 °	6.6 °	0.2 °	-0.7 °	1.22	0.08	-0.02	-0.03
Jan	267.5 °	30.3 °	0.4 °	-0.6 °	1.59	0.37	-0.01	-0.03
Feb	269.9 °	26.0 °	0.1 °	-0.3 °	1.49	0.38	-0.02	-0.03
Mar	274.3 °	32.2 °	-0.0 °	-0.6 °	1.28	0.30	-0.02	-0.03
Apr	293.6 °	29.2 °	-2.1 °	-0.8 °	1.00	0.21	-0.01	-0.02
May	294.3 °	30.4 °	0.2 °	-1.0 °	0.93	0.16	-0.02	-0.05
Jun	294.1 °	16.9 °	-0.5 °	-1.1 °	0.98	0.16	-0.02	-0.03
Jul	288.0 °	16.4 °	-0.1 °	-0.9 °	0.96	0.20	-0.02	-0.04
Aug	279.4 °	19.8 °	0.5 °	0.2 °	1.00	0.18	-0.02	-0.03
Sep	278.4 °	28.0 °	0.2 °	-0.5 °	1.17	0.24	-0.02	-0.03
Oct	262.3 °	24.4 °	1.2 °	-0.0 °	1.37	0.22	-0.02	-0.03
Nov	265.1 °	35.9 °	-3.3 °	-6.6 °	1.43	0.29	-0.01	-0.03
Dec	267.8 °	26.9 °	-0.1 °	-0.0 °	1.52	0.32	-0.02	-0.03

P1																																					
	Direction, average [°N]																																				
Annual	275	278	273	276	276	282	272	283	280	278	273	270	265	280	266	273	278	283	268	283	268	267	293	268	275	275	258	269	273	278	270	272	265	283	273	276	270
Jan	231	266	249	254	291	260	268	267	67	276	263	249	281	222	236	277	249	281	243	277	248	243	308	280	281	272	67	270	235	285	248	297	262	269	288	260	280
Feb	271	259	244	270	278	295	60	265	274	254	269	292	266	296	263	236	290	314	268	257	276	297	258	225	287	304	324	277	271	246	262	258	261	240	272	291	264
Mar	283	259	280	239	277	290	268	265	345	278	285	259	300	220	281	257	264	282	277	279	281	274	284	287	297	36	270	268	275	273	19	275	259	279	264	275	241
Apr	302	314	283	273	285	220	269	310	266	302	225	285	281	276	258	280	255	262	278	305	267	254	281	292	278	267	292	297	289	296	267	10	301	310	307	296	262
May	223	294	315	316	273	308	311	306	302	281	306	271	272	302	246	249	312	285	261	278	333	269	319	260	305	283	295	267	300	288	276	302	303	266	282	320	154
Jun	314	295	271	276	327	305	284	297	293	307	270	264	286	297	267	279	282	300	299	282	275	310	317	273	275	283	314	288	267	272	302	258	267	305	275	305	282
Jul	242	312	295	268	286	284	311	287	295	300	282	284	309	262	282	273	290	293	281	272	258	259	264	296	274	300	280	280	270	280	311	297	279	293	293	264	264
Aug	265	287	284	289	292	290	290	320	237	273	291	293	289	299	286	302	253	288	298	291	253	258	280	269	268	266	264	223	275	267	266	244	262	295	290	283	254
Sep	276	294	298	290	248	310	281	288	314	290	233	219	69	310	304	284	279	278	255	298	240	298	285	256	273	284	310	271	244	252	269	282	276	286	299	250	292
Oct	258	274	239	269	300	296	266	253	259	297	277	267	224	236	292	294	227	226	248	281	269	284	265	261	266	239	243	259	349	276	290	256	242	256	248	279	258
Nov	295	266	299	239	267	61	270	268	274	66	252	268	210	299	64	223	293	267	266	300	264	229	310	230	249	287	60	265	273	284	66	222	254	283	209	255	286
Dec	289	279	273	285	257	267	256	325	281	244	264	273	246	298	60	282	275	301	257	257	275	251	310	274	234	254	276	246	274	280	268	253	223	260	253	270	277

P1																																					
	H _{ms} , average [m]																																				
	988	989	990	991	992	993	994	995	996	997	998	999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	131	13	138	124	126	122	126	127	105	117	13	126	132	116	109	108	124	127	117	133	128	116	109	128	127	117	117	134	116	127	111	124	129	116	125	123	12
Jan	138	169	192	162	163	229	187	187	691	694	143	154	197	107	16	154	133	229	113	236	189	13	103	139	181	117	149	196	143	143	138	174	179	126	191	163	174
Feb	188	195	216	111	15	145	688	199	128	198	177	177	188	136	2101	682	148	144	108	121	169	106	687	143	136	698	163	147	157	126	12	13	210	107	216	185	154
Mar	137	151	196	687	159	124	183	187	688	138	154	113	158	688	134	696	117	116	106	137	176	117	119	116	126	693	12	124	164	118	687	175	135	127	682	141	103
Apr	695	694	112	114	105	681	108	132	68	148	681	116	685	1	674	694	682	699	114	106	679	68	699	101	102	699	694	125	114	133	683	68	106	135	154	682	117
May	673	697	675	117	689	685	681	679	698	691	694	689	687	687	684	687	105	113	106	1	655	1	105	104	104	693	683	133	688	678	665	117	109	619	103	1	663
Jun	699	673	681	109	68	106	127	101	112	684	102	692	127	105	102	68	109	108	691	689	113	102	68	68	119	116	1	117	682	108	112	685	672	673	68	673	114
Jul	103	114	131	678	691	117	695	678	105	667	134	687	108	677	696	68	685	689	688	117	689	102	687	116	698	679	665	113	102	693	677	104	12	688	123	119	697
Aug	115	128	1	105	127	116	104	103	674	685	133	68	681	697	696	105	685	116	687	118	112	104	105	103	689	697	123	672	11	103	1	698	681	108	675	698	105
Sep	115	104	142	138	116	693	123	122	697	148	689	689	102	119	103	108	162	106	104	115	692	143	125	132	154	108	688	119	689	692	152	144	119	698	116	1	695
Oct	125	163	134	138	111	107	131	131	143	157	154	149	146	151	119	12	134	109	127	103	176	138	138	154	142	135	133	682	162	161	153	127	136	159	138	154	123
Nov	156	134	113	167	18	69	141	125	162	114	107	151	156	184	694	119	16	155	214	188	172	161	129	114	141	161	107	161	135	177	108	106	158	154	142	129	138
Dec	195	139	166	164	137	175	181	688	108	118	155	216	138	13	696	156	169	154	182	134	116	114	117	225	13	211	18	195	157	178	142	16	126	124	137	16	162

P2

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	276.6 °	5.8 °	0.2 °	0.4 °	1.46	0.11	-0.01	-0.05
Jan	269.8 °	26.8 °	0.0 °	-0.1 °	1.95	0.50	0.00	-0.03
Feb	272.3 °	21.2 °	0.4 °	1.5 °	1.83	0.50	-0.01	-0.05
Mar	276.2 °	26.7 °	0.2 °	0.4 °	1.54	0.38	-0.01	-0.04
Apr	292.1 °	20.1 °	-1.0 °	9.0 °	1.18	0.27	-0.01	-0.03
May	292.1 °	19.3 °	0.0 °	-0.2 °	1.09	0.21	-0.01	-0.06
Jun	292.4 °	14.5 °	0.4 °	1.7 °	1.16	0.20	-0.02	-0.05
Jul	287.3 °	14.2 °	0.4 °	0.5 °	1.13	0.23	-0.01	-0.06
Aug	279.3 °	16.9 °	0.9 °	1.5 °	1.17	0.22	-0.02	-0.06
Sep	279.1 °	24.4 °	0.4 °	0.2 °	1.39	0.32	0.00	-0.06
Oct	265.5 °	22.7 °	0.7 °	0.6 °	1.61	0.27	-0.01	-0.05
Nov	268.1 °	30.0 °	-9.1 °	-6.9 °	1.71	0.37	-0.02	-0.05
Dec	270.1 °	24.1 °	0.0 °	0.3 °	1.83	0.45	-0.01	-0.04

		P2																																				
		Direction, average [°N]																																				
		888	889	890	891	892	893	894	895	896	897	898	899	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	276	278	274	278	276	283	274	283	281	277	275	272	267	280	270	274	277	283	270	283	269	289	293	270	276	276	262	270	275	278	273	273	267	283	274	278	273	
	Jan	236	267	252	238	290	263	271	267	166	290	267	252	281	232	239	278	251	290	251	278	251	248	336	279	280	277	204	271	244	284	252	296	264	288	268	262	281
Feb	271	261	248	267	278	293	199	267	279	257	271	291	267	298	265	245	269	311	290	264	271	296	290	237	286	300	228	276	271	258	266	260	262	259	273	290	269	
Mar	283	261	280	246	278	287	269	268	335	277	284	263	299	242	281	262	268	283	281	280	280	274	285	288	295	7	273	271	277	276	42	275	260	280	270	278	255	
Apr	300	308	282	274	287	241	267	311	265	299	236	284	284	276	268	282	259	272	278	301	268	263	280	290	279	270	291	295	288	295	276	346	298	310	306	294	264	
May	296	292	310	310	272	299	305	302	302	283	302	274	272	300	249	252	309	283	265	278	320	271	315	262	302	283	294	268	294	285	287	301	300	268	280	310	257	
Jun	310	291	271	278	317	303	283	298	290	303	270	269	285	294	269	278	283	297	295	290	275	310	310	274	278	282	311	287	286	275	300	262	271	300	278	302	281	
Jul	244	308	293	273	285	284	301	285	293	297	281	285	307	264	281	272	290	293	282	273	280	280	285	293	278	297	279	281	270	280	305	295	279	293	292	287	269	
Aug	287	285	282	288	254	280	290	315	290	272	289	292	288	270	285	301	298	288	298	289	254	298	279	289	272	268	266	236	278	267	268	248	269	295	285	285	258	
Sep	277	281	298	289	230	342	286	289	308	289	249	225	210	308	299	283	278	276	280	296	247	285	287	298	275	284	308	271	248	258	260	282	279	287	298	298	294	
Oct	264	275	245	272	301	298	270	296	282	296	279	269	228	241	298	294	234	235	250	279	269	283	268	263	270	245	246	271	346	277	288	260	246	258	251	287	258	
Nov	293	268	298	262	268	238	272	292	278	185	295	268	217	297	210	231	292	268	267	298	284	235	312	241	252	286	201	267	277	284	209	244	297	282	216	283	285	
Dec	289	280	274	285	280	269	298	310	281	248	267	274	249	297	82	283	274	300	260	259	277	253	310	274	241	257	278	248	275	281	270	256	229	267	256	274	277	

P2																																					
		H _{ms} , average [m]																																			
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	152	154	164	146	148	143	147	149	121	137	152	147	154	131	125	125	144	149	136	158	149	132	125	151	148	137	136	159	135	149	13	144	152	134	146	143	14
Jan	137	209	237	195	199	248	226	221	0.93	10	189	184	243	123	191	18	151	282	133	289	226	154	112	165	215	136	169	244	164	172	161	207	216	146	231	92	208
Feb	221	238	268	128	18	173	0.98	239	146	24	216	216	227	199	238	0.93	171	166	121	137	209	118	0.92	164	162	111	195	175	186	145	14	154	239	122	238	189	18
Mar	153	179	24	1	185	148	216	217	0.97	166	178	133	185	0.94	159	112	135	129	117	18	203	137	139	139	151	0.99	147	144	123	141	0.94	206	16	151	0.94	161	117
Apr	11	108	134	135	125	0.88	124	154	0.69	172	0.85	133	0.94	110	0.84	103	0.93	1	134	129	0.89	0.65	114	12	117	12	108	149	129	107	0.93	0.61	124	156	119	0.92	134
May	0.78	114	0.88	138	101	0.94	0.91	0.91	114	105	107	101	0.99	0.99	0.95	102	123	133	117	119	0.6	116	123	123	124	109	0.94	157	1	0.86	0.71	137	13	0.97	12	116	0.67
Jun	117	0.86	0.95	125	0.93	128	152	116	132	0.96	119	105	147	123	116	104	125	127	107	0.78	132	118	107	104	139	132	114	14	0.92	126	134	1	0.79	0.87	0.94	0.84	137
Jul	12	133	153	0.89	107	137	0.84	0.92	124	0.76	157	101	121	0.87	109	0.91	0.99	103	0.7	136	0.97	116	102	128	112	0.97	0.72	13	119	109	0.92	12	147	101	148	136	113
Aug	135	154	118	125	146	133	12	122	0.82	0.85	156	1	111	112	0.83	125	0.94	136	0.96	133	125	121	122	118	102	112	147	0.83	13	12	119	118	0.91	123	0.87	118	127
Sep	177	123	169	164	132	107	141	141	107	183	0.99	0.99	119	133	112	138	193	127	119	257	103	117	143	159	182	124	0.99	136	105	103	187	17	138	109	129	11	108
Oct	147	19	152	158	124	121	149	16	171	182	229	169	171	177	135	14	154	123	143	12	212	159	159	182	165	152	158	105	116	213	184	146	154	188	183	143	
Nov	184	157	129	2	217	0.99	166	144	19	124	122	18	175	215	104	132	169	163	245	223	202	19	153	135	165	193	119	214	159	21	12	114	188	182	182	144	164
Dec	235	184	198	199	185	202	216	0.96	127	131	182	246	199	147	1	185	205	18	224	18	136	125	129	216	146	254	215	236	194	21	164	189	144	141	156	19	192

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	277.4°	5.6°	-0.1°	1.0°	1.46	0.11	-0.01	-0.02
<i>Jan</i>	272.0°	26.6°	-0.0°	0.3°	1.95	0.50	0.00	-0.02
<i>Feb</i>	275.3°	25.2°	-0.0°	1.2°	1.83	0.50	0.00	-0.03
<i>Mar</i>	278.3°	22.9°	-0.1°	0.4°	1.54	0.38	0.00	-0.02
<i>Apr</i>	292.2°	18.4°	-0.9°	0.9°	1.18	0.27	-0.01	-0.02
<i>May</i>	290.1°	19.2°	-0.1°	-0.0°	1.09	0.21	-0.01	-0.02
<i>Jun</i>	288.3°	13.2°	-1.0°	1.6°	1.16	0.20	-0.01	-0.04
<i>Jul</i>	283.8°	12.5°	0.6°	0.9°	1.13	0.23	0.00	-0.02
<i>Aug</i>	277.1°	15.2°	0.2°	2.2°	1.17	0.22	-0.01	-0.04
<i>Sep</i>	279.3°	23.5°	0.3°	0.5°	1.39	0.32	0.00	-0.02
<i>Oct</i>	267.6°	23.5°	0.1°	1.6°	1.61	0.27	-0.01	-0.03
<i>Nov</i>	271.0°	28.0°	-1.9°	1.0°	1.71	0.37	-0.03	-0.04
<i>Dec</i>	272.3°	25.1°	0.0°	0.7°	1.83	0.45	-0.01	-0.02

P3																																						
Direction, average [°N]																																						
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	277	277	273	278	277	284	275	283	282	277	276	272	268	280	275	275	277	282	271	283	270	269	295	270	278	277	285	271	278	279	278	275	268	284	275	281	275	
Jan	240	268	254	258	269	265	272	266	26	279	269	253	280	236	260	278	254	251	259	278	254	252	351	277	281	283	269	272	250	262	257	294	265	292	288	262	284	
Feb	272	263	249	306	278	291	31	267	281	281	271	290	267	302	265	258	288	313	298	285	272	296	347	247	285	304	230	277	271	268	274	260	264	271	273	289	275	
Mar	283	262	280	251	278	285	270	269	310	268	297	260	282	288	277	278	290	265	277	275	297	266	266	278	288	282	271	293	291	287	293	280	3	293	308	307	298	
Apr	300	309	280	273	287	265	265	310	268	297	260	282	288	277	278	290	265	277	275	297	266	266	278	288	282	271	293	291	287	293	280	3	293	308	307	298		
May	274	287	307	310	270	299	301	297	306	284	299	284	270	295	230	231	306	279	266	277	314	269	311	281	299	281	292	266	291	286	294	299	295	272	278	306	346	
Jun	305	284	267	272	312	299	280	298	284	300	268	268	280	289	268	275	280	291	287	296	272	305	308	272	274	280	308	283	284	275	295	263	271	286	278	296	277	
Jul	244	302	288	274	282	281	289	281	289	293	278	280	302	263	278	271	288	290	277	271	281	238	262	288	273	281	280	281	266	280	297	291	277	292	287	268	268	
Aug	265	280	278	284	235	287	290	308	258	269	286	289	284	267	286	297	263	284	301	286	230	237	278	268	273	267	264	244	275	265	265	245	273	294	279	285	258	
Sep	276	286	296	284	250	301	297	295	303	286	279	232	215	311	296	280	276	272	263	294	254	282	292	255	275	280	308	269	250	258	268	282	282	290	302	297	293	
Oct	270	275	247	276	307	299	270	295	261	293	262	271	231	243	312	293	238	240	251	277	268	284	267	263	273	250	247	278	303	276	288	260	250	258	251	300	259	
Nov	281	268	299	262	268	290	273	298	278	293	260	269	220	294	288	238	289	268	267	296	283	238	323	247	254	283	208	267	283	284	218	307	258	282	219	281	284	
Dec	288	280	278	284	262	272	257	310	281	252	268	275	252	298	311	283	274	301	261	258	279	257	310	274	256	257	279	250	278	281	272	258	234	275	262	280	279	

			P3																																		
			Hm0, average [m]																																		
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	154	159	168	15	152	149	15	152	124	14	154	148	155	133	128	128	146	154	138	181	152	134	129	156	152	141	139	163	138	154	134	146	157	138	149	147	145
Jan	157	227	244	196	210	302	23	221	0.93	124	174	167	232	124	195	186	151	293	141	292	231	18	12	169	219	143	169	230	162	18	165	209	228	15	24	193	215
Feb	221	25	287	137	19	184	105	245	152	25	226	219	233	168	237	0.98	173	175	125	142	223	118	0.95	168	172	119	198	184	189	15	147	159	218	132	262	202	186
Mar	147	181	235	103	186	109	224	216	103	177	18	143	191	0.95	168	117	136	129	12	167	2	143	141	149	16	107	16	152	127	152	101	205	166	157	898	157	124
Apr	114	114	143	14	13	0.91	127	162	0.77	174	0.86	133	0.95	115	0.88	105	0.95	105	136	134	0.9	0.87	118	126	119	13	112	155	13	165	0.98	0.68	131	16	12	0.96	136
May	0.78	12	0.91	142	103	0.95	0.9	0.92	122	109	107	104	1	101	0.97	105	128	135	117	123	0.82	118	125	127	13	114	0.95	158	103	0.88	0.75	142	135	0.95	121	122	0.7
Jun	118	0.9	0.96	122	0.99	131	158	118	134	0.98	122	104	148	124	116	104	124	129	111	0.8	135	119	114	107	139	132	112	144	0.91	129	137	105	0.79	0.96	0.99	0.84	142
Jul	122	134	152	0.92	11	137	0.87	0.95	129	0.76	195	101	121	0.87	108	0.92	102	105	0.75	137	0.95	118	106	122	110	103	0.73	131	118	112	1	123	158	103	153	135	116
Aug	137	157	12	136	143	132	119	124	0.84	0.66	185	0.95	119	112	0.85	127	0.93	14	0.95	135	119	124	125	119	104	114	149	0.87	134	121	119	118	0.91	125	0.89	118	132
Sep	179	128	167	167	13	116	145	144	106	191	1	0.98	110	132	118	132	193	124	121	212	105	172	147	159	187	123	1	131	11	104	195	175	146	118	129	124	106
Oct	151	188	151	150	125	124	149	169	174	179	228	167	169	177	142	144	153	123	137	124	22	150	159	187	165	149	16	11	129	210	101	144	153	119	185	192	146
Nov	187	159	133	206	218	104	17	146	191	122	125	185	168	219	111	132	191	188	249	224	203	191	164	139	17	2	122	212	169	218	121	119	195	189	162	147	169
Dec	242	166	205	21	175	205	218	0.97	13	128	188	246	159	146	101	189	217	167	233	161	142	125	13	283	146	242	222	241	204	218	163	191	148	143	158	193	158

P4

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	280.4 °	5.8 °	-0.1 °	0.3 °	1.42	0.09	0.00	-0.01
<i>Jan</i>	277.5 °	33.9 °	0.2 °	0.1 °	1.88	0.47	0.00	0.00
<i>Feb</i>	281.9 °	32.5 °	-0.0 °	0.6 °	1.80	0.44	0.00	-0.01
<i>Mar</i>	282.5 °	27.9 °	-0.1 °	-0.1 °	1.51	0.34	0.00	0.00
<i>Apr</i>	295.3 °	26.2 °	-0.5 °	-0.3 °	1.17	0.24	0.00	0.00
<i>May</i>	289.0 °	21.8 °	-0.1 °	0.0 °	1.07	0.20	0.00	-0.01
<i>Jun</i>	281.5 °	10.6 °	-2.1 °	-0.0 °	1.12	0.18	-0.01	-0.02
<i>Jul</i>	278.2 °	9.0 °	0.4 °	0.5 °	1.10	0.21	0.00	-0.01
<i>Aug</i>	275.7 °	14.0 °	-0.0 °	0.5 °	1.13	0.20	-0.01	-0.01
<i>Sep</i>	281.5 °	30.5 °	0.0 °	0.1 °	1.37	0.30	0.01	0.00
<i>Oct</i>	273.4 °	27.2 °	0.0 °	0.7 °	1.55	0.26	0.00	-0.01
<i>Nov</i>	278.1 °	46.3 °	0.2 °	0.0 °	1.65	0.31	-0.03	0.00
<i>Dec</i>	277.9 °	29.6 °	0.1 °	0.2 °	1.77	0.41	0.00	0.00

P4																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	279	276	275	279	279	287	279	283	290	279	278	276	270	284	284	279	277	281	274	282	273	273	303	272	280	279	273	274	282	279	282	279	272	286	277	285	279
Jan	251	271	262	263	283	269	275	269	98	278	276	263	278	248	265	281	257	260	272	277	262	260	35	272	282	298	92	276	274	281	272	295	269	306	285	267	288
Feb	279	267	257	343	281	287	39	268	323	268	273	285	270	309	269	277	284	327	38	345	273	297	33	272	283	339	238	280	273	282	301	265	270	311	274	288	280
Mar	293	266	277	270	281	286	273	271	7	276	278	273	292	284	281	272	279	284	325	288	278	277	280	286	287	31	280	282	276	279	37	273	268	278	276	282	295
Apr	296	326	282	279	289	320	272	304	280	292	45	278	299	285	290	320	279	281	276	288	268	279	278	287	286	271	296	284	287	289	295	35	288	300	310	311	284
May	310	282	297	303	271	306	309	295	310	278	293	297	274	288	239	256	298	271	267	276	317	272	301	267	290	284	291	267	302	291	305	291	286	286	274	295	35
Jun	295	277	264	268	305	286	277	294	275	303	267	269	274	281	269	275	274	281	281	299	270	294	293	270	269	274	284	276	285	275	284	269	276	285	278	289	274
Jul	290	290	279	271	277	276	283	279	281	288	273	275	292	286	276	274	282	284	274	268	287	239	264	281	268	282	288	278	265	278	287	283	274	287	279	268	272
Aug	265	276	276	276	261	285	291	294	310	271	280	285	278	269	316	288	276	278	303	280	280	239	279	271	271	268	282	296	273	266	266	259	277	293	280	282	260
Sep	275	284	296	277	256	22	314	313	314	280	317	245	66	328	294	278	274	271	268	290	283	279	302	299	277	280	310	274	282	277	271	282	285	295	312	265	291
Oct	276	277	295	283	336	307	272	261	264	290	282	278	242	252	335	292	242	252	257	278	267	297	268	268	276	260	257	312	22	279	289	268	259	261	258	316	282
Nov	288	272	314	267	271	5	275	315	284	98	276	274	223	290	34	247	284	269	269	293	284	244	359	261	267	281	16	271	292	281	227	21	265	283	224	341	280
Dec	285	280	279	283	268	279	263	335	290	278	278	279	263	304	59	285	274	310	265	265	291	354	332	274	341	262	280	257	278	279	272	260	245	287	272	290	283

P4																																						
	Hm., average [m]																																					
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	279	276	275	276	276	287	279	283	290	279	278	276	270	284	284	279	277	281	274	282	273	273	303	272	280	279	273	274	282	279	282	279	272	286	277	285	279	
Jan	251	271	262	263	283	269	275	269	98	278	276	275	263	278	248	265	281	257	260	272	277	262	260	35	272	282	298	92	276	274	281	272	295	269	306	285	267	288
Feb	279	267	257	343	281	287	39	268	323	268	273	285	270	309	269	277	284	327	38	345	273	297	33	272	283	339	238	280	273	282	301	265	270	311	274	288	280	
Mar	293	266	277	270	281	286	273	271	7	276	278	273	292	284	281	272	279	284	325	288	278	277	280	286	287	31	280	282	276	279	37	273	288	278	276	282	295	
Apr	296	326	282	279	289	320	272	304	280	292	45	278	299	285	290	320	279	281	276	288	268	279	278	287	286	271	296	284	287	289	295	35	288	300	310	311	284	
May	310	282	297	303	271	306	309	295	310	278	293	297	274	288	239	256	298	271	267	276	317	272	301	267	290	284	291	267	302	291	305	291	286	286	274	295	35	
Jun	295	277	264	268	305	286	277	294	275	303	267	269	274	281	269	276	274	281	281	299	270	294	293	270	269	274	284	276	285	275	284	269	276	285	278	289	274	
Jul	290	290	279	271	277	276	283	279	281	288	273	275	292	286	276	274	282	284	274	268	287	239	264	281	268	282	288	278	265	278	287	283	274	287	279	268	272	
Aug	265	276	276	276	261	285	291	294	316	271	280	285	276	269	316	288	276	278	303	280	280	239	279	271	271	268	282	296	273	266	266	259	277	293	280	282	260	
Sep	275	284	296	277	256	22	314	313	314	280	317	245	66	328	294	278	274	271	268	290	283	279	302	299	277	280	310	274	282	277	271	282	285	295	312	265	291	
Oct	276	277	295	283	336	307	272	261	264	290	282	278	242	252	335	292	242	252	257	278	267	297	268	268	276	260	257	312	22	279	289	268	259	261	258	316	282	
Nov	288	272	314	287	271	5	275	315	284	98	276	274	223	290	34	247	284	269	269	293	284	244	359	261	267	281	16	271	292	281	227	21	265	283	224	341	280	
Dec	285	280	279	283	268	279	263	335	290	278	278	279	263	304	59	285	274	310	265	265	291	354	332	274	341	262	280	257	278	279	272	260	245	287	272	290	283	

P5

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	272.3 °	5.5 °	0.0 °	0.2 °	1.08	0.08	0.00	0.00
<i>Jan</i>	272.4 °	34.7 °	0.1 °	0.0 °	1.41	0.41	0.00	0.00
<i>Feb</i>	275.0 °	35.8 °	0.0 °	0.2 °	1.35	0.40	0.00	0.00
<i>Mar</i>	272.0 °	31.5 °	0.1 °	-0.0 °	1.11	0.30	0.00	0.00
<i>Apr</i>	286.6 °	29.4 °	-0.3 °	0.1 °	0.86	0.20	0.00	0.00
<i>May</i>	276.7 °	21.3 °	0.1 °	-0.1 °	0.80	0.19	0.00	0.00
<i>Jun</i>	267.4 °	7.8 °	-0.5 °	0.6 °	0.87	0.18	0.00	0.00
<i>Jul</i>	264.6 °	5.6 °	0.3 °	0.0 °	0.88	0.20	0.00	0.00
<i>Aug</i>	265.3 °	9.2 °	-0.2 °	0.3 °	0.88	0.18	-0.01	-0.01
<i>Sep</i>	272.9 °	29.5 °	0.0 °	0.1 °	1.04	0.27	0.01	0.01
<i>Oct</i>	269.2 °	26.7 °	0.1 °	0.0 °	1.17	0.22	-0.01	0.00
<i>Nov</i>	275.4 °	48.2 °	0.4 °	-0.8 °	1.23	0.27	-0.01	0.00
<i>Dec</i>	274.1 °	33.5 °	0.4 °	0.5 °	1.33	0.34	0.01	0.02

P5																																					
		Direction, average [°N]																																			
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	279	276	275	276	276	287	279	283	290	279	278	276	270	284	284	279	277	281	274	282	273	273	303	272	280	279	273	274	282	278	282	276	272	286	277	285	279
Jan	251	271	262	263	263	269	275	269	98	278	275	263	278	248	265	281	257	260	272	277	262	260	35	272	282	298	82	276	274	281	272	295	269	306	285	267	288
Feb	279	267	257	343	281	287	39	268	323	268	273	285	270	309	269	277	284	327	338	345	273	297	33	272	283	339	238	280	273	282	301	265	270	311	274	288	280
Mar	293	266	277	270	281	286	273	271	7	276	278	273	292	284	281	272	279	284	325	288	278	277	280	286	287	31	280	282	276	279	37	273	288	278	276	282	295
Apr	296	326	282	279	289	320	272	304	280	292	45	278	299	285	290	320	279	281	278	288	268	279	278	287	286	271	296	284	287	289	295	35	288	300	310	311	284
May	310	282	297	303	271	306	309	295	310	278	293	297	274	288	239	256	298	271	267	276	317	272	301	267	290	284	291	267	302	291	305	291	286	286	274	295	35
Jun	295	277	264	268	305	286	277	294	275	303	267	269	274	281	269	275	274	281	281	299	270	284	283	270	269	274	284	276	285	275	284	269	276	285	278	289	274
Jul	290	290	279	271	277	276	283	279	281	288	273	275	292	286	276	274	282	284	274	268	287	239	264	281	268	282	288	278	265	278	287	283	274	287	279	266	272
Aug	265	276	276	279	261	285	291	294	310	271	280	285	276	269	316	288	276	278	303	280	260	258	279	271	271	268	262	256	273	266	266	259	277	293	280	282	260
Sep	275	284	296	277	256	22	314	313	314	280	317	245	156	328	294	278	274	271	268	290	283	279	302	259	277	280	316	274	262	277	271	282	285	295	312	265	291
Oct	276	277	295	283	336	307	272	261	264	290	282	278	242	252	335	292	242	252	257	278	267	297	268	268	276	260	257	312	22	279	289	268	259	261	258	316	282
Nov	288	272	314	267	271	5	275	315	284	98	276	274	223	290	34	247	284	269	269	293	284	244	359	261	267	281	16	271	292	281	227	21	265	283	224	341	280
Dec	285	280	279	283	268	279	263	335	290	278	278	279	263	304	59	285	274	310	265	265	291	354	332	274	341	262	280	257	276	279	272	260	245	287	272	290	283

P5																																					
	Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	107	119	124	108	112	112	111	111	0.92	1	114	105	108	1	0.99	0.94	105	107	0.99	118	111	0.98	0.97	110	111	103	11	125	104	117	0.99	11	123	103	111	112	109
Jan	106	2.02	179	136	163	2.35	156	139	0.8	0.9	128	132	177	0.79	152	143	0.94	2.09	0.94	2.39	171	1.8	0.92	11	145	112	122	151	118	131	121	136	163	107	15	138	159
Feb	139	196	198	116	155	13	0.79	186	106	194	192	143	172	13	178	0.57	115	131	0.86	1	178	0.76	0.73	12	133	0.85	126	132	129	112	107	129	2.04	112	187	156	137
Mar	0.92	132	2.09	0.7	126	128	166	142	0.74	144	116	105	135	0.69	137	0.76	0.94	0.79	0.83	117	13	101	0.96	118	123	0.84	129	116	0.75	119	0.85	145	122	118	0.65	108	0.91
Apr	0.86	0.83	106	0.96	0.98	0.67	0.87	114	0.65	125	0.64	0.9	0.61	0.82	0.62	0.75	0.65	0.69	0.91	103	0.59	0.48	0.9	0.88	0.79	101	0.81	116	0.79	128	0.81	0.56	109	105	0.84	0.72	101
May	0.54	0.96	0.76	0.97	0.79	0.63	0.57	0.75	0.81	0.79	0.72	0.78	0.76	0.8	0.85	0.8	0.95	0.89	0.81	0.99	0.42	0.96	0.84	109	0.92	0.8	0.82	12	0.71	0.68	0.45	113	109	0.73	1	0.76	0.54
Jun	0.66	0.73	0.67	0.91	0.69	1	135	0.82	0.87	0.61	0.99	0.78	101	1	0.97	0.91	0.96	0.87	0.81	0.68	113	0.73	0.9	0.94	0.89	0.87	0.69	122	0.6	114	0.9	0.88	0.59	0.9	0.91	0.58	108
Jul	0.95	0.87	103	0.75	0.92	114	0.92	0.81	11	0.93	11	0.81	0.81	0.71	0.81	0.88	0.84	0.77	0.65	112	0.65	0.92	0.83	0.87	0.86	0.83	0.81	108	0.95	0.91	0.78	0.94	142	0.73	116	12	0.96
Aug	1	110	0.93	1	116	1	0.81	0.78	0.63	0.43	119	0.55	0.95	0.94	0.92	0.93	0.73	0.93	0.66	102	0.92	102	101	0.84	0.72	0.84	108	0.74	110	0.94	0.92	0.92	0.67	0.99	0.7	0.94	109
Sep	136	0.94	121	119	0.85	0.93	117	107	0.72	143	0.75	0.69	0.8	0.98	0.81	103	135	103	0.92	152	0.79	131	11	126	156	0.78	0.69	0.91	0.86	0.74	164	138	132	0.85	0.88	0.95	0.73
Oct	106	129	105	0.95	0.98	0.85	104	131	121	126	163	122	105	13	121	102	111	0.82	0.93	0.85	167	112	101	136	113	109	123	0.87	119	107	144	105	108	136	125	166	106
Nov	131	105	0.92	145	147	0.78	129	118	152	0.75	0.88	133	0.99	161	0.9	0.87	132	128	173	14	141	134	147	104	129	14	0.92	153	139	153	0.78	0.87	147	14	126	109	126
Dec	17	118	137	158	127	15	147	0.68	0.99	0.82	147	175	111	106	0.78	151	162	135	18	114	0.95	0.86	0.97	2.02	114	191	153	187	158	161	11	14	0.96	102	112	149	148

P6

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	198.7°	15.0°	-0.0°	0.6°	0.67	0.03	0.00	0.00
<i>Jan</i>	199.1°	40.5°	-0.8°	2.1°	0.85	0.15	0.00	0.00
<i>Feb</i>	194.8°	59.6°	-0.1°	-0.6°	0.81	0.16	0.00	0.00
<i>Mar</i>	191.2°	45.8°	0.1°	0.1°	0.68	0.12	0.00	0.00
<i>Apr</i>	190.5°	59.4°	-0.1°	-1.3°	0.55	0.07	0.00	-0.01
<i>May</i>	196.1°	60.1°	0.3°	-0.3°	0.50	0.08	0.00	0.00
<i>Jun</i>	217.1°	31.9°	-0.3°	-0.5°	0.50	0.10	0.00	0.00
<i>Jul</i>	220.6°	23.6°	-0.1°	2.1°	0.49	0.11	0.00	-0.01
<i>Aug</i>	209.2°	35.9°	-0.2°	0.5°	0.55	0.11	0.00	0.00
<i>Sep</i>	200.5°	47.3°	0.2°	1.5°	0.67	0.11	0.00	0.00
<i>Oct</i>	190.9°	39.5°	0.5°	0.1°	0.79	0.11	0.00	-0.01
<i>Nov</i>	185.8°	55.7°	-2.3°	-0.7°	0.82	0.11	0.00	-0.01
<i>Dec</i>	198.9°	47.9°	0.6°	0.1°	0.82	0.13	0.00	0.00

P6																																						
Direction, average [°N]																																						
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	204	224	220	212	220	210	205	215	195	221	216	201	200	202	199	209	211	210	199	229	217	191	197	210	216	203	194	212	203	220	189	210	215	207	211	208	197	
Jan	199	241	216	211	207	227	216	191	195	230	209	200	241	197	229	222	199	243	197	245	192	188	190	203	200	194	190	220	190	232	185	283	225	197	296	202	201	
Feb	191	221	209	190	246	205	198	232	192	220	242	258	225	274	224	192	245	191	198	193	238	198	191	191	252	30	196	209	219	190	141	219	225	198	229	258	202	
Mar	196	207	200	190	214	208	220	200	195	234	228	190	269	195	244	191	199	190	190	196	218	209	191	222	237	252	184	222	193	198	216	192	229	209	232	190	208	146
Apr	249	17	228	177	209	195	190	304	09	293	191	242	192	199	194	194	190	199	211	272	144	199	219	251	198	205	193	268	201	259	195	108	253	298	313	191	199	
May	192	253	288	327	227	196	147	281	198	191	255	194	205	283	145	201	268	218	198	232	184	240	321	211	228	195	199	224	190	192	199	254	275	194	233	191	191	
Jun	330	231	193	228	241	265	254	235	233	198	235	190	242	253	231	240	239	263	234	290	241	303	263	241	225	219	280	247	199	240	290	192	194	223	239	234	227	
Jul	196	275	292	239	246	203	203	247	295	307	243	210	210	225	228	241	253	240	242	240	228	223	212	199	233	267	198	239	236	235	298	294	248	238	254	228	228	
Aug	226	239	226	254	210	247	214	303	198	198	290	147	240	219	199	274	211	235	190	226	229	214	219	212	223	226	209	147	240	220	229	209	198	291	243	252	207	
Sep	238	225	270	233	193	195	197	198	193	259	149	191	195	195	214	210	232	207	196	247	195	239	193	216	236	191	192	195	219	145	232	236	236	195	199	198	198	
Oct	192	225	190	190	190	194	190	213	200	290	210	205	197	195	198	237	191	193	205	228	224	145	199	214	214	192	195	199	190	230	223	211	193	208	212	198	199	
Nov	251	193	213	201	210	196	219	190	205	145	194	198	191	265	199	190	250	198	214	277	234	195	193	198	232	197	220	205	231	149	196	214	216	190	191	234		
Dec	296	208	196	252	200	195	210	192	199	192	213	218	194	198	198	238	231	190	221	198	204	197	199	226	194	214	220	208	242	230	191	206	196	193	193	204	207	

P6																																						
		Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	0.68	0.68	0.71	0.66	0.65	0.68	0.7	0.69	0.61	0.62	0.7	0.67	0.71	0.64	0.65	0.61	0.66	0.66	0.63	0.68	0.68	0.65	0.62	0.7	0.69	0.64	0.71	0.74	0.65	0.71	0.65	0.67	0.72	0.62	0.66	0.67	0.68	
Jan	0.88	0.94	102	0.8	0.76	1.6	0.88	0.97	0.99	0.92	0.73	0.81	0.9	0.73	0.85	0.82	0.83	0.97	0.99	0.76	106	1.6	0.75	0.39	0.65	0.84	0.7	121	0.95	0.84	0.72	0.82	0.79	0.93	0.67	0.94	0.85	0.9
Feb	0.95	103	113	0.82	0.78	0.66	0.72	0.98	0.76	108	0.95	0.72	0.9	0.79	0.98	0.47	0.63	0.87	0.99	0.91	0.93	0.96	0.95	0.7	0.5	0.96	0.81	0.72	0.8	0.79	0.76	1	0.75	0.92	0.77	0.8		
Mar	0.72	0.84	0.95	0.62	0.75	0.68	0.81	1	0.98	0.76	0.73	0.69	0.7	0.57	0.74	0.85	0.66	0.53	0.54	0.69	0.83	0.63	0.51	0.66	0.69	0.67	0.71	0.68	0.45	0.67	0.63	0.76	0.75	0.63	0.51	0.65	0.66	
Apr	0.5	0.64	0.63	0.62	0.54	0.55	0.62	0.62	0.37	0.68	0.58	0.57	0.47	0.53	0.44	0.62	0.51	0.5	0.54	0.56	0.4	0.42	0.59	0.49	0.55	0.58	0.53	0.58	0.55	0.67	0.56	0.49	0.62	0.56	0.47	0.54	0.62	
May	0.38	0.35	0.37	0.49	0.45	0.43	0.48	0.49	0.33	0.52	0.46	0.33	0.33	0.44	0.31	0.49	0.56	0.49	0.62	0.35	0.28	0.35	0.48	0.65	0.53	0.53	0.39	0.72	0.46	0.48	0.35	0.8	0.55	0.43	0.56	0.48	0.47	
Jun	0.31	0.38	0.39	0.6	0.37	0.49	0.7	0.52	0.38	0.48	0.59	0.54	0.59	0.54	0.62	0.59	0.59	0.52	0.45	0.39	0.63	0.47	0.42	0.52	0.59	0.6	0.37	0.62	0.51	0.71	0.47	0.52	0.4	0.4	0.47	0.31	0.59	
Jul	0.62	0.41	0.6	0.39	0.52	0.99	0.29	0.44	0.96	0.35	0.99	0.41	0.4	0.4	0.49	0.4	0.43	0.43	0.3	0.87	0.38	0.95	0.5	0.45	0.5	0.44	0.4	0.62	0.49	0.95	0.35	0.49	0.75	0.45	0.58	0.69	0.55	
Aug	0.62	0.63	0.55	0.53	0.72	0.99	0.55	0.44	0.57	0.25	0.67	0.37	0.56	0.62	0.42	0.46	0.52	0.53	0.36	0.61	0.6	0.65	0.6	0.59	0.43	0.53	0.74	0.6	0.68	0.59	0.59	0.54	0.42	0.61	0.37	0.53	0.62	
Sep	0.73	0.5	0.89	0.64	0.6	0.71	0.78	0.71	0.96	0.74	0.58	0.69	0.81	0.66	0.5	0.62	0.74	0.61	0.74	0.8	0.59	0.74	0.7	0.77	0.91	0.52	0.51	0.73	0.51	0.59	0.85	0.8	0.67	0.53	0.66	0.65	0.52	
Oct	0.77	0.75	0.8	0.68	0.69	0.63	0.79	0.73	0.77	0.71	103	0.85	0.85	0.9	0.81	0.64	0.9	0.73	0.7	0.49	0.94	0.79	0.79	0.89	0.75	0.83	0.95	0.72	0.84	0.95	0.83	0.64	0.79	0.84	0.73	0.95	0.74	
Nov	0.71	0.76	0.98	0.93	0.88	0.73	0.82	0.79	0.97	0.77	0.89	0.82	102	0.88	0.74	0.77	0.76	0.87	1	0.81	0.85	105	0.88	0.71	0.87	0.77	0.9	0.84	0.93	0.81	0.74	0.76	0.87	0.77	103	0.75	0.66	
Dec	0.93	0.76	0.86	0.81	0.78	0.99	0.92	0.96	0.62	0.64	0.85	1	0.78	0.7	0.75	0.89	0.81	0.87	0.98	0.65	0.58	0.7	0.66	105	0.87	104	0.84	106	0.79	0.92	0.8	0.87	0.52	0.76	0.75	0.85	0.81	

P7

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	236.4 °	25.2 °	0.3 °	0.3 °	0.70	0.03	0.00	0.00
<i>Jan</i>	232.6 °	57.5 °	0.3 °	0.0 °	0.91	0.17	0.00	-0.01
<i>Feb</i>	235.2 °	82.4 °	-0.1 °	0.3 °	0.86	0.17	0.00	0.00
<i>Mar</i>	225.0 °	68.0 °	0.1 °	0.1 °	0.73	0.15	0.00	0.00
<i>Apr</i>	314.5 °	89.6 °	0.6 °	-0.0 °	0.58	0.10	0.00	0.00
<i>May</i>	267.6 °	73.5 °	-0.4 °	-0.7 °	0.52	0.09	0.00	-0.01
<i>Jun</i>	254.5 °	35.9 °	1.9 °	-2.2 °	0.52	0.11	0.00	0.00
<i>Jul</i>	250.8 °	26.7 °	-0.8 °	-1.5 °	0.50	0.11	0.00	-0.01
<i>Aug</i>	237.0 °	45.6 °	-1.6 °	0.8 °	0.55	0.11	0.00	0.00
<i>Sep</i>	239.9 °	72.2 °	0.3 °	1.1 °	0.70	0.13	0.00	-0.01
<i>Oct</i>	213.3 °	64.2 °	-0.0 °	0.8 °	0.82	0.12	0.00	-0.01
<i>Nov</i>	215.6 °	83.4 °	-0.4 °	0.0 °	0.88	0.12	-0.01	-0.01
<i>Dec</i>	237.6 °	68.7 °	1.7 °	3.5 °	0.88	0.15	0.01	0.00

P7																																					
		Direction, average [°N]																																			
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	237	253	250	247	253	261	238	273	207	267	253	232	223	270	236	249	245	246	224	267	245	27	358	236	251	240	51	237	249	258	220	253	243	289	249	261	227
Jan	67	258	233	223	277	248	241	84	16	253	276	211	267	53	250	259	56	266	57	272	204	83	72	222	252	299	83	240	58	266	80	333	244	356	286	74	248
Feb	63	241	220	68	272	279	92	250	61	239	257	284	244	337	248	97	279	5	86	51	257	242	65	93	279	0	98	246	240	57	102	246	248	69	255	290	234
Mar	50	224	268	57	254	284	236	218	30	264	278	93	303	12	274	94	220	252	77	261	273	95	268	261	281	62	254	221	92	244	76	254	236	260	93	285	15
Apr	301	93	256	263	293	52	19	329	338	32	15	271	98	280	56	24	58	202	228	296	10	59	264	281	211	224	316	291	227	290	346	81	291	320	344	47	222
May	62	276	312	333	245	279	38	311	355	237	295	62	230	301	57	216	309	236	240	260	0	281	335	230	256	56	54	239	9	258	58	276	297	63	260	241	89
Jun	330	247	214	251	290	277	270	299	260	60	255	200	270	269	298	263	258	296	266	282	257	329	282	296	244	236	325	261	201	263	268	237	61	256	261	277	236
Jul	207	293	281	251	269	266	298	272	272	324	257	264	267	245	244	290	275	253	262	298	248	237	241	64	250	287	222	257	250	263	272	291	259	272	274	241	248
Aug	233	262	251	269	226	276	272	310	51	94	268	50	291	249	203	294	235	250	52	265	252	228	242	236	237	243	220	52	264	234	246	222	228	319	274	275	223
Sep	264	263	323	252	206	57	291	27	7	279	66	60	50	31	284	231	254	223	225	287	210	262	293	233	257	205	322	84	238	57	250	268	271	5	71	215	230
Oct	165	260	168	163	11	139	221	233	214	310	248	240	98	212	62	265	58	90	222	261	240	335	87	237	240	83	86	138	46	264	275	238	80	229	229	342	89
Nov	297	241	353	215	226	137	248	347	250	55	154	228	137	298	85	166	288	215	238	316	281	63	22	215	201	262	50	244	302	259	142	97	232	269	166	57	268
Dec	284	251	217	276	215	233	224	68	286	53	258	259	204	3	106	282	251	9	238	212	252	98	8	248	83	231	241	222	263	260	251	222	151	91	204	293	246

P7																																					
	H _{ms} average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	0.71	0.73	0.76	0.69	0.7	0.72	0.73	0.74	0.63	0.67	0.74	0.7	0.73	0.7	0.67	0.64	0.69	0.7	0.65	0.74	0.72	0.67	0.67	0.72	0.73	0.67	0.72	0.78	0.68	0.75	0.67	0.71	0.77	0.68	0.71	0.72	0.69
Jan	0.85	106	106	0.8	0.88	127	0.9	109	0.64	0.55	0.81	0.82	101	0.7	0.89	0.88	0.82	12	0.76	121	12	0.8	0.82	0.69	0.92	0.75	118	104	0.85	0.82	0.85	0.95	0.99	0.73	12	0.88	0.97
Feb	1	11	111	0.87	0.89	0.75	0.71	104	0.81	115	103	0.83	0.93	0.92	103	0.45	0.88	101	0.66	0.93	101	0.53	0.98	0.95	0.78	0.96	0.91	0.88	0.73	0.79	0.84	0.81	11	0.8	1	0.92	0.84
Mar	0.77	0.85	111	0.57	0.8	0.76	0.95	106	0.58	0.88	0.82	0.71	0.83	0.86	0.82	0.86	0.64	0.52	0.58	0.71	0.92	0.64	0.85	0.71	0.78	0.71	0.77	0.69	0.46	0.71	0.65	0.62	0.78	0.7	0.52	0.74	0.64
Apr	0.57	0.65	0.65	0.67	0.59	0.52	0.64	0.73	0.39	0.84	0.96	0.6	0.46	0.55	0.44	0.64	0.5	0.48	0.52	0.66	0.41	0.38	0.6	0.54	0.55	0.59	0.57	0.68	0.54	0.76	0.59	0.5	0.75	0.71	0.95	0.56	0.64
May	0.34	0.6	0.44	0.6	0.48	0.42	0.48	0.52	0.38	0.52	0.48	0.52	0.35	0.49	0.49	0.49	0.62	0.5	0.6	0.59	0.27	0.38	0.52	0.64	0.35	0.53	0.42	0.73	0.51	0.49	0.33	0.66	0.64	0.47	0.81	0.49	0.45
Jun	0.34	0.41	0.38	0.6	0.41	0.52	0.78	0.55	0.38	0.44	0.62	0.52	0.64	0.58	0.61	0.63	0.58	0.54	0.48	0.44	0.66	0.47	0.47	0.53	0.55	0.59	0.42	0.66	0.48	0.73	0.49	0.52	0.4	0.45	0.49	0.31	0.59
Jul	0.59	0.44	0.66	0.36	0.96	0.6	0.29	0.46	0.63	0.36	0.58	0.43	0.45	0.4	0.48	0.41	0.44	0.42	0.32	0.67	0.36	0.54	0.5	0.45	0.49	0.49	0.42	0.63	0.49	0.95	0.37	0.54	0.81	0.46	0.62	0.7	0.56
Aug	0.59	0.65	0.59	0.56	0.71	0.6	0.57	0.48	0.53	0.25	0.72	0.39	0.56	0.64	0.38	0.54	0.52	0.52	0.38	0.64	0.62	0.64	0.62	0.54	0.44	0.53	0.71	0.55	0.73	0.59	0.57	0.52	0.44	0.67	0.4	0.56	0.64
Sep	0.81	0.55	0.81	0.68	0.58	0.74	0.85	0.75	0.59	0.84	0.54	0.63	0.77	0.72	0.53	0.67	0.79	0.64	0.72	0.91	0.59	0.8	0.75	0.79	1	0.5	0.52	0.74	0.52	0.59	0.93	0.87	0.78	0.57	0.69	0.66	0.51
Oct	0.77	0.8	0.81	0.68	0.81	0.64	0.79	0.79	0.78	0.87	105	0.92	0.85	0.91	0.89	0.67	0.92	0.75	0.68	0.53	0.97	0.85	0.79	0.94	0.79	0.82	0.94	0.7	0.94	111	0.91	0.66	0.76	0.89	0.76	107	0.75
Nov	0.83	0.83	0.83	0.95	0.9	0.72	0.87	0.9	104	0.75	0.72	0.86	0.95	108	0.77	0.75	0.89	0.9	107	0.95	0.95	103	106	0.69	0.93	0.86	0.87	0.9	105	0.9	0.7	0.78	0.92	0.85	105	0.83	0.75
Dec	104	0.81	0.91	0.93	0.79	107	0.93	0.57	0.69	0.81	0.95	113	0.8	0.83	0.75	103	0.9	1	103	0.65	0.64	0.74	0.74	118	0.95	112	0.87	112	0.88	1	0.83	0.88	0.9	0.83	0.78	0.94	0.91

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	167.9 °	25.0 °	-0.1 °	-0.2 °	0.55	0.02	0.00	-0.01
<i>Jan</i>	171.0 °	54.8 °	0.2 °	-1.4 °	0.70	0.13	0.00	-0.01
<i>Feb</i>	158.2 °	79.2 °	-0.3 °	-0.9 °	0.68	0.12	0.00	-0.01
<i>Mar</i>	158.1 °	55.5 °	-0.6 °	-0.3 °	0.58	0.09	0.00	0.00
<i>Apr</i>	104.3 °	78.9 °	-0.9 °	-1.2 °	0.48	0.07	0.00	-0.01
<i>May</i>	146.9 °	87.9 °	0.5 °	2.4 °	0.42	0.06	0.00	0.00
<i>Jun</i>	201.3 °	69.7 °	0.6 °	-0.2 °	0.39	0.07	0.00	0.00
<i>Jul</i>	208.3 °	56.0 °	3.2 °	4.4 °	0.37	0.07	0.00	-0.01
<i>Aug</i>	188.2 °	62.0 °	-0.1 °	0.6 °	0.42	0.08	-0.01	-0.01
<i>Sep</i>	166.7 °	63.8 °	2.3 °	3.3 °	0.55	0.10	0.00	-0.01
<i>Oct</i>	161.9 °	43.1 °	0.8 °	-0.2 °	0.66	0.11	0.00	-0.01
<i>Nov</i>	151.5 °	64.7 °	-0.4 °	-0.7 °	0.70	0.10	-0.01	-0.01
<i>Dec</i>	168.4 °	58.4 °	0.8 °	0.9 °	0.67	0.09	-0.01	-0.01

			P8																																			
			Direction, average [°N]																																			
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	56	56	59	61	62	58	58	55	59	67	59	59	63	50	54	50	64	51	55	54	64	59	49	61	71	61	65	69	58	75	51	55	66	54	61	62	62	
Jan	61	238	205	67	262	27	62	63	16	226	57	76	244	61	222	66	55	242	57	248	67	57	66	73	63	66	67	22	52	69	58	39	23	65	266	61	60	
Feb	56	207	58	60	252	281	99	225	88	210	240	288	210	51	209	52	296	57	97	18	232	66	61	65	269	44	61	52	68	57	62	88	214	90	218	256	65	
Mar	10	60	252	53	79	287	205	71	67	210	63	131	332	66	244	67	51	53	66	57	66	58	55	223	254	60	62	62	60	75	64	217	79	219	59	63	51	
Apr	340	66	66	54	90	52	111	10	58	326	17	247	10	10	15	73	52	66	60	299	57	59	52	280	52	74	87	286	52	278	97	95	256	341	17	90	54	
May	57	264	342	339	203	57	58	20	78	52	335	59	68	316	111	64	351	68	64	223	77	225	29	69	203	102	82	205	59	502	111	269	297	55	206	59	67	
Jun	28	221	54	230	61	274	258	92	240	51	224	63	239	298	60	230	233	293	239	223	233	4	272	221	203	67	354	243	68	237	259	61	51	69	201	62	207	
Jul	65	309	244	227	294	281	56	252	263	25	244	245	236	205	226	75	260	212	232	231	68	69	74	68	233	297	65	240	226	62	60	241	243	202	268	219	204	
Aug	212	257	69	254	66	270	56	346	16	54	257	59	222	63	16	292	57	209	97	68	214	65	65	74	208	61	62	51	236	64	209	66	68	5	75	265	61	
Sep	236	67	48	209	64	79	66	89	70	267	55	51	19	72	83	68	220	57	67	308	54	227	66	63	223	67	76	55	68	54	220	217	241	68	65	71	59	
Oct	60	209	54	52	72	100	67	65	67	346	72	68	68	69	83	84	55	69	75	62	205	73	60	66	63	66	65	16	74	67	59	67	52	65	69	66	66	
Nov	327	55	45	177	63	54	62	56	62	59	61	66	64	303	61	63	268	94	95	5	230	64	59	66	66	225	53	203	63	215	57	66	67	64	52	90	220	
Dec	205	50	60	206	177	69	64	63	60	18	68	68	51	54	58	251	221	78	206	62	61	63	45	217	52	200	68	61	241	224	59	63	57	68	63	52	64	

P8																																						
		Hm0, average [m]																																				
		1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual		0.56	0.53	0.56	0.54	0.52	0.58	0.59	0.57	0.56	0.52	0.58	0.54	0.58	0.55	0.57	0.5	0.54	0.55	0.51	0.55	0.54	0.56	0.57	0.54	0.56	0.54	0.61	0.59	0.54	0.56	0.55	0.56	0.56	0.53	0.54	0.56	0.55
Jan		0.73	0.63	0.73	0.6	0.57	0.64	0.59	0.62	0.74	0.39	0.61	0.66	0.68	0.63	0.61	0.67	0.73	0.75	0.68	0.79	0.93	0.63	0.63	0.52	0.72	0.64	1.07	0.68	0.77	0.57	0.67	0.7	0.66	0.62	0.72	0.66	0.76
Feb		0.62	0.74	0.62	0.63	0.58	0.51	0.78	0.69	0.78	0.61	0.67	0.55	0.67	0.71	0.72	0.41	0.49	0.65	0.57	0.68	0.68	0.53	0.57	0.66	0.57	0.52	0.77	0.68	0.55	0.73	0.73	0.58	0.74	0.72	0.67	0.61	0.66
Mar		0.67	0.63	0.68	0.55	0.61	0.54	0.7	0.62	0.65	0.6	0.59	0.59	0.57	0.56	0.59	0.46	0.59	0.5	0.54	0.59	0.69	0.56	0.42	0.52	0.56	0.74	0.58	0.57	0.4	0.52	0.69	0.57	0.61	0.5	0.45	0.52	0.62
Apr		0.43	0.63	0.5	0.58	0.46	0.52	0.55	0.52	0.32	0.59	0.59	0.46	0.44	0.47	0.42	0.62	0.5	0.43	0.4	0.45	0.36	0.38	0.5	0.4	0.48	0.48	0.5	0.43	0.47	0.52	0.49	0.53	0.49	0.49	0.44	0.53	0.57
May		0.38	0.41	0.29	0.45	0.35	0.41	0.48	0.41	0.49	0.47	0.41	0.47	0.41	0.36	0.48	0.36	0.5	0.36	0.52	0.4	0.28	0.44	0.44	0.51	0.43	0.48	0.35	0.51	0.46	0.44	0.33	0.48	0.43	0.38	0.42	0.44	0.49
Jun		0.27	0.28	0.32	0.44	0.35	0.37	0.5	0.45	0.28	0.45	0.45	0.4	0.44	0.42	0.48	0.45	0.44	0.42	0.36	0.32	0.45	0.41	0.53	0.38	0.45	0.48	0.31	0.44	0.44	0.54	0.36	0.43	0.38	0.32	0.34	0.27	0.43
Jul		0.48	0.34	0.46	0.3	0.4	0.41	0.26	0.34	0.43	0.32	0.43	0.28	0.33	0.29	0.37	0.33	0.32	0.34	0.22	0.47	0.3	0.38	0.38	0.39	0.35	0.33	0.35	0.46	0.32	0.44	0.29	0.4	0.5	0.38	0.43	0.45	0.4
Aug		0.44	0.47	0.41	0.38	0.53	0.43	0.47	0.37	0.35	0.23	0.5	0.33	0.41	0.48	0.42	0.34	0.42	0.38	0.32	0.5	0.45	0.49	0.46	0.46	0.31	0.39	0.54	0.54	0.48	0.42	0.42	0.38	0.34	0.52	0.3	0.39	0.44
Sep		0.55	0.39	0.6	0.46	0.49	0.78	0.66	0.67	0.56	0.52	0.54	0.58	0.61	0.65	0.44	0.49	0.53	0.46	0.58	0.65	0.51	0.54	0.63	0.53	0.65	0.44	0.48	0.62	0.38	0.53	0.61	0.6	0.52	0.46	0.62	0.49	0.45
Oct		0.67	0.58	0.7	0.62	0.65	0.55	0.64	0.52	0.58	0.61	0.64	0.71	0.71	0.71	0.78	0.54	0.8	0.59	0.51	0.39	0.63	0.72	0.69	0.68	0.6	0.68	0.78	0.69	0.69	0.75	0.67	0.5	0.67	0.63	0.52	0.63	0.6
Nov		0.59	0.66	0.53	0.72	0.68	0.75	0.66	0.75	0.82	0.76	0.67	0.67	0.84	0.7	0.74	0.65	0.61	0.66	0.77	0.7	0.71	0.6	0.88	0.58	0.7	0.57	0.87	0.61	0.78	0.59	0.68	0.73	0.63	0.63	0.95	0.73	0.49
Dec		0.74	0.63	0.71	0.8	0.8	0.82	0.67	0.54	0.54	0.8	0.7	0.78	0.65	0.65	0.8	0.72	0.56	0.62	0.66	0.49	0.5	0.71	0.66	0.73	0.68	0.75	0.8	0.79	0.57	0.68	0.68	0.66	0.8	0.68	0.64	0.75	0.64

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	167.9°	25.0°	-0.1°	-0.2°	0.59	0.02	-0.01	-0.02
<i>Jan</i>	171.0°	54.8°	0.2°	-1.4°	0.74	0.13	-0.01	-0.02
<i>Feb</i>	158.2°	79.2°	-0.3°	-0.9°	0.72	0.11	-0.01	-0.01
<i>Mar</i>	158.1°	55.5°	-0.6°	-0.3°	0.62	0.10	0.00	-0.01
<i>Apr</i>	104.3°	78.9°	-0.9°	-1.2°	0.52	0.08	-0.01	-0.03
<i>May</i>	146.9°	87.9°	0.5°	2.4°	0.46	0.07	-0.01	-0.01
<i>Jun</i>	201.3°	69.7°	0.6°	-0.2°	0.45	0.08	-0.01	-0.02
<i>Jul</i>	208.3°	56.0°	3.2°	4.4°	0.42	0.08	0.00	-0.01
<i>Aug</i>	188.2°	62.0°	-0.1°	0.6°	0.47	0.08	0.00	-0.01
<i>Sep</i>	166.7°	63.8°	2.3°	3.3°	0.59	0.10	0.00	-0.01
<i>Oct</i>	161.9°	43.1°	0.8°	-0.2°	0.69	0.11	-0.01	-0.02
<i>Nov</i>	151.5°	64.7°	-0.4°	-0.7°	0.73	0.09	-0.01	-0.02
<i>Dec</i>	168.4°	58.4°	0.8°	0.9°	0.72	0.10	0.00	-0.01

P9																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	211	260	247	243	248	238	209	305	92	277	251	200	165	24	100	233	234	237	89	278	234	149	58	221	249	173	143	223	188	259	135	230	230	9	226	225	175
Jan	65	260	227	220	284	240	240	60	102	261	296	202	271	160	249	263	154	268	55	271	183	189	186	202	218	186	124	237	150	265	168	351	239	45	288	204	183
Feb	57	233	214	81	283	301	96	249	76	234	263	296	239	0	242	154	297	48	88	102	258	74	89	145	295	32	165	261	239	123	99	237	240	82	251	302	206
Mar	152	211	274	131	236	306	237	208	79	258	299	133	312	100	287	149	99	89	90	253	262	115	294	269	296	75	240	150	141	237	90	254	209	273	135	241	120
Apr	319	85	270	82	329	19	92	344	34	319	114	283	93	75	112	54	114	154	231	305	95	122	293	296	114	118	62	297	221	294	90	90	298	328	342	69	121
May	119	292	325	342	239	75	91	341	13	110	322	127	197	314	131	166	322	231	164	289	17	270	355	204	267	13	13	241	40	69	108	297	306	148	268	115	105
Jun	1	279	142	275	38	295	280	322	284	164	261	163	278	351	256	274	269	307	290	294	263	332	296	269	260	252	324	274	167	281	286	140	109	249	276	320	248
Jul	195	311	281	287	290	279	89	283	291	303	270	277	286	239	270	323	287	280	288	267	236	240	246	45	261	303	117	269	260	291	315	297	271	305	290	240	248
Aug	253	293	265	291	216	292	304	332	119	121	283	118	274	296	154	301	216	267	66	292	298	224	261	216	252	237	219	118	270	240	258	119	171	337	340	284	207
Sep	270	286	340	261	166	77	29	74	56	285	122	129	111	12	32	226	253	219	160	302	154	270	66	224	261	193	53	140	227	125	249	263	280	73	183	162	111
Oct	163	268	135	130	12	98	167	222	189	325	214	164	162	162	75	329	137	161	207	287	232	32	141	224	234	160	146	107	71	292	309	224	133	210	220	74	166
Nov	312	102	21	189	220	130	234	30	200	125	87	160	166	306	99	145	302	192	232	328	291	148	51	195	191	272	119	235	348	256	124	109	221	267	132	183	260
Dec	293	159	119	287	150	119	217	81	70	114	260	235	168	28	133	291	248	50	231	200	150	99	31	240	96	222	228	212	266	258	122	209	138	81	157	172	210

P9																																					
	Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	0.6	0.58	0.61	0.59	0.57	0.62	0.64	0.62	0.58	0.57	0.63	0.58	0.61	0.58	0.61	0.54	0.56	0.56	0.54	0.61	0.58	0.58	0.4	0.58	0.61	0.58	0.63	0.64	0.58	0.62	0.58	0.58	0.6	0.58	0.58	0.6	0.58
Jan	0.71	0.71	0.76	0.65	0.68	0.53	0.76	0.85	0.72	0.42	0.63	0.68	0.76	0.63	0.68	0.71	0.71	0.55	0.68	0.52	0.56	0.63	0.5	0.50	0.79	0.68	118	0.75	0.78	0.61	0.68	0.78	0.71	0.65	0.63	0.68	0.63
Feb	0.84	0.81	0.86	0.83	0.86	0.58	0.77	0.79	0.78	0.86	0.77	0.63	0.74	0.76	0.8	0.4	0.56	0.87	0.58	0.9	0.77	0.58	0.58	0.89	0.66	0.54	0.73	0.72	0.61	0.76	0.72	0.63	0.81	0.72	0.76	0.7	0.71
Mar	0.69	0.67	0.8	0.56	0.67	0.59	0.77	0.87	0.67	0.67	0.65	0.59	0.65	0.36	0.65	0.43	0.63	0.54	0.52	0.63	0.76	0.59	0.43	0.58	0.65	0.76	0.61	0.6	0.4	0.57	0.69	0.64	0.63	0.54	0.46	0.56	0.61
Apr	0.49	0.66	0.55	0.57	0.5	0.52	0.56	0.61	0.33	0.7	0.57	0.51	0.45	0.49	0.43	0.64	0.51	0.43	0.44	0.53	0.35	0.39	0.56	0.46	0.52	0.51	0.53	0.53	0.51	0.62	0.52	0.54	0.56	0.57	0.47	0.56	0.57
May	0.39	0.47	0.35	0.51	0.37	0.44	0.51	0.48	0.52	0.49	0.47	0.5	0.43	0.45	0.48	0.38	0.58	0.4	0.57	0.44	0.28	0.48	0.53	0.53	0.48	0.48	0.38	0.58	0.47	0.48	0.34	0.56	0.53	0.39	0.46	0.49	0.5
Jun	0.3	0.32	0.35	0.49	0.41	0.47	0.6	0.49	0.33	0.49	0.5	0.43	0.51	0.47	0.58	0.52	0.53	0.5	0.39	0.36	0.52	0.5	0.39	0.43	0.51	0.52	0.37	0.53	0.46	0.64	0.43	0.45	0.38	0.34	0.38	0.3	0.47
Jul	0.47	0.43	0.55	0.35	0.45	0.49	0.28	0.39	0.5	0.37	0.5	0.33	0.38	0.32	0.44	0.37	0.38	0.39	0.24	0.55	0.33	0.42	0.4	0.44	0.36	0.4	0.36	0.53	0.37	0.5	0.35	0.46	0.59	0.42	0.52	0.5	0.44
Aug	0.49	0.55	0.47	0.47	0.36	0.49	0.12	0.43	0.36	0.23	0.39	0.37	0.48	0.33	0.43	0.42	0.44	0.44	0.35	0.38	0.5	0.33	0.49	0.5	0.35	0.43	0.58	0.35	0.35	0.48	0.49	0.38	0.34	0.38	0.32	0.44	0.47
Sep	0.6	0.43	0.67	0.54	0.5	0.79	0.69	0.7	0.6	0.63	0.54	0.57	0.8	0.66	0.49	0.53	0.61	0.49	0.63	0.73	0.52	0.62	0.67	0.58	0.73	0.47	0.52	0.65	0.41	0.53	0.66	0.68	0.55	0.48	0.65	0.51	0.47
Oct	0.71	0.65	0.7	0.64	0.67	0.57	0.71	0.53	0.6	0.7	0.94	0.77	0.68	0.72	0.8	0.6	0.8	0.59	0.64	0.43	0.69	0.78	0.7	0.74	0.65	0.69	0.76	0.71	0.8	0.63	0.74	0.53	0.67	0.66	0.55	0.8	0.62
Nov	0.67	0.68	0.58	0.75	0.75	0.74	0.74	0.78	0.86	0.73	0.87	0.68	0.79	0.81	0.74	0.65	0.7	0.69	0.82	0.79	0.8	0.87	0.89	0.59	0.71	0.62	0.85	0.67	0.82	0.66	0.66	0.71	0.67	0.67	0.91	0.75	0.56
Dec	0.88	0.68	0.74	0.7	0.62	0.86	0.72	0.57	0.26	0.8	0.72	0.85	0.64	0.7	0.8	0.79	0.62	0.87	0.75	0.52	0.53	0.7	0.7	0.81	0.85	0.79	0.66	0.82	0.65	0.77	0.74	0.7	0.77	0.71	0.63	0.8	0.67

P10

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
Annual	231.5 °	13.6 °	1.2 °	1.9 °	0.74	0.04	-0.01	-0.02
Jan	227.3 °	45.2 °	-0.1 °	-0.4 °	0.95	0.19	0.00	-0.01
Feb	227.1 °	68.5 °	1.2 °	1.9 °	0.90	0.20	-0.01	-0.03
Mar	222.5 °	54.1 °	0.5 °	1.3 °	0.75	0.16	-0.01	-0.02
Apr	264.1 °	64.3 °	0.2 °	0.1 °	0.60	0.11	0.00	-0.03
May	248.0 °	59.0 °	0.1 °	0.3 °	0.55	0.10	0.00	-0.02
Jun	256.2 °	26.8 °	2.2 °	3.8 °	0.57	0.13	-0.01	-0.02
Jul	254.8 °	19.5 °	-0.1 °	-0.7 °	0.55	0.13	0.00	-0.01
Aug	240.7 °	36.5 °	-0.7 °	0.2 °	0.60	0.13	-0.01	-0.02
Sep	235.7 °	53.1 °	0.4 °	-0.3 °	0.73	0.14	0.00	-0.01
Oct	216.9 °	51.2 °	1.2 °	1.9 °	0.86	0.13	-0.01	-0.03
Nov	215.9 °	68.5 °	-2.3 °	-0.9 °	0.90	0.12	-0.01	-0.04
Dec	228.7 °	56.7 °	1.3 °	1.5 °	0.92	0.17	-0.01	-0.01

P10																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	237	250	246	244	247	247	237	239	220	238	247	231	227	230	230	244	243	245	228	259	241	226	273	236	246	238	24	236	242	251	226	246	240	262	245	245	227
Jan	81	252	231	227	272	241	241	210	18	253	250	220	260	84	246	247	88	257	83	260	211	203	89	228	244	235	89	236	89	258	201	309	241	292	278	220	238
Feb	216	236	222	87	266	274	83	246	73	235	254	279	241	306	240	85	266	9	96	96	251	222	89	82	273	346	87	245	241	82	80	244	240	89	248	278	231
Mar	94	227	263	86	246	275	237	227	71	256	269	80	289	96	266	200	232	240	84	245	252	207	266	258	279	89	244	222	84	241	92	251	234	255	96	249	81
Apr	286	82	252	229	252	88	207	311	32	301	59	264	88	232	87	11	58	209	234	291	57	81	262	274	234	221	259	283	232	277	86	93	283	306	303	234	216
May	88	274	305	317	242	249	85	300	278	234	282	89	226	294	86	221	291	238	240	257	351	263	318	230	250	82	358	242	360	220	82	275	290	212	260	249	18
Jun	311	257	222	251	291	281	268	284	262	81	256	210	265	275	253	261	258	287	263	271	258	307	281	282	252	249	304	263	220	261	270	225	209	256	282	287	247
Jul	212	280	278	295	270	287	272	268	275	18	257	265	278	243	230	274	268	263	272	257	243	240	248	252	247	288	218	295	252	263	231	285	281	284	274	240	245
Aug	244	263	256	274	228	267	263	312	84	80	269	83	261	248	84	290	235	256	259	264	250	235	246	238	245	247	230	86	260	243	250	225	228	300	285	268	229
Sep	255	260	298	255	224	73	240	206	354	276	81	85	57	48	279	235	251	231	226	272	215	258	246	235	253	223	301	207	242	89	247	258	262	271	88	217	236
Oct	205	253	84	81	358	82	226	233	222	303	238	235	88	210	73	277	98	82	227	261	238	304	83	236	242	202	93	86	82	268	262	236	89	230	232	285	210
Nov	280	243	331	219	234	10	248	331	236	85	89	224	99	287	10	81	281	216	237	299	262	95	21	209	210	255	86	238	262	252	81	87	232	254	86	98	259
Dec	278	243	218	269	216	224	228	83	255	89	245	242	205	329	10	266	246	352	236	218	245	83	355	240	84	232	236	221	259	250	242	225	83	181	208	256	233

P10																																					
	Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	0.75	0.76	0.8	0.73	0.73	0.75	0.78	0.77	0.65	0.7	0.78	0.73	0.77	0.71	0.71	0.67	0.74	0.72	0.68	0.78	0.76	0.71	0.68	0.77	0.77	0.71	0.75	0.83	0.71	0.8	0.7	0.75	0.81	0.69	0.74	0.75	0.72
Jan	0.88	109	115	0.9	0.92	135	0.99	107	0.83	0.57	0.8	0.88	108	0.73	0.98	0.91	0.83	1.9	0.76	13	126	0.79	0.76	0.72	0.96	0.77	123	11	0.88	0.81	0.88	0.95	107	0.75	1.8	0.92	104
Feb	102	116	125	0.85	0.91	0.78	0.89	116	0.79	122	111	0.86	102	0.89	114	0.45	0.73	0.95	0.81	0.95	107	0.8	0.57	1	0.82	0.94	0.93	0.89	0.82	0.84	0.8	0.85	118	0.77	109	0.92	0.9
Mar	0.77	0.9	1.0	0.82	0.84	0.74	103	11	0.89	0.87	0.84	0.7	0.84	0.86	0.85	0.8	0.72	0.57	0.54	0.75	0.97	0.67	0.6	0.76	0.81	0.71	0.77	0.72	0.46	0.75	0.62	0.91	0.82	0.72	0.52	0.76	0.66
Apr	0.61	0.67	0.68	0.63	0.62	0.63	0.63	0.76	0.38	0.86	0.55	0.66	0.49	0.86	0.44	0.67	0.51	0.5	0.57	0.67	0.39	0.4	0.65	0.96	0.6	0.64	0.57	0.72	0.62	0.81	0.61	0.5	0.74	0.73	0.56	0.67	0.84
May	0.38	0.61	0.44	0.6	0.48	0.45	0.5	0.58	0.58	0.55	0.52	0.56	0.57	0.54	0.51	0.51	0.66	0.52	0.66	0.59	0.28	0.81	0.59	0.67	0.59	0.53	0.43	0.8	0.47	0.52	0.35	0.72	0.7	0.48	0.63	0.54	0.46
Jun	0.34	0.43	0.41	0.67	0.42	0.6	0.82	0.59	0.44	0.51	0.66	0.56	0.69	0.6	0.7	0.67	0.68	0.62	0.49	0.43	0.72	0.59	0.49	0.57	0.67	0.68	0.47	0.73	0.53	0.83	0.57	0.58	0.44	0.42	0.51	0.35	0.66
Jul	0.61	0.52	0.73	0.44	0.86	0.7	0.3	0.5	0.87	0.4	0.87	0.46	0.48	0.44	0.96	0.43	0.5	0.48	0.32	0.76	0.41	0.8	0.52	0.51	0.56	0.52	0.4	0.71	0.56	0.62	0.41	0.59	0.88	0.51	0.71	0.78	0.62
Aug	0.68	0.72	0.61	0.62	0.78	0.65	0.61	0.82	0.86	0.24	0.81	0.4	0.63	0.89	0.41	0.8	0.54	0.61	0.39	0.88	0.7	0.7	0.64	0.64	0.47	0.58	0.82	0.8	0.79	0.67	0.67	0.57	0.43	0.73	0.39	0.8	0.66
Sep	0.83	0.84	0.81	0.73	0.62	0.73	0.87	0.76	0.6	0.91	0.58	0.68	0.78	0.69	0.55	0.69	0.87	0.64	0.78	0.95	0.81	0.86	0.78	0.85	106	0.56	0.57	0.78	0.54	0.61	0.97	0.94	0.75	0.57	0.71	0.69	0.55
Oct	0.84	0.87	0.82	0.73	0.75	0.67	0.9	0.78	0.82	0.88	12	0.95	0.85	0.94	0.85	0.72	0.93	0.74	0.76	0.55	107	0.88	0.82	0.99	0.85	0.87	0.95	0.73	0.89	1.2	0.95	0.72	0.83	0.94	0.81	109	0.78
Nov	0.85	0.82	0.87	101	1	0.72	0.94	0.88	108	0.74	0.71	0.86	0.98	108	0.75	0.78	0.93	0.93	1.0	0.98	102	106	0.97	0.73	0.94	0.89	0.88	0.96	105	0.95	0.71	0.77	0.97	0.86	103	0.82	0.77
Dec	1.0	0.85	0.94	0.97	0.84	1.1	104	0.99	0.88	0.66	0.94	1.07	0.81	0.81	0.73	105	0.94	0.96	1.0	0.73	0.63	0.71	0.75	123	0.92	1.07	0.94	1.07	0.94	1.08	0.88	0.98	0.9	0.82	0.8	0.97	0.89

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	218.0°	11.4°	-1.3°	0.4°	0.76	0.05	-0.01	-0.04
<i>Jan</i>	216.7°	38.6°	0.3°	3.2°	0.99	0.21	-0.01	-0.04
<i>Feb</i>	214.1°	60.6°	-0.8°	0.6°	0.93	0.22	-0.02	-0.05
<i>Mar</i>	210.1°	47.4°	-1.1°	0.4°	0.77	0.16	-0.01	-0.05
<i>Apr</i>	222.7°	60.3°	0.4°	0.9°	0.60	0.10	-0.01	-0.02
<i>May</i>	221.6°	60.9°	-1.4°	0.2°	0.54	0.10	-0.02	-0.04
<i>Jun</i>	240.7°	26.9°	-3.2°	-5.5°	0.58	0.14	-0.01	-0.04
<i>Jul</i>	243.0°	20.3°	0.4°	2.7°	0.56	0.14	-0.01	-0.03
<i>Aug</i>	229.2°	36.5°	-1.6°	-1.1°	0.62	0.14	0.00	-0.05
<i>Sep</i>	221.8°	49.1°	-0.5°	0.4°	0.75	0.15	-0.01	-0.03
<i>Oct</i>	209.4°	41.3°	-0.9°	-0.1°	0.89	0.14	-0.01	-0.05
<i>Nov</i>	206.5°	59.2°	-1.4°	-2.6°	0.92	0.13	-0.01	-0.03
<i>Dec</i>	217.0°	48.3°	-0.2°	0.7°	0.95	0.19	0.00	-0.04

P11																																						
	Direction, average [°N]																																					
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	227	241	235	233	237	232	227	242	86	243	235	222	229	232	235	230	232	231	239	246	232	238	222	228	236	225	81	228	228	240	238	234	231	240	233	231	238	
Jan	83	245	226	222	262	234	235	205	19	244	233	236	251	86	239	233	91	248	83	249	208	201	84	222	230	213	81	232	92	247	98	290	230	233	227	267	236	227
Feb	210	229	219	18	256	264	69	239	61	229	246	267	233	282	232	85	252	48	99	99	243	97	97	81	262	360	80	234	235	80	69	235	232	92	240	266	226	
Mar	96	222	254	86	235	264	232	219	87	240	250	89	274	92	253	94	238	205	64	229	237	98	253	248	268	81	235	209	86	231	97	241	226	243	92	235	82	
Apr	268	191	241	210	235	168	194	298	309	291	133	252	164	211	81	91	155	200	228	281	156	154	248	263	223	214	221	274	224	268	91	100	269	294	274	81	201	
May	83	267	296	307	233	223	161	289	259	235	267	87	217	287	82	236	282	233	225	246	47	254	316	224	242	82	86	237	39	98	87	268	283	207	249	204	194	
Jun	295	247	299	244	283	276	262	265	255	69	249	210	258	270	244	254	251	274	251	282	250	298	276	256	246	242	289	259	211	254	264	210	64	246	257	278	241	
Jul	207	285	287	290	281	281	244	260	268	192	251	257	268	236	246	258	258	258	268	253	236	237	240	230	242	282	203	250	247	257	287	278	258	259	268	233	248	
Aug	241	259	250	269	223	259	248	309	82	93	262	94	259	238	82	279	227	250	239	252	241	228	237	230	242	241	224	163	253	236	244	221	234	286	274	262	224	
Sep	247	243	278	249	219	85	225	94	65	269	87	84	97	79	265	229	243	225	238	259	202	249	223	229	245	236	267	202	233	87	240	251	248	229	85	238	291	
Oct	201	243	82	89	195	80	238	238	269	229	229	80	207	99	262	99	82	221	251	232	277	83	229	235	89	94	132	76	256	247	227	86	228	227	234	207		
Nov	266	221	320	235	229	10	237	330	223	168	97	235	83	274	121	82	269	210	231	281	247	98	49	203	206	244	97	231	239	242	83	156	227	242	89	132	247	
Dec	265	230	210	260	212	238	224	163	228	90	231	232	203	281	194	250	239	346	230	216	231	134	5	233	143	227	228	216	250	241	228	219	87	147	204	232	223	

P11																																					
	Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	0.77	0.78	0.82	0.75	0.75	0.78	0.81	0.78	0.66	0.71	0.81	0.75	0.79	0.71	0.73	0.68	0.76	0.74	0.7	0.75	0.75	0.73	0.68	0.8	0.75	0.72	0.76	0.66	0.73	0.82	0.72	0.77	0.82	0.69	0.74	0.77	0.74
Jan	0.94	1.2	1.24	0.97	0.94	1.45	1.05	1.09	0.87	0.96	0.81	0.93	1.2	0.77	1.03	0.94	0.87	1.25	0.79	1.38	1.34	0.82	0.77	0.73	1	0.8	1.32	1.17	0.95	0.81	0.91	0.93	1.1	0.77	1.14	0.97	1.09
Feb	1.08	1.24	1.37	0.87	0.92	0.77	0.73	1.24	0.81	1.3	1.16	0.85	1.08	0.87	1.23	0.48	0.74	0.94	0.8	0.98	1.3	0.83	0.99	1.07	0.84	0.92	0.99	0.9	0.86	0.91	0.82	0.87	1.25	0.78	1.15	0.91	0.94
Mar	0.78	0.94	1.16	0.64	0.87	0.73	1.09	1.15	0.6	0.87	0.83	0.7	0.83	0.58	0.87	0.59	0.76	0.6	0.54	0.78	0.99	0.68	0.61	0.77	0.8	0.73	0.78	0.75	0.46	0.76	0.64	0.94	0.85	0.71	0.91	0.76	0.69
Apr	0.6	0.68	0.7	0.63	0.63	0.55	0.64	0.74	0.36	0.84	0.98	0.68	0.49	0.96	0.45	0.67	0.52	0.52	0.59	0.65	0.37	0.42	0.65	0.96	0.6	0.65	0.57	0.7	0.65	0.8	0.62	0.5	0.7	0.68	0.53	0.56	0.64
May	0.41	0.39	0.42	0.35	0.49	0.46	0.51	0.57	0.36	0.36	0.38	0.38	0.32	0.32	0.32	0.32	0.85	0.53	0.69	0.58	0.27	0.81	0.58	0.68	0.59	0.53	0.42	0.82	0.46	0.53	0.35	0.71	0.66	0.48	0.81	0.54	0.48
Jun	0.33	0.42	0.41	0.69	0.42	0.6	0.84	0.59	0.45	0.52	0.67	0.59	0.7	0.58	0.73	0.68	0.71	0.62	0.48	0.43	0.74	0.59	0.46	0.57	0.7	0.69	0.45	0.74	0.54	0.86	0.57	0.56	0.45	0.41	0.48	0.35	0.68
Jul	0.64	0.53	0.73	0.46	0.57	0.72	0.3	0.5	0.87	0.39	0.71	0.46	0.45	0.45	0.39	0.43	0.51	0.49	0.3	0.78	0.45	0.62	0.51	0.53	0.58	0.49	0.4	0.73	0.58	0.63	0.4	0.59	0.89	0.5	0.71	0.82	0.63
Aug	0.71	0.73	0.61	0.62	0.82	0.65	0.62	0.5	0.39	0.24	0.82	0.41	0.64	0.71	0.44	0.54	0.36	0.62	0.4	0.68	0.74	0.73	0.65	0.69	0.48	0.59	0.87	0.64	0.76	0.69	0.69	0.59	0.42	0.72	0.36	0.81	0.67
Sep	0.86	0.53	0.8	0.74	0.64	0.76	0.89	0.78	0.61	0.92	0.61	0.71	0.81	0.69	0.35	0.69	0.91	0.64	0.8	0.97	0.65	0.88	0.8	0.89	1.09	0.58	0.59	0.81	0.54	0.63	1	0.96	0.73	0.56	0.72	0.7	0.57
Oct	0.9	0.9	0.85	0.77	0.74	0.68	0.94	0.79	0.84	0.85	1.3	0.99	0.92	0.98	0.88	0.73	0.95	0.76	0.62	0.55	1.3	0.9	0.85	1.02	0.89	0.93	0.99	0.75	0.9	1.2	0.98	0.74	0.67	0.97	0.83	1.2	0.79
Nov	0.84	0.81	0.87	1.08	1.06	0.76	0.97	0.88	1.5	0.77	0.71	0.89	1.04	1.04	0.77	0.82	0.92	0.96	1.7	0.97	1.05	1.4	0.95	0.78	0.96	0.89	0.93	1.02	1.05	0.98	0.74	0.78	1.01	0.87	1.07	0.84	0.77
Dec	1.5	0.87	0.97	0.97	0.89	1.6	1.1	0.8	0.68	0.7	0.95	1.24	0.84	0.79	0.77	1.07	0.97	0.95	1.21	0.78	0.64	0.73	0.74	1.32	0.94	1.24	1.01	1.27	0.96	1.6	0.91	1.03	0.93	0.82	0.82	1	0.82

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	292.4 °	9.3 °	-0.3 °	-1.5 °	0.65	0.05	0.00	-0.02
<i>Jan</i>	277.5 °	42.0 °	0.1 °	-0.1 °	0.83	0.23	0.00	-0.03
<i>Feb</i>	286.9 °	47.9 °	-0.3 °	-1.0 °	0.80	0.21	0.00	-0.03
<i>Mar</i>	291.0 °	38.7 °	-0.4 °	-1.8 °	0.66	0.18	-0.01	-0.03
<i>Apr</i>	325.7 °	48.0 °	1.0 °	1.6 °	0.54	0.14	-0.01	-0.01
<i>May</i>	318.2 °	30.3 °	-0.3 °	-1.4 °	0.50	0.11	0.00	-0.03
<i>Jun</i>	302.9 °	13.4 °	0.9 °	-1.3 °	0.54	0.13	-0.01	-0.03
<i>Jul</i>	299.5 °	16.2 °	-0.9 °	-2.5 °	0.53	0.14	0.00	-0.03
<i>Aug</i>	294.2 °	30.6 °	0.2 °	-0.7 °	0.55	0.14	0.00	-0.02
<i>Sep</i>	296.9 °	35.1 °	-0.4 °	-1.8 °	0.64	0.16	0.00	-0.03
<i>Oct</i>	281.1 °	33.4 °	-0.7 °	-2.4 °	0.72	0.15	0.00	-0.01
<i>Nov</i>	288.5 °	62.0 °	-1.2 °	0.1 °	0.74	0.16	0.00	-0.01
<i>Dec</i>	279.4 °	40.4 °	0.4 °	-0.4 °	0.80	0.19	0.01	-0.01

P12																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	281	282	277	281	279	286	278	295	310	290	282	275	275	295	296	285	280	286	274	287	274	285	312	271	283	284	287	273	286	283	290	283	274	298	281	284	278
Jan	220	268	254	257	284	262	271	279	84	273	286	256	282	220	268	273	224	275	87	276	252	258	40	265	290	297	18	264	240	264	247	314	263	317	294	254	284
Feb	272	261	247	344	287	293	68	266	97	260	267	293	263	315	264	219	285	351	340	65	270	322	1	249	295	335	223	286	266	276	326	275	262	16	270	293	274
Mar	307	281	278	229	280	286	264	270	15	281	296	269	300	61	285	276	280	301	334	277	288	301	292	282	299	36	276	266	336	276	51	272	274	278	269	282	295
Apr	306	29	287	297	288	276	293	318	339	305	86	282	326	291	261	336	73	260	263	303	38	109	297	293	296	261	297	295	277	291	310	60	302	309	292	328	271
May	324	293	311	316	265	307	322	310	299	298	300	290	263	304	336	259	306	268	283	281	340	285	323	260	278	316	325	269	338	331	295	292	302	278	282	319	79
Jun	319	294	276	277	323	294	284	304	285	318	281	281	289	298	283	285	279	300	288	294	275	312	294	283	280	283	307	283	275	286	289	283	293	295	287	306	274
Jul	251	303	294	274	289	283	311	289	294	324	274	282	290	271	278	299	281	287	302	278	288	285	294	322	270	306	281	278	273	292	311	298	280	293	292	260	274
Aug	273	292	287	293	260	287	290	320	63	287	287	303	288	283	303	297	267	277	311	292	277	276	280	271	277	276	262	160	283	273	278	250	281	305	307	284	268
Sep	277	293	305	285	276	318	279	316	337	292	241	64	48	349	317	277	271	272	286	295	269	281	292	263	276	278	314	286	270	250	271	284	286	297	314	260	287
Oct	272	282	260	276	320	331	282	269	266	307	269	281	228	235	341	300	264	247	299	290	263	314	250	274	279	248	219	12	271	292	299	267	271	270	286	300	276
Nov	299	316	325	293	267	68	281	328	281	92	350	267	208	299	26	244	301	253	273	308	289	227	347	253	254	281	93	265	301	275	92	51	261	285	85	342	277
Dec	293	282	262	287	293	268	289	22	297	289	280	268	252	319	72	290	269	330	260	252	294	13	340	263	329	263	260	251	276	272	300	260	85	338	254	285	271

P12																																						
	H _{max} average [m]																																					
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	0.77	0.78	0.82	0.75	0.75	0.78	0.81	0.78	0.66	0.71	0.81	0.75	0.76	0.79	0.71	0.73	0.68	0.76	0.74	0.7	0.79	0.79	0.73	0.68	0.8	0.79	0.72	0.76	0.86	0.73	0.82	0.72	0.77	0.82	0.69	0.74	0.77	0.74
Jan	0.94	1.2	1.24	0.97	0.94	1.45	1.05	1.09	0.87	0.98	0.81	0.93	1.2	0.77	1.03	0.94	0.87	1.25	0.79	1.38	1.34	0.82	0.77	0.73	1	0.8	1.32	1.17	0.95	0.81	0.91	0.93	1.1	0.77	1.14	0.97	1.09	
Feb	1.08	1.24	1.37	0.87	0.92	0.77	0.73	1.24	0.81	1.3	1.6	0.85	1.08	0.87	1.23	0.48	0.74	0.94	0.8	0.98	1.3	0.83	0.99	1.07	0.84	0.92	0.99	0.9	0.86	0.91	0.82	0.87	1.25	0.78	1.5	0.91	0.94	
Mar	0.78	0.94	1.6	0.64	0.87	0.73	1.09	1.5	0.6	0.87	0.83	0.7	0.83	0.8	0.87	0.89	0.76	0.6	0.54	0.78	0.99	0.68	0.61	0.77	0.8	0.73	0.78	0.75	0.48	0.76	0.64	0.94	0.85	0.71	0.51	0.76	0.69	
Apr	0.6	0.68	0.7	0.63	0.63	0.65	0.64	0.74	0.36	0.84	0.98	0.68	0.49	0.96	0.45	0.67	0.52	0.59	0.65	0.37	0.42	0.65	0.96	0.6	0.65	0.57	0.7	0.65	0.8	0.62	0.5	0.7	0.68	0.53	0.56	0.64		
May	0.41	0.38	0.42	0.35	0.49	0.46	0.51	0.57	0.36	0.36	0.33	0.38	0.38	0.32	0.32	0.32	0.85	0.53	0.69	0.58	0.27	0.81	0.58	0.68	0.59	0.53	0.42	0.82	0.46	0.53	0.35	0.71	0.66	0.48	0.81	0.54	0.48	
Jun	0.33	0.42	0.41	0.69	0.42	0.6	0.84	0.59	0.45	0.52	0.67	0.59	0.7	0.58	0.73	0.68	0.71	0.62	0.48	0.43	0.74	0.59	0.46	0.57	0.7	0.69	0.45	0.74	0.54	0.86	0.57	0.56	0.45	0.41	0.48	0.35	0.68	
Jul	0.64	0.53	0.73	0.46	0.57	0.72	0.3	0.5	0.87	0.39	0.71	0.46	0.45	0.45	0.99	0.43	0.51	0.49	0.3	0.78	0.45	0.62	0.51	0.53	0.58	0.48	0.4	0.73	0.58	0.63	0.4	0.59	0.88	0.5	0.71	0.82	0.63	
Aug	0.71	0.75	0.61	0.62	0.82	0.65	0.62	0.5	0.99	0.24	0.82	0.41	0.64	0.71	0.44	0.94	0.96	0.62	0.4	0.68	0.74	0.75	0.65	0.69	0.48	0.59	0.87	0.64	0.76	0.69	0.69	0.89	0.42	0.72	0.36	0.81	0.67	
Sep	0.86	0.53	0.8	0.74	0.64	0.76	0.89	0.78	0.61	0.92	0.61	0.71	0.81	0.69	0.35	0.69	0.91	0.64	0.8	0.97	0.65	0.88	0.8	0.89	1.09	0.58	0.59	0.81	0.54	0.63	1	0.96	0.73	0.56	0.72	0.7	0.57	
Oct	0.9	0.9	0.85	0.77	0.74	0.68	0.94	0.79	0.84	0.85	1.3	0.99	0.92	0.98	0.88	0.73	0.95	0.76	0.62	0.55	1.3	0.9	0.85	1.02	0.89	0.93	0.99	0.75	0.9	1.2	0.98	0.74	0.87	0.97	0.83	1.2	0.79	
Nov	0.84	0.81	0.87	1.08	1.06	0.76	0.97	0.88	1.5	0.77	0.71	0.89	1.04	1.04	0.77	0.82	0.92	0.96	1.7	0.97	1.05	1.4	0.95	0.78	0.96	0.89	0.93	1.02	1.05	0.96	0.74	0.78	1.01	0.87	1.07	0.84	0.77	
Dec	1.5	0.87	0.97	0.97	0.89	1.6	1.1	0.8	0.68	0.7	0.95	1.24	0.84	0.79	0.77	1.07	0.97	0.95	1.21	0.78	0.64	0.73	0.74	1.32	0.94	1.24	1.01	1.27	0.96	1.6	0.91	1.03	0.93	0.82	0.82	1	0.82	

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	188.2 °	29.5 °	0.7 °	3.3 °	0.68	0.04	-0.01	-0.04
<i>Jan</i>	201.7 °	45.5 °	-0.6 °	-2.2 °	0.87	0.18	-0.01	-0.02
<i>Feb</i>	186.9 °	68.9 °	-0.0 °	-0.2 °	0.85	0.16	-0.01	-0.03
<i>Mar</i>	173.2 °	59.6 °	-0.1 °	-0.3 °	0.73	0.14	0.00	-0.01
<i>Apr</i>	112.7 °	73.5 °	0.5 °	0.7 °	0.62	0.09	-0.03	-0.10
<i>May</i>	116.9 °	74.6 °	0.3 °	5.3 °	0.55	0.09	-0.01	-0.04
<i>Jun</i>	206.9 °	72.4 °	0.8 °	4.4 °	0.49	0.12	-0.02	-0.05
<i>Jul</i>	214.2 °	58.6 °	1.4 °	2.2 °	0.50	0.09	-0.01	-0.03
<i>Aug</i>	202.0 °	55.7 °	1.5 °	3.6 °	0.52	0.09	-0.02	-0.02
<i>Sep</i>	187.4 °	64.3 °	0.6 °	0.7 °	0.67	0.11	-0.01	-0.03
<i>Oct</i>	189.2 °	49.3 °	-0.0 °	0.9 °	0.79	0.15	-0.01	-0.03
<i>Nov</i>	187.8 °	56.8 °	1.0 °	2.7 °	0.80	0.10	0.01	-0.04
<i>Dec</i>	203.0 °	47.4 °	0.1 °	0.0 °	0.85	0.12	0.00	-0.02

P13																																						
	Direction, average [°N]																																					
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	202	217	219	209	208	186	188	203	155	203	210	202	200	182	155	185	204	187	152	224	202	155	177	205	212	189	184	206	179	222	134	205	212	155	208	156	189	
Jan	180	238	223	219	241	227	226	187	185	211	222	210	247	185	233	211	153	241	184	241	189	185	180	215	217	188	151	220	154	218	180	258	227	142	248	211	209	
Feb	201	225	218	185	248	262	186	232	182	228	239	252	230	185	228	155	242	184	109	107	235	185	184	184	244	172	187	202	223	149	104	224	229	189	234	250	213	
Mar	185	218	244	150	222	237	225	213	185	222	225	144	242	186	229	184	149	184	110	142	228	158	240	214	245	181	223	151	144	225	186	231	184	231	188	227	103	
Apr	218	188	100	180	208	186	186	118	118	188	272	100	232	103	184	193	182	197	185	218	268	185	184	149	109	109	148	184	257	208	259	186	186	234	264	158	184	188
May	187	204	275	281	142	184	188	170	172	183	177	182	141	258	188	208	158	182	187	231	183	184	181	179	117	180	188	221	179	184	189	234	262	210	218	184	185	
Jun	178	208	155	239	181	261	251	250	255	181	234	202	257	265	203	223	231	257	210	181	220	187	280	235	245	249	273	245	182	238	248	182	182	181	218	184	235	
Jul	200	258	248	223	183	203	182	188	200	188	244	201	257	218	180	209	233	237	185	247	181	230	183	229	235	284	182	235	247	187	189	261	245	184	253	227	238	
Aug	241	275	231	252	225	243	219	309	189	189	251	182	245	218	182	265	189	236	244	219	230	204	225	207	228	188	224	186	239	226	233	207	185	241	185	237	214	
Sep	234	188	250	236	178	183	214	182	178	257	181	188	183	185	185	183	234	184	183	242	183	235	189	225	235	181	186	181	188	185	238	241	235	182	185	187	188	
Oct	180	239	170	183	180	184	181	228	207	268	203	205	180	183	182	249	180	186	210	180	225	180	189	218	224	186	181	186	186	186	186	186	186	186	186	186		
Nov	250	184	189	209	223	185	207	184	204	180	189	187	177	254	184	184	256	189	234	264	217	184	184	187	189	238	185	229	218	233	187	180	223	230	186	184	233	
Dec	244	187	188	248	182	204	224	187	183	185	227	225	204	183	189	231	233	180	226	218	187	189	189	227	149	226	221	212	240	230	214	217	182	187	184	213	218	

P13																																					
	H _{max} , average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	0.69	0.64	0.71	0.64	0.69	0.75	0.73	0.69	0.69	0.64	0.74	0.65	0.67	0.63	0.75	0.62	0.69	0.67	0.62	0.71	0.74	0.7	0.69	0.7	0.67	0.69	0.74	0.73	0.66	0.69	0.68	0.69	0.68	0.65	0.63	0.71	0.67
Jan	0.8	0.78	0.98	0.81	0.75	1.17	0.94	0.96	0.9	0.86	0.76	0.77	0.86	0.83	0.84	0.86	0.79	1.04	0.73	1.22	1.1	0.67	1.08	0.55	0.94	0.93	1.31	0.94	0.93	0.66	0.84	0.79	0.86	0.8	0.88	0.95	1.01
Feb	0.98	0.93	1.07	0.93	0.7	0.61	0.91	0.99	0.91	1.01	0.95	0.67	0.87	0.77	1.06	0.46	0.68	0.99	0.59	0.94	0.85	0.77	0.75	1.15	0.89	0.72	0.74	0.71	0.74	0.93	0.73	0.71	1.09	0.91	1.01	0.74	0.86
Mar	0.73	0.71	0.9	0.63	0.74	0.66	0.91	0.97	0.84	0.66	0.72	0.57	0.84	0.83	0.74	0.86	0.78	0.77	0.73	0.84	0.84	0.63	0.94	0.66	0.6	1.16	0.6	0.78	0.48	0.58	0.94	0.81	0.71	0.95	0.52	0.64	0.73
Apr	0.52	0.75	0.64	0.98	0.98	0.75	0.6	0.73	0.42	0.65	0.73	0.62	0.48	0.52	0.98	0.78	0.6	0.65	0.52	0.54	0.49	0.6	0.54	0.64	0.59	0.58	0.75	0.51	0.58	0.61	0.71	0.76	0.51	0.62	0.69	0.73	0.61
May	0.62	0.42	0.33	0.42	0.39	0.85	0.89	0.93	0.98	0.83	0.81	0.52	0.54	0.43	0.57	0.44	0.6	0.41	0.67	0.45	0.43	0.6	0.58	0.6	0.81	0.54	0.59	0.57	0.58	0.62	0.54	0.63	0.49	0.53	0.47	0.6	0.69
Jun	0.36	0.26	0.38	0.53	0.64	0.52	0.6	0.43	0.41	0.58	0.49	0.38	0.58	0.36	0.72	0.6	0.65	0.48	0.35	0.58	0.64	0.71	0.37	0.45	0.52	0.48	0.36	0.48	0.52	0.65	0.48	0.52	0.58	0.34	0.33	0.4	0.47
Jul	0.47	0.59	0.57	0.45	0.54	0.95	0.36	0.49	0.49	0.46	0.63	0.46	0.44	0.39	0.63	0.38	0.48	0.47	0.3	0.63	0.62	0.48	0.4	0.96	0.5	0.37	0.62	0.61	0.47	0.47	0.36	0.58	0.58	0.51	0.56	0.63	0.43
Aug	0.47	0.59	0.42	0.48	0.63	0.51	0.98	0.43	0.65	0.35	0.62	0.51	0.42	0.6	0.98	0.47	0.58	0.53	0.45	0.57	0.63	0.57	0.5	0.66	0.42	0.5	0.61	0.69	0.53	0.49	0.5	0.41	0.43	0.63	0.3	0.49	0.42
Sep	0.69	0.49	0.74	0.49	0.85	0.83	0.74	0.8	0.84	0.64	0.65	0.64	0.81	0.64	0.6	0.57	0.7	0.5	0.64	0.78	0.81	0.71	0.84	0.62	0.7	0.54	0.75	0.69	0.54	0.71	0.7	0.74	0.5	0.6	0.67	0.46	0.66
Oct	0.87	0.78	0.82	0.77	0.86	0.71	0.78	0.5	0.94	0.73	1.33	0.88	0.74	0.8	0.89	0.69	0.67	0.65	0.71	0.57	0.83	0.84	0.63	0.78	0.7	0.86	0.7	0.82	1.05	0.84	0.79	0.68	0.68	0.75	0.59	1	0.63
Nov	0.68	0.63	0.7	0.85	0.89	0.95	0.86	0.77	0.98	0.73	0.67	0.73	0.78	0.82	0.79	0.74	0.77	0.73	0.86	0.79	0.98	1.01	0.96	0.68	0.72	0.72	0.89	0.89	0.77	0.77	0.78	0.8	0.76	0.74	0.93	0.83	0.64
Dec	1.05	0.79	0.93	0.77	0.73	1.07	0.87	0.75	0.69	0.83	0.76	1.04	0.68	0.78	0.99	0.91	0.77	0.85	0.94	0.69	0.63	0.81	0.91	1.06	0.87	0.9	0.97	1.05	0.73	0.93	0.79	0.84	0.82	0.78	0.69	0.98	0.88

P14

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	184.5 °	37.6 °	-0.5 °	0.2 °	0.58	0.03	-0.01	-0.01
<i>Jan</i>	200.6 °	50.9 °	-2.6 °	1.8 °	0.73	0.15	-0.01	-0.01
<i>Feb</i>	182.7 °	81.6 °	-0.9 °	-0.7 °	0.72	0.13	-0.01	-0.01
<i>Mar</i>	163.6 °	73.2 °	-0.1 °	-0.2 °	0.62	0.13	-0.01	-0.01
<i>Apr</i>	73.1 °	79.1 °	0.0 °	0.7 °	0.54	0.09	-0.01	-0.02
<i>May</i>	76.4 °	78.1 °	-0.6 °	-0.5 °	0.48	0.08	-0.01	-0.02
<i>Jun</i>	219.3 °	86.7 °	-0.3 °	3.7 °	0.42	0.10	-0.02	-0.03
<i>Jul</i>	221.2 °	86.5 °	0.2 °	-1.1 °	0.43	0.08	-0.01	-0.02
<i>Aug</i>	202.4 °	64.6 °	-0.2 °	-1.2 °	0.44	0.08	-0.01	0.00
<i>Sep</i>	182.0 °	77.3 °	0.1 °	-0.0 °	0.56	0.09	0.00	-0.01
<i>Oct</i>	187.0 °	59.4 °	-0.3 °	-0.1 °	0.66	0.13	-0.01	-0.01
<i>Nov</i>	185.4 °	67.2 °	0.9 °	-0.1 °	0.67	0.09	0.01	-0.01
<i>Dec</i>	201.1 °	53.5 °	-0.5 °	-0.5 °	0.71	0.10	0.00	-0.01

P14																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	201	216	220	206	205	146	166	167	90	166	210	202	200	172	174	179	200	172	87	227	166	158	140	205	210	150	154	206	164	224	110	208	214	166	210	163	167
Jan	166	237	222	216	240	227	228	204	87	211	221	209	249	163	232	216	162	240	160	241	169	161	162	212	214	164	162	231	168	212	161	280	227	167	250	210	208
Feb	201	225	218	83	250	271	89	232	82	225	239	253	231	16	226	160	242	65	98	96	235	67	102	102	322	65	161	201	222	169	164	226	229	169	234	256	218
Mar	169	219	245	163	229	243	228	216	78	222	227	160	263	86	232	160	169	95	95	102	232	86	249	202	261	76	227	163	167	225	160	230	167	231	165	229	160
Apr	229	79	79	66	214	85	98	22	64	284	91	239	89	139	85	73	87	83	222	262	76	83	108	64	83	125	73	260	217	268	83	78	267	295	111	75	165
May	79	160	282	288	119	76	77	61	58	78	69	80	127	274	78	212	44	166	145	246	77	95	52	163	86	79	168	224	68	74	81	257	280	217	229	83	77
Jun	70	201	169	244	74	271	257	239	258	83	244	209	273	268	167	235	234	269	210	83	161	162	346	239	250	250	278	244	81	246	285	160	81	74	163	62	236
Jul	201	348	247	235	83	296	83	69	296	63	247	163	267	218	176	214	237	240	77	252	118	231	219	250	236	337	73	236	252	74	48	288	244	169	268	227	231
Aug	244	295	239	254	226	248	268	167	161	87	254	160	234	222	74	275	163	239	245	201	230	219	223	214	236	163	222	166	241	229	234	219	168	269	67	246	229
Sep	235	160	251	240	173	75	211	168	73	257	162	165	162	84	65	168	235	162	162	252	83	238	165	227	234	164	166	168	162	162	239	247	237	163	111	162	160
Oct	167	239	165	160	169	162	167	233	212	271	201	206	164	169	168	310	161	160	212	79	223	163	160	219	225	162	163	166	76	294	233	216	163	225	229	161	160
Nov	256	83	163	208	224	166	206	38	202	166	72	168	177	216	163	146	265	169	238	278	223	168	78	162	202	237	164	230	220	232	169	119	226	234	168	168	233
Dec	248	160	175	249	162	206	226	160	160	110	228	226	212	166	160	231	232	168	216	216	171	166	166	226	169	226	220	210	238	228	229	217	161	163	165	218	221

P14																																						
	H _{ms} , average [m]																																					
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	0.98	0.54	0.99	0.54	0.98	0.64	0.62	0.99	0.99	0.95	0.63	0.95	0.96	0.93	0.64	0.93	0.98	0.57	0.53	0.61	0.62	0.99	0.99	0.99	0.98	0.99	0.63	0.66	0.98	0.57	0.98	0.99	0.99	0.98	0.98	0.55	0.61	0.57
Jan	0.66	0.63	0.81	0.68	0.63	0.97	0.78	0.79	0.77	0.37	0.66	0.65	0.73	0.5	0.7	0.73	0.66	0.68	0.61	104	0.91	0.55	0.94	0.45	0.79	0.61	108	0.76	0.76	0.54	0.7	0.69	0.72	0.72	0.75	0.81	0.88	
Feb	0.85	0.76	0.89	0.8	0.99	0.51	0.79	0.82	0.79	0.84	0.78	0.56	0.73	0.66	0.89	0.38	0.58	0.87	0.51	0.78	0.7	0.66	0.65	0.98	0.76	0.65	0.6	0.59	0.63	0.78	0.63	0.6	0.94	0.82	0.87	0.63	0.74	
Mar	0.62	0.39	0.74	0.52	0.63	0.57	0.76	0.61	0.74	0.56	0.61	0.47	0.73	0.73	0.61	0.48	0.67	0.68	0.65	0.73	0.71	0.54	0.46	0.56	0.5	104	0.5	0.67	0.41	0.48	0.63	0.69	0.62	0.47	0.45	0.53	0.64	
Apr	0.45	0.66	0.96	0.5	0.48	0.67	0.53	0.63	0.37	0.96	0.63	0.52	0.41	0.45	0.53	0.7	0.52	0.58	0.44	0.46	0.43	0.53	0.47	0.56	0.51	0.49	0.66	0.42	0.48	0.52	0.61	0.7	0.44	0.56	0.63	0.67	0.53	
May	0.57	0.36	0.28	0.37	0.32	0.39	0.61	0.46	0.5	0.54	0.54	0.46	0.47	0.37	0.5	0.36	0.51	0.35	0.58	0.38	0.39	0.31	0.31	0.31	0.53	0.47	0.53	0.47	0.52	0.55	0.49	0.56	0.42	0.46	0.4	0.53	0.64	
Jun	0.32	0.23	0.31	0.45	0.50	0.44	0.49	0.37	0.34	0.51	0.41	0.32	0.47	0.31	0.62	0.49	0.54	0.41	0.29	0.49	0.54	0.63	0.31	0.38	0.43	0.4	0.31	0.37	0.44	0.54	0.41	0.45	0.53	0.3	0.29	0.36	0.4	
Jul	0.38	0.53	0.48	0.39	0.46	0.45	0.32	0.42	0.4	0.42	0.53	0.4	0.38	0.33	0.54	0.33	0.4	0.4	0.26	0.52	0.56	0.4	0.32	0.5	0.42	0.31	0.57	0.51	0.39	0.4	0.32	0.48	0.46	0.46	0.47	0.53	0.36	
Aug	0.38	0.49	0.35	0.4	0.52	0.43	0.48	0.37	0.56	0.3	0.51	0.44	0.33	0.5	0.51	0.39	0.48	0.44	0.4	0.48	0.52	0.47	0.41	0.36	0.35	0.41	0.5	0.59	0.42	0.4	0.4	0.34	0.39	0.58	0.28	0.41	0.34	
Sep	0.58	0.42	0.63	0.41	0.54	0.72	0.62	0.68	0.75	0.53	0.56	0.54	0.67	0.54	0.53	0.47	0.58	0.41	0.53	0.66	0.7	0.59	0.72	0.49	0.56	0.46	0.65	0.57	0.45	0.6	0.57	0.62	0.43	0.52	0.58	0.38	0.56	
Oct	0.74	0.65	0.69	0.66	0.75	0.61	0.63	0.39	0.44	0.63	1.2	0.74	0.61	0.66	0.76	0.59	0.72	0.53	0.59	0.45	0.68	0.73	0.68	0.64	0.58	0.72	0.57	0.67	0.6	0.69	0.65	0.59	0.76	0.63	0.49	0.66	0.52	
Nov	0.57	0.53	0.6	0.69	0.73	0.81	0.7	0.67	0.82	0.61	0.56	0.62	0.63	0.69	0.67	0.61	0.64	0.59	0.71	0.67	0.83	0.82	0.83	0.56	0.58	0.6	0.73	0.74	0.64	0.65	0.68	0.63	0.62	0.79	0.73	0.54		
Dec	0.88	0.65	0.78	0.64	0.59	0.8	0.72	0.64	0.58	0.73	0.64	0.87	0.57	0.65	0.85	0.76	0.64	0.72	0.77	0.58	0.52	0.68	0.8	0.88	0.73	0.75	0.81	0.86	0.59	0.76	0.66	0.71	0.66	0.67	0.58	0.84	0.72	

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	177.7°	18.8°	1.2°	6.8°	0.59	0.03	-0.01	-0.05
<i>Jan</i>	187.1°	39.8°	-1.4°	1.4°	0.77	0.17	-0.01	-0.06
<i>Feb</i>	177.0°	54.8°	-0.3°	-0.8°	0.73	0.15	-0.01	-0.04
<i>Mar</i>	169.5°	46.3°	-0.0°	-0.5°	0.63	0.12	0.00	-0.03
<i>Apr</i>	145.8°	58.4°	-0.6°	-2.1°	0.53	0.08	-0.02	-0.08
<i>May</i>	149.0°	58.0°	2.7°	9.2°	0.47	0.08	-0.01	-0.04
<i>Jun</i>	189.1°	63.5°	-1.9°	-0.7°	0.42	0.10	-0.02	-0.07
<i>Jul</i>	196.8°	55.4°	5.2°	11.7°	0.41	0.07	-0.01	-0.04
<i>Aug</i>	186.6°	48.9°	1.5°	7.1°	0.44	0.09	-0.02	-0.05
<i>Sep</i>	176.4°	47.8°	1.7°	4.5°	0.58	0.09	0.00	-0.03
<i>Oct</i>	176.8°	34.6°	-0.6°	1.9°	0.70	0.13	-0.01	-0.04
<i>Nov</i>	175.4°	41.0°	0.2°	-1.3°	0.72	0.11	-0.01	-0.09
<i>Dec</i>	187.6°	38.7°	0.7°	2.6°	0.74	0.11	-0.01	-0.06

P15																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	50	58	203	57	58	50	79	50	54	55	57	54	77	54	45	71	54	71	73	59	55	54	50	57	59	53	45	55	55	59	42	54	55	75	55	77	55
Jan	57	239	26	208	239	221	24	78	70	20	207	58	244	50	227	201	52	237	59	240	53	73	55	56	50	52	59	28	51	55	55	249	28	52	249	55	79
Feb	59	215	207	18	246	259	12	226	10	28	239	254	221	50	219	149	233	51	51	50	231	15	57	143	221	53	50	57	210	148	58	205	222	54	228	254	53
Mar	54	205	248	51	207	238	24	58	55	208	58	145	211	10	222	145	59	57	55	148	201	51	230	58	249	56	210	51	143	206	52	224	55	219	51	208	51
Apr	52	57	55	53	72	16	54	52	84	274	114	230	52	55	15	55	12	19	57	281	57	111	146	141	54	58	56	263	58	264	114	104	237	256	52	105	59
May	55	215	294	290	56	52	58	50	54	18	57	10	50	239	55	55	50	51	51	201	51	52	50	55	54	10	55	205	58	105	109	215	267	57	55	111	51
Jun	58	52	51	225	59	264	258	231	262	16	222	53	251	267	57	204	221	257	56	52	209	59	252	211	242	241	280	234	54	224	233	58	114	52	149	55	216
Jul	79	205	242	59	18	257	59	54	230	52	237	201	230	56	70	51	220	210	18	242	59	210	111	55	227	312	54	225	244	145	53	268	245	72	248	215	208
Aug	230	272	205	239	209	232	54	337	16	55	245	54	77	53	53	259	56	222	201	59	220	53	202	79	220	51	21	12	232	207	206	57	142	245	10	236	58
Sep	228	53	234	214	57	53	56	54	55	260	56	59	19	19	58	58	225	50	145	215	50	53	214	224	55	55	57	141	51	226	230	224	143	59	71	149	
Oct	56	226	54	53	149	51	72	211	79	263	53	55	58	50	17	50	59	57	55	145	211	58	52	54	201	56	55	10	52	238	54	202	51	203	207	149	59
Nov	234	75	149	52	209	10	58	141	58	52	15	53	53	247	51	52	235	52	217	245	52	54	59	55	79	220	56	220	56	221	59	55	211	219	145	52	215
Dec	241	70	70	246	76	56	209	54	111	59	212	211	56	57	15	212	223	53	208	200	53	52	52	214	147	211	207	200	237	217	52	203	50	149	50	201	200

P16																																						
		Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	
Annual	0.6	0.56	0.61	0.55	0.59	0.64	0.65	0.61	0.59	0.55	0.63	0.56	0.56	0.55	0.64	0.54	0.56	0.57	0.54	0.6	0.64	0.61	0.59	0.61	0.59	0.59	0.67	0.65	0.57	0.59	0.6	0.5	0.5	0.5	0.54	0.56	0.61	0.59
Jan	0.79	0.66	0.67	0.71	0.63	103	0.8	0.88	0.85	0.38	0.6	0.66	0.72	0.63	0.71	0.73	0.75	0.85	0.7	1	106	0.61	0.93	0.49	0.6	0.76	131	0.84	0.86	0.54	0.79	0.66	0.73	0.64	0.75	0.63	0.83	0.88
Feb	0.86	0.84	0.98	0.81	0.57	0.5	0.83	0.87	0.8	0.9	0.77	0.54	0.75	0.66	0.9	0.43	0.54	0.81	0.48	0.92	0.74	0.65	0.66	104	0.71	0.51	0.76	0.67	0.63	0.84	0.62	0.62	0.92	0.77	0.85	0.61	0.75	
Mar	0.63	0.65	0.75	0.59	0.63	0.52	0.81	0.89	0.73	0.59	0.63	0.54	0.68	0.73	0.63	0.5	0.65	0.59	0.58	0.7	0.75	0.57	0.44	0.58	0.54	0.56	0.55	0.67	0.42	0.51	0.64	0.69	0.63	0.47	0.48	0.56	0.67	
Apr	0.65	0.68	0.54	0.49	0.52	0.64	0.53	0.6	0.33	0.55	0.66	0.52	0.44	0.45	0.47	0.65	0.55	0.54	0.45	0.45	0.42	0.53	0.47	0.53	0.52	0.52	0.63	0.43	0.52	0.51	0.63	0.64	0.47	0.49	0.52	0.58	0.53	
May	0.52	0.36	0.26	0.33	0.48	0.54	0.57	0.45	0.46	0.54	0.49	0.47	0.47	0.35	0.49	0.42	0.52	0.36	0.35	0.39	0.34	0.49	0.45	0.52	0.53	0.47	0.47	0.52	0.49	0.53	0.44	0.53	0.41	0.46	0.42	0.54	0.8	
Jun	0.26	0.24	0.34	0.45	0.52	0.41	0.52	0.38	0.32	0.5	0.43	0.37	0.47	0.29	0.61	0.52	0.55	0.4	0.3	0.44	0.53	0.56	0.3	0.39	0.45	0.41	0.29	0.41	0.46	0.57	0.4	0.48	0.48	0.3	0.29	0.33	0.42	
Jul	0.44	0.48	0.48	0.35	0.47	0.45	0.3	0.41	0.42	0.35	0.51	0.37	0.32	0.33	0.52	0.31	0.39	0.38	0.26	0.51	0.48	0.43	0.37	0.47	0.41	0.31	0.49	0.51	0.37	0.4	0.27	0.46	0.5	0.41	0.48	0.54	0.38	
Aug	0.4	0.48	0.37	0.39	0.53	0.41	0.49	0.34	0.51	0.31	0.51	0.42	0.36	0.54	0.51	0.37	0.47	0.42	0.35	0.49	0.58	0.51	0.43	0.57	0.36	0.43	0.53	0.54	0.45	0.44	0.46	0.38	0.36	0.49	0.24	0.41	0.4	
Sep	0.59	0.4	0.59	0.42	0.55	0.76	0.55	0.59	0.69	0.67	0.55	0.59	0.6	0.78	0.56	0.48	0.5	0.58	0.45	0.59	0.67	0.65	0.5	0.68	0.55	0.64	0.47	0.62	0.63	0.47	0.5	0.62	0.64	0.43	0.49	0.62	0.45	0.57
Oct	0.79	0.64	0.75	0.68	0.7	0.64	0.74	0.47	0.51	0.56	1.0	0.76	0.73	0.74	0.75	0.53	0.79	0.63	0.62	0.45	0.72	0.7	0.79	0.69	0.63	0.8	0.71	0.78	0.68	0.71	0.71	0.58	0.77	0.69	0.55	0.68	0.62	
Nov	0.57	0.56	0.56	0.79	0.79	0.89	0.78	0.65	0.86	0.71	0.6	0.6	0.82	0.86	0.75	0.72	0.64	0.68	0.76	0.65	0.86	0.99	0.77	0.65	0.68	0.5	0.89	0.76	0.69	0.65	0.75	0.77	0.68	0.62	0.9	0.74	0.54	
Dec	0.88	0.69	0.83	0.63	0.7	0.95	0.77	0.65	0.57	0.7	0.66	0.9	0.61	0.62	0.9	0.79	0.63	0.71	0.63	0.61	0.56	0.72	0.72	0.92	0.8	0.64	0.81	0.56	0.61	0.62	0.69	0.75	0.87	0.69	0.62	0.84	0.73	

	AENWS		NIRAS Model		AENWS		NIRAS Model	
	Mean MWD	St.Dev.	Impact on Direction		Mean Hm0	St. Dev.	Impact on Hm0	
	[°N]	[°]	Current Scenario - Baseline	Future Scenario - Baseline	[m]	[m]	Current Scenario - Baseline	Future Scenario - Baseline
<i>Annual</i>	243.2°	28.0°	-0.2°	-0.6°	0.66	0.03	0.00	-0.01
<i>Jan</i>	234.5°	55.2°	-0.5°	-0.8°	0.84	0.16	0.00	-0.02
<i>Feb</i>	239.5°	85.3°	-0.2°	-0.8°	0.81	0.16	0.00	-0.02
<i>Mar</i>	232.4°	67.5°	-0.4°	-0.4°	0.68	0.13	0.00	-0.01
<i>Apr</i>	332.5°	88.6°	0.8°	1.4°	0.55	0.10	-0.01	-0.02
<i>May</i>	303.2°	90.2°	-0.5°	-1.0°	0.49	0.08	0.00	-0.02
<i>Jun</i>	267.6°	36.0°	6.2°	8.6°	0.50	0.11	-0.01	-0.02
<i>Jul</i>	260.9°	30.6°	-0.4°	-1.8°	0.48	0.10	0.00	-0.01
<i>Aug</i>	248.3°	50.1°	-0.2°	-0.7°	0.52	0.11	0.00	-0.02
<i>Sep</i>	250.9°	70.0°	0.0°	-0.4°	0.64	0.12	0.00	-0.01
<i>Oct</i>	224.4°	61.9°	-0.6°	-1.7°	0.75	0.12	0.00	-0.01
<i>Nov</i>	227.0°	83.5°	-3.3°	-1.7°	0.79	0.10	0.00	0.00
<i>Dec</i>	236.4°	66.3°	0.3°	-0.1°	0.81	0.13	0.01	-0.01

P16																																					
	Direction, average [°N]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	250	296	250	254	232	262	245	277	42	270	255	240	235	276	248	235	252	257	236	268	244	244	139	239	257	251	89	242	254	258	239	255	243	281	230	253	236
Jan	80	246	231	230	272	239	245	231	91	251	262	229	263	83	244	250	81	257	83	258	217	210	72	233	267	329	81	241	89	266	203	322	239	350	277	227	255
Feb	234	235	226	57	269	284	79	244	60	235	249	281	240	329	240	87	271	28	64	94	249	5	72	93	283	6	89	267	243	84	16	246	240	64	248	281	240
Mar	290	231	260	145	205	282	239	238	36	253	273	86	291	93	270	82	262	6	79	247	260	273	277	258	288	66	247	222	82	245	72	247	240	253	146	252	91
Apr	308	61	269	291	268	125	334	333	11	307	94	263	50	264	108	29	96	87	238	298	72	107	281	285	284	225	0	283	251	281	72	67	288	315	293	33	214
May	99	281	314	329	238	21	59	328	314	85	309	140	227	301	87	225	312	244	239	265	53	265	344	228	263	65	358	241	23	31	104	281	294	226	252	26	85
Jun	349	275	225	261	26	284	264	297	267	72	255	232	266	287	258	264	257	290	275	278	253	319	287	260	260	258	306	263	289	267	279	235	238	270	259	315	248
Jul	212	302	278	230	276	265	9	274	278	348	257	281	274	242	262	295	281	268	297	298	236	241	264	338	248	297	235	258	252	277	302	286	259	279	278	238	244
Aug	249	279	263	278	232	272	277	330	82	10	269	334	271	259	70	287	239	255	321	278	252	238	254	240	251	248	234	89	261	246	254	228	238	312	322	271	232
Sep	256	272	309	260	227	59	253	0	31	279	51	111	93	38	345	250	249	238	246	285	82	259	289	236	250	235	5	236	236	87	248	263	266	308	48	225	276
Oct	217	298	82	201	5	80	237	237	227	308	238	239	87	214	45	301	93	86	230	270	236	347	84	240	252	203	96	84	96	275	279	238	205	237	235	297	221
Nov	290	283	356	225	238	82	251	356	239	19	52	228	89	291	80	98	291	223	245	310	274	82	34	207	215	259	111	242	279	251	132	100	235	261	87	66	256
Dec	279	253	224	271	216	230	233	63	291	132	253	243	217	345	84	273	244	12	236	225	271	89	11	241	78	236	236	223	256	249	272	229	65	48	214	259	239

P16																																					
	Hm0, average [m]																																				
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024
Annual	0.67	0.66	0.71	0.66	0.65	0.69	0.7	0.69	0.62	0.63	0.71	0.64	0.66	0.63	0.66	0.58	0.66	0.65	0.6	0.7	0.67	0.64	0.64	0.67	0.68	0.65	0.66	0.72	0.64	0.7	0.63	0.65	0.67	0.61	0.62	0.65	0.62
Jan	0.72	0.67	0.57	0.79	0.81	1.17	0.89	0.92	0.64	0.5	0.72	0.8	0.96	0.61	0.85	0.83	0.72	107	0.66	1.9	107	0.66	0.78	0.61	0.89	0.76	109	0.93	0.83	0.7	0.73	0.85	0.89	0.75	0.96	0.8	0.95
Feb	0.97	0.98	109	0.87	0.79	0.7	0.72	101	0.82	105	0.99	0.75	0.91	0.8	104	0.4	0.88	0.95	0.56	0.84	0.93	0.63	0.58	0.91	0.77	0.58	0.75	0.79	0.73	0.81	0.75	0.71	103	0.73	0.96	0.79	0.78
Mar	0.73	0.73	0.86	0.55	0.77	0.65	0.9	0.97	0.66	0.72	0.74	0.57	0.74	0.56	0.75	0.51	0.72	0.61	0.55	0.72	0.86	0.62	0.56	0.65	0.7	0.79	0.65	0.66	0.39	0.64	0.67	0.79	0.72	0.61	0.4	0.65	0.59
Apr	0.54	0.65	0.62	0.59	0.56	0.5	0.6	0.67	0.34	0.81	0.53	0.61	0.43	0.53	0.43	0.69	0.47	0.46	0.5	0.58	0.33	0.37	0.57	0.51	0.56	0.57	0.56	0.61	0.57	0.68	0.57	0.49	0.57	0.65	0.51	0.54	0.57
May	0.4	0.49	0.37	0.54	0.43	0.46	0.51	0.52	0.54	0.52	0.53	0.51	0.47	0.5	0.48	0.42	0.61	0.45	0.62	0.51	0.27	0.35	0.39	0.58	0.51	0.49	0.4	0.67	0.46	0.51	0.35	0.63	0.57	0.43	0.5	0.47	0.46
Jun	0.33	0.35	0.38	0.61	0.42	0.56	0.69	0.54	0.42	0.48	0.54	0.47	0.58	0.5	0.66	0.58	0.62	0.57	0.42	0.4	0.62	0.6	0.41	0.47	0.6	0.6	0.43	0.59	0.49	0.72	0.51	0.45	0.41	0.34	0.37	0.32	0.58
Jul	0.51	0.52	0.64	0.42	0.48	0.61	0.28	0.43	0.57	0.39	0.62	0.39	0.4	0.37	0.52	0.38	0.45	0.42	0.25	0.64	0.43	0.5	0.42	0.5	0.49	0.42	0.39	0.61	0.5	0.53	0.36	0.52	0.68	0.45	0.58	0.63	0.49
Aug	0.57	0.65	0.51	0.54	0.66	0.57	0.55	0.48	0.53	0.21	0.69	0.4	0.51	0.38	0.4	0.49	0.48	0.52	0.38	0.5	0.65	0.57	0.54	0.58	0.4	0.51	0.71	0.54	0.63	0.57	0.55	0.46	0.33	0.68	0.32	0.46	0.5
Sep	0.71	0.46	0.76	0.62	0.53	0.78	0.75	0.72	0.68	0.75	0.54	0.57	0.72	0.68	0.54	0.56	0.75	0.5	0.64	0.87	0.58	0.73	0.73	0.68	0.86	0.51	0.58	0.7	0.44	0.55	0.62	0.76	0.58	0.48	0.63	0.52	0.51
Oct	0.78	0.8	0.71	0.69	0.73	0.6	0.77	0.61	0.65	0.83	1.2	0.83	0.7	0.78	0.83	0.67	0.62	0.57	0.66	0.5	0.67	0.86	0.71	0.83	0.73	0.78	0.74	0.67	0.93	0.93	0.65	0.63	0.73	0.76	0.64	0.96	0.62
Nov	0.77	0.71	0.67	0.87	0.87	0.73	0.8	0.84	0.96	0.66	0.67	0.76	0.79	0.92	0.7	0.65	0.81	0.77	0.97	0.91	0.94	0.92	0.97	0.63	0.77	0.74	0.79	0.86	0.89	0.79	0.62	0.66	0.78	0.72	0.87	0.78	0.64
Dec	104	0.75	0.83	0.83	0.69	0.98	0.89	0.81	0.62	0.66	0.83	102	0.7	0.75	0.74	0.93	0.78	0.9	0.94	0.63	0.56	0.7	0.8	105	0.9	0.98	0.83	101	0.81	0.95	0.79	0.83	0.7	0.71	0.71	0.9	0.78